

Management of Spontaneous Abortion

CRAIG P. GRIEBEL, M.D., JOHN HALVORSEN, M.D., THOMAS B. GOLEMOM, M.D., and ANTHONY A. DAY, M.D., *University of Illinois College of Medicine at Peoria, Peoria, Illinois*

Spontaneous abortion, which is the loss of a pregnancy without outside intervention before 20 weeks' gestation, affects up to 20 percent of recognized pregnancies. Spontaneous abortion can be subdivided into threatened abortion, inevitable abortion, incomplete abortion, missed abortion, septic abortion, complete abortion, and recurrent spontaneous abortion. Ultrasonography is helpful in the diagnosis of spontaneous abortion, but other testing may be needed if an ectopic pregnancy cannot be ruled out. Chromosomal abnormalities are causative in approximately 50 percent of spontaneous abortions; multiple other factors also may play a role. Traditional treatment consisting of surgical evacuation of the uterus remains the treatment of choice in unstable patients. Recent studies suggest that expectant or medical management is appropriate in selected patients. Patients with a completed spontaneous abortion rarely require medical or surgical intervention. For women with incomplete spontaneous abortion, expectant management for up to two weeks usually is successful, and medical therapy provides little additional benefit. When patients are allowed to choose between treatment options, a large percentage will choose expectant management. Expectant management of missed spontaneous abortion has variable success rates, but medical therapy with intravaginal misoprostol has an 80 percent success rate. Physicians should be aware of psychologic issues that patients and their partners face after completing a spontaneous abortion. Women are at increased risk for significant depression and anxiety for up to one year after spontaneous abortion. Counseling to address feelings of guilt, the grief process, and how to cope with friends and family should be provided. (*Am Fam Physician* 2005;72:1243-50. Copyright © 2005 American Academy of Family Physicians.)

Spontaneous abortion refers to pregnancy loss at less than 20 weeks' gestation in the absence of elective medical or surgical measures to terminate the pregnancy. The term "miscarriage" is synonymous and often is used with patients because the word "abortion" is associated

with elective termination. "Spontaneous pregnancy loss" has been recommended to avoid the term "abortion" and acknowledge the emotional aspects of losing a pregnancy.¹ Another emotionally neutral term is "early pregnancy failure."²

For clinical purposes, spontaneous abortion often is subdivided into threatened abortion, inevitable abortion, incomplete abortion, missed abortion, septic abortion, recurrent spontaneous abortion, and complete abortion (*Table 1*).

Incidence

Approximately 20 percent of pregnant women will have some bleeding before 20 weeks' gestation, and roughly one half of these pregnancies will end in spontaneous abortion.³ Up to 20 percent of recognized pregnancies will end in miscarriage. However, when women were followed with serial serum human chorionic gonadotropin (hCG) measurements, the actual miscarriage rate was found to be 31 percent.⁴ Many pregnancies are lost spontaneously before a woman recognizes that she is pregnant, and the clinical signs of miscarriage are mistaken for a heavy or late menses.

TABLE 1
Spontaneous Abortion: Definitions of Subcategories

Complete abortion: all products of conception have been passed without the need for surgical or medical intervention
Incomplete abortion: some, but not all, of the products of conception have been passed; retained products may be part of the fetus, placenta, or membranes
Inevitable abortion: the cervix has dilated, but the products of conception have not been expelled
Missed abortion: a pregnancy in which there is a fetal demise (usually for a number of weeks) but no uterine activity to expel the products of conception
Recurrent spontaneous abortion: three or more consecutive pregnancy losses
Septic abortion: a spontaneous abortion that is complicated by intrauterine infection
Threatened abortion: a pregnancy complicated by bleeding before 20 weeks' gestation

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
The possibility of ectopic pregnancy should be considered when transvaginal ultrasonography reveals an empty uterus and the quantitative serum human chorionic gonadotropin level is greater than 1,800 mIU per mL (1,800 IU per L).	C	5
Transvaginal ultrasound should be performed in the first trimester of pregnancy when incomplete abortion is suspected and is extremely reliable in identifying intrauterine products of conception.	C	7, 8
Expectant management should be considered for women with incomplete spontaneous abortions. It has an 82 to 96 percent success rate without the need for surgical or medical intervention.	A	17-22, 24
When misoprostol (Cytotec) is used to treat women with a missed spontaneous abortion, it should be given vaginally rather than orally.	B	27
Patients who have had a spontaneous abortion should be given the opportunity to choose a treatment option.	B	28
A 50-mcg dose of Rh ₀ (D) immune globulin (Rhogam) should be administered to Rh-negative patients who have a threatened abortion or have completed a spontaneous abortion.	C	5
Physicians should be alert to the development of psychologic symptoms that frequently occur following spontaneous abortion (e.g., depression, anxiety).	C	31-34

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, see page 1154 or <http://www.aafp.org/afpsort.xml>.

Diagnosis

Threatened abortion is defined by vaginal bleeding in a woman with a confirmed pregnancy. First-trimester bleeding in a pregnant woman has an extensive differential diagnosis (Table 2) and should be evaluated with a full history and physical examination. Laboratory tests should include potassium hydroxide and “wet prep” microscopy of any vaginal discharge, complete blood count, blood typing and Rh testing, and quantitative serum hCG testing. Gonorrhea and chlamydia testing also should be considered. Ultrasonography is crucial in identifying the status of the pregnancy and verifying that the pregnancy is intrauterine. When transvaginal ultrasonography reveals an empty uterus and the quantitative

serum hCG level is greater than 1,800 mIU per mL (1,800 IU per L), an ectopic pregnancy should be considered.⁵ When transabdominal ultrasonography is performed, an empty uterus should raise suspicion of an ectopic pregnancy if quantitative hCG levels are greater than 3,500 mIU per mL (3,500 IU per L). A uterus found to be empty on ultrasound examination may signal a completed spontaneous abortion, but the diagnosis is not definitive until ectopic pregnancy is excluded. If an ultrasound examination finds an intrauterine pregnancy, ectopic pregnancy is unlikely, although heterotopic pregnancy has been reported (i.e., simultaneous intrauterine and ectopic pregnancies).⁵ The risk for spontaneous abortion decreases from 50 to 3 percent when a fetal heartbeat is identified on ultrasound examination.¹

When the clinical examination reveals a dilated cervix, spontaneous abortion is inevitable. However, cervical evaluation is not reliable for distinguishing between complete and incomplete abortion.^{6,7} Transvaginal ultrasonography should be performed and is extremely reliable for finding products of conception, with a 90 to 100 percent sensitivity and 80 to 92 percent specificity.^{7,8}

A missed spontaneous abortion usually is diagnosed by routine ultrasonography or when an ultrasound scan is obtained because the symptoms and physical signs of pregnancy are regressing. Figure 1 presents an algorithm for diagnosing spontaneous abortion.¹

TABLE 2
Differential Diagnosis of First-Trimester Vaginal Bleeding

- Cervical abnormalities (e.g., excessive friability, malignancy, polyps, trauma)
- Ectopic pregnancy
- Idiopathic bleeding in a viable pregnancy
- Infection of the vagina or cervix
- Molar pregnancy
- Spontaneous abortion
- Subchorionic hemorrhage
- Vaginal trauma

Diagnosis of Spontaneous Abortion




Figure 1. Algorithm for the diagnosis of spontaneous pregnancy loss. (ED = emergency department; hCG = human chorionic gonadotropin.)

Adapted with permission from Scroggins KM, Smucker WD, Krishen AE. Spontaneous pregnancy loss: evaluation, management, and follow-up counseling. *Prim Care* 2000;27:157.

Spontaneous Abortion

The risk for spontaneous abortion decreases from 50 to 3 percent when a fetal heartbeat is identified on ultrasound examination.

Etiology and Risk Factors

Chromosomal abnormalities are a direct cause of spontaneous abortion. One meta-analysis⁹ found that a chromosomal abnormality occurs in 49 percent of spontaneous abortions.

Autosomal trisomy was the most commonly identified anomaly (52 percent), followed by polyploidy (21 percent) and monosomy X (13 percent).⁹ Most chromosomal abnormalities that result in spontaneous abortion are random events, such as maternal and paternal gametogenesis errors, dispermy, and nondisjunction. Structural abnormalities of individual chromosomes (e.g., translocations, inversions) were reported in 6 percent of women who had spontaneous abortions, and approximately one half of these abnormalities were inherited.⁹ Chromosomal abnormalities are more likely to be associated with recurrent spontaneous abortion, but are uncommon even in that instance (4 to 6 percent).⁹

Risk factors for spontaneous abortion are

listed in *Table 3*.^{1,10-14} However, other factors are notable for their lack of association with miscarriage. One study¹⁵ that examined the influence of stress on early pregnancy loss failed to find a clear association. Marijuana use, likewise, has not been proven to increase the risk for spontaneous abortion.¹¹ Sexual activity also does not elevate risk in women with uncomplicated pregnancies.

Treatment

Dilatation and curettage is the traditional treatment for spontaneous abortion; manual vacuum aspiration is another surgical option. Prompt surgical evacuation of the uterus has been recommended in the past because of the risk for infection and concerns about coagulation disorders that result from retained products of conception.^{1,2} However, the need for immediate surgical evacuation in all patients with a spontaneous abortion has been questioned. Many recent studies¹⁶⁻²⁴ have examined the outcomes of expectant and medical management for women with spontaneous abortions.

Prompt surgical evacuation of the uterus is the treatment of choice when the patient is unstable because of heavy bleeding or has evidence of a septic abortion. Patient choice is another reason to proceed with surgical evacuation.

Some women may have already completed a spontaneous abortion by the time they present for clinical evaluation. If the ultrasound examination shows an empty uterus and evaluation of the expelled tissue confirms the presence of products of conception, no further action is needed; in these instances, patients have a completed spontaneous abortion and can be managed expectantly.¹⁶ If the products of conception are not physically confirmed when the uterus is empty, an ectopic pregnancy must be ruled out.

Many studies¹⁷⁻²⁴ have compared expectant management, medical therapy, and surgical management for women with incomplete spontaneous abortion. Expectant management proved to be successful, with no need for surgical intervention in 82 to 96 percent of women.^{17-22,24} Most patients who had surgical intervention were followed expectantly

TABLE 3
Risk Factors for Spontaneous Abortion

Advanced maternal age
Alcohol use
Anesthetic gas use (e.g., nitrous oxide)
Caffeine use (heavy)
Chronic maternal diseases: poorly controlled diabetes, celiac disease, autoimmune diseases (particularly antiphospholipid antibody syndrome)
Cigarette smoking
Cocaine use
Conception within three to six months after delivery
Intrauterine device use
Maternal infections: bacterial vaginosis; mycoplasmosis, herpes simplex virus, toxoplasmosis, listeriosis, chlamydia, human immunodeficiency virus, syphilis, parvovirus B19, malaria, gonorrhea, rubella, cytomegalovirus
Medications: misoprostol (Cytotec), retinoids, methotrexate, nonsteroidal anti-inflammatory drugs
Multiple previous elective abortions
Previous spontaneous abortion
Toxins: arsenic, lead, ethylene glycol, carbon disulfide, polyurethane, heavy metals, organic solvents
Uterine abnormalities: congenital anomalies, adhesions, leiomyoma

Information from references 1 and 10 through 14.




Figure 2. Algorithm for the management of spontaneous pregnancy loss. (hCG = human chorionic gonadotropin.)

Adapted with permission from Scroggins KM, Smucker WD, Krishen AE. Spontaneous pregnancy loss: evaluation, management, and follow-up counseling. *Prim Care* 2000;27:161.

for two weeks before intervention was recommended.^{17,19,21} Medical therapy with misoprostol (Cytotec) or mifepristone (Mifeprex) does not confer significant additional benefit.²³ The average time to completion of the miscarriage was nine days.²⁰

In women with missed spontaneous abortions, expectant management has a variable but generally lower success rate than medical therapy, ranging from 16 to 76 percent.^{17,20,25,26} In contrast, medical therapy for missed spontaneous abortion results in high success rates for completion of a spontaneous abortion without surgical intervention. One study²⁵ found that patients had an 80 percent success rate after using 800 mcg of misoprostol, administered intravaginally and repeated after four hours, if necessary. Intravaginal administration of misoprostol causes less diarrhea than oral administration.²⁷

Patient preferences should be considered when choosing a treatment for spontane-

ous abortion. Physicians should discuss the available options and the evidence to support each option with the patient. There is evidence to suggest that women who are given the opportunity to choose a treatment option have better subsequent mental health than women who are not allowed to choose their therapy.²⁸ However, patients express less happiness with the mode of treatment they receive and are less willing to have the same care again when they begin with noninvasive management and later require surgical intervention.²⁹ When patients are allowed to choose their therapy, 38 to 75 percent choose expectant management.^{20,26,30}

An algorithm for managing women with spontaneous abortion is presented in *Figure 2*.¹ A 50-mcg dose of Rh₀(D)

Patients express less happiness with the mode of treatment they receive and are less willing to have the same care again when they begin with noninvasive management and later require surgical intervention.

Spontaneous Abortion

immune globulin (Rhogam) should be given to patients who are Rh-negative and have a threatened abortion or have completed a spontaneous abortion.⁵

Psychologic Issues After Spontaneous Abortion

Physicians should recognize the psychologic issues that affect a patient who experiences a spontaneous abortion. Although the literature lacks good evidence to support psychologic counseling for women after a spontaneous abortion, it is thought that patients will have better outcomes if these issues are addressed. The patient and her partner may be dealing with feelings of guilt, and they typically will go through a grieving process and have symptoms of anxiety and depression.

Women who have a spontaneous abortion frequently struggle with guilt over what they may have done to cause or prevent the loss. Physicians should address the issue of guilt

with their patients and allay any concerns that they may have “caused” the spontaneous abortion.

Physicians should encourage the patient and her partner to allow themselves to grieve. The woman and her partner may grieve differently; specifically, they may go through the stages of grief in different orders or at different rates. They also should be aware that friends and family members may not recognize the magnitude of their loss. Friends and family members may ignore the subject of miscarriage, or they may make well-meaning comments that try to minimize the event. Connecting the couple with a counselor who has experience in helping couples cope with pregnancy loss may be beneficial. Many hospitals offer programs that provide follow-up care and literature to the woman and her partner. Two national organizations, the Compassionate Friends (<http://www.compassionatefriends.org>; telephone: 877-969-0010) and SHARE Pregnancy and Infant Loss Support, Inc. (<http://www.nationalshareoffice.com>; telephone: 800-821-6819), provide support for women and their partners as they progress through the grieving process after a miscarriage.

Most studies³¹⁻³⁴ have found that a significant percentage of women experience psychiatric symptoms in the weeks to months after spontaneous abortion. Women who were found to be especially prone to these symptoms are childless and have lost a wanted pregnancy.³¹ One study²⁸ showed that women who are managed expectantly have better overall mental health 12 weeks after a spontaneous abortion.

Physicians should realize the importance of providing care that is sensitive to the medical and psychologic aspects of a couple who experiences spontaneous abortion. Many patients report dissatisfaction with the medical care they receive.^{35,36} The Advanced Life Support in Obstetrics⁵ provider course offered by the American Academy of Family Physicians summarizes issues to discuss with women and their partners after a spontaneous abortion (Table 4).⁵

The Authors

CRAIG P. GRIEBEL, M.D., is clinical assistant professor and director of obstetrics and gynecology at the University of Illinois College of Medicine at Peoria family medicine residency program at Methodist Medical Center. Dr. Griebel received his medical degree from the University of Missouri School of Medicine, Columbia, and completed his residency at Quad Cities Genesis Family Practice Residency Program in Davenport, Iowa.

JOHN HALVORSEN, M.D., is the Thomas and Ellen Foster chair and professor of family medicine at the University of Illinois College of Medicine at Peoria. He also is associate dean for community health. He received his medical degree from Ohio State University School of Medicine and Public Health, Columbus, and completed a family medicine residency at the University of Minnesota Medical School at Hennepin County Medical Center, Minneapolis.

THOMAS B. GOLEMON, M.D., is executive director of the University of Illinois College of Medicine at Peoria family medicine residency program at Methodist Medical Center. He received his medical degree from the University of Texas Southwestern Medical School, Dallas, and completed a family medicine residency at The Medical Center in Columbus, Ga.

ANTHONY A. DAY, M.D., is assistant executive director of the University of Illinois College of Medicine at Peoria family medicine residency program at Methodist Medical Center. He received his medical degree from the University of Iowa Roy J. and Lucille A. Carver College of Medicine, Iowa City, and completed a family medicine residency at the University of Illinois College of Medicine at Rockford.

Address correspondence to Craig P. Griebel, M.D., Department of Family and Community Medicine, University of Illinois College of Medicine at Peoria, 815 Main St., Suite C, Peoria, IL 61602 (e-mail: cgriebel@mnci.org). Reprints are not available from the authors.

TABLE 4
Points to Cover with Women and Their Partners After Spontaneous Abortion

Acknowledge and attempt to dispel guilt
 Acknowledge and legitimize grief
 Assess level of grief and adjust counseling accordingly
 Counsel how to tell family and friends of the miscarriage
 Include the patient's partner in psychologic care
 Provide comfort, empathy, and ongoing support
 Reassure about the future
 Warn about the "anniversary phenomenon"

Information from reference 5.

Members of various family medicine departments develop articles for "Practical Therapeutics." This article is one in a series coordinated by the Department of Family Practice at the University of Illinois College of Medicine at Chicago–Rockford. Coordinator of the series is Eric Henley, M.D.

Author disclosure: Nothing to disclose.

REFERENCES

- Scroggins KM, Smucker WD, Krishen AE. Spontaneous pregnancy loss: evaluation, management, and follow-up counseling. *Prim Care* 2000;27:153-67.
- Creinin MD, Schwartz JL, Guido RS, Pymar HC. Early pregnancy failure—current management concepts. *Obstet Gynecol Surv* 2001;56:105-13.
- Everett C. Incidence and outcome of bleeding before the 20th week of pregnancy: prospective study from general practice. *BMJ* 1997;315:32-4.
- Wilcox AJ, Weinberg CR, O'Connor JF, Baird DD, Schlatterer JP, Canfield RE, et al. Incidence of early loss of pregnancy. *N Engl J Med* 1988;319:189-94.
- Deutchman M, Eisinger S, Kelber M. First trimester pregnancy complications. In: *ALSO: Advanced Life Support in Obstetrics course syllabus*. 4th ed. Leawood, Kan.: American Academy of Family Physicians, 2000:1-27.
- Wieringa-de Waard M, Bonsel GJ, Ankum WM, Vos J, Bindels PJ. Threatened miscarriage in general practice: diagnostic value of history taking and physical examination. *Br J Gen Pract* 2002;52:825-9.
- Wong SF, Lam MH, Ho LC. Transvaginal sonography in the detection of retained products of conception after first-trimester spontaneous abortion. *J Clin Ultrasound* 2002;30:428-32.
- Rulin MC, Bornstein SG, Campbell JD. The reliability of ultrasonography in the management of spontaneous abortion, clinically thought to be complete: a prospective study. *Am J Obstet Gynecol* 1993;168(1 pt 1):12-5.
- Goddijn M, Leschot NJ. Genetic aspects of miscarriage. *Baillieres Best Pract Res Clin Obstet Gynaecol* 2000;14:855-65.
- Cunningham FG, Gant NF, Leveno KJ, Gilstrap LC, Hauth JC, Wenstrom KD. Spontaneous abortion. In: *Cunningham FG, Williams JW. Williams Obstetrics*. 21st ed. New York: McGraw-Hill, 2001:856-69.
- Garcia-Enguidanos A, Calle ME, Valero J, Luna S, Dominguez-Rojas V. Risk factors in miscarriage: a review. *Eur J Obstet Gynecol Reprod Biol* 2002;102:111-9.
- Rasch V. Cigarette, alcohol, and caffeine consumption: risk factors for spontaneous abortion. *Acta Obstet Gynecol Scand* 2003;82:182-8.
- Donders GG, Van Bulck B, Caudron J, Londers L, Vereecken A, Spitz B. Relationship of bacterial vaginosis and mycoplasmas to the risk of spontaneous abortion. *Am J Obstet Gynecol* 2000;183:431-7.
- Li DK, Liu L, Odouli R. Exposure to non-steroidal anti-inflammatory drugs during pregnancy and risk of miscarriage: population based cohort study. *BMJ* 2003;327:368.
- Nelson DB, Grisso JA, Joffe MM, Brensinger C, Shaw L, Datner E. Does stress influence early pregnancy loss? *Ann Epidemiol* 2003;13:223-9.
- Chung TK, Cheung LP, Sahota DS, Haines CJ, Chang AM. Spontaneous abortion: short-term complications following either conservative or surgical management. *Aust N Z J Obstet Gynaecol* 1998;38:61-4.
- Sairam S, Khare M, Michailidis G, Thilaganathan B. The role of ultrasound in the expectant management of early pregnancy loss. *Ultrasound Obstet Gynecol* 2001;17:506-9.
- Blohm F, Friden B, Platz-Christensen JJ, Milsom I, Nielsen S. Expectant management of first-trimester miscarriage in clinical practice. *Acta Obstet Gynecol Scand* 2003;82:654-8.
- Gronlund L, Gronlund AL, Clevin L, Andersen B, Palmgren N, Lidsgaard O. Spontaneous abortion: expectant management, medical treatment or surgical evacuation. *Acta Obstet Gynecol Scand* 2002;81:781-2.
- Luise C, Jermy K, Collons WP, Bourne TH. Expectant management of incomplete, spontaneous first-trimester miscarriage: outcome according to initial ultrasound criteria and value of follow-up visits. *Ultrasound Obstet Gynecol* 2002;19:580-2.
- Nielsen S, Hahlin M. Expectant management of first-trimester spontaneous abortion. *Lancet* 1995;345:84-6.
- Ankum WM, Wieringa-De Waard M, Bindels PJ. Management of spontaneous miscarriage in the first trimester: an example of putting informed shared decision making into practice. *BMJ* 2001;322:1343-6.
- Nielsen S, Hahlin M, Platz-Christensen J. Randomised trial comparing expectant with medical management for first trimester miscarriages. *Br J Obstet Gynaecol* 1999;106:804-7.
- Geyman JP, Oliver LM, Sullivan SD. Expectant medical or surgical treatment of spontaneous abortion in first trimester of pregnancy? A pooled quantitative literature evaluation. *J Am Board Fam Pract* 1999;12:55-64.
- Wood SL, Brain PH. Medical management of missed abortion: a randomized clinical trial [published correction appears in *Obstet Gynecol* 2002;100:175]. *Obstet Gynecol* 2002;99:563-6.

Spontaneous Abortion

26. Jurkovic D, Ross JA, Nicolaides KH. Expectant management of missed miscarriage. *Br J Obstet Gynaecol* 1998;105:670-1.
27. Pang MW, Lee TS, Chung TK. Incomplete miscarriage: a randomized controlled trial comparing oral with vaginal misoprostol for medical evacuation. *Hum Reprod* 2001;16:2283-7.
28. Wieringa-De Waard M, Hartman EE, Ankum WM, Reitsma JB, Bindels PJ, Bonsel GJ. Expectant management versus surgical evacuation in first trimester miscarriage: health-related quality of life in randomized and non-randomized patients. *Hum Reprod* 2002;17:1638-42.
29. Lee DT, Cheung LP, Haines CJ, Chan KP, Chung TK. A comparison of the psychologic impact and client satisfaction of surgical treatment with medical treatment of spontaneous abortion: a randomized controlled trial. *Am J Obstet Gynecol* 2001;185:953-8.
30. Molnar AM, Oliver LM, Geyman JP. Patient preferences for management of first-trimester incomplete spontaneous abortion. *J Am Board Fam Pract* 2000;13:333-7.
31. Neugebauer R, Kline J, O'Connor P, Shrout P, Johnson J, Skodol A, et al. Determinants of depressive symptoms in the early weeks after miscarriage. *Am J Public Health* 1992;82:1332-9.
32. Neugebauer R, Kline J, Shrout P, Skodol A, O'Connor P, Geller PA, et al. Major depressive disorder in the 6 months after miscarriage. *JAMA* 1997;277:383-8.
33. Janssen HJ, Cuisinier MC, Hoogduin KA, de Graauw KP. Controlled prospective study on the mental health of women following pregnancy loss. *Am J Psychiatry* 1996;153:226-30.
34. Thapar AK, Thapar A. Psychological sequelae of miscarriage: a controlled study using the general health questionnaire and the hospital anxiety and depression scale. *Br J Gen Pract* 1992;42:94-6.
35. Speraw SR. The experience of miscarriage: how couples define quality in health care delivery. *J Perinatol* 1994;14:208-15.
36. Lee C, Slade P. Miscarriage as a traumatic event: a review of the literature and new implications for intervention. *J Psychosom Res* 1996;40:235-44.