





**Figure 1.** Images of the ureter obtained during the excretory phase. (A) Urogram shows a radiolucent filling defect (*arrow*) in the distal left ureter. (B) Transaxial computed tomography (CT) scan shows a soft-tissue-filling defect (*arrow*) in the left ureter. (C) Curved reformatted image from the CT shows a stalk (*black arrowhead*) in the filling defect; an additional small filling defect (*white arrow*) proximal to the lesion; and a defect (*black arrow*) near the ureteric insertion into the bladder.

revealed a soft-tissue–filling defect in the left ureter (*Figure 1B*). A curved reformatted image was obtained from the excretory-phase enhanced transaxial CT urography and showed that the polypoid filling defect has a stalk, an additional small polypoid ureteral filling defect was present, and there was a defect near the ureteric insertion into the bladder (*Figure 1C*).

The patient underwent a left distal ureterectomy with a left ureteroneocystostomy. The tumors were noninvasive grade 2 (of 3) papillary urothelial carcinomas (transitional cell carcinoma). The bladder lesions were also noninvasive grade 2 (of 3)

papillary urothelial carcinomas and were treated with fulguration.

### Must Imaging Always Be Performed?

There may be specific circumstances in which a complete radiologic work-up is not necessary.<sup>6</sup> Young women with a clinical picture of simple cystitis and whose hematuria completely and permanently resolves after successful therapy will probably not require any imaging.<sup>7,8</sup> Patients who have clear-cut evidence of glomerulopathy also constitute a special group—chest radiography<sup>8</sup> to search for any of the numerous manifestations of glomerulonephritis (including cardiac

enlargement, pleural and pericardial effusions, pulmonary congestion and edema, and pulmonary bleeding) and ultrasonography (to display the site and number of kidneys before biopsy and to screen for renal morphologic abnormalities that may coexist by chance in a patient with glomerulonephritis) should be obtained.<sup>9-13</sup> However, the decision to pursue this course requires firm demonstration that the glomerular abnormality is responsible for the hematuria; such evidence includes heavy proteinuria (sufficient to indicate that plasma proteins, rather than proteins in red cells, account for the protein in the urine), red cell casts, or evidence of severe red cell dysmorphism (only given reliable methods of identifying such abnormalities).

Although hematuria in patients taking anticoagulants is often blamed on excessive anticoagulation, significant pathologies have been detected, justifying radiologic work-ups in this group as well.<sup>14</sup> All other adult patients require imaging evaluation<sup>7,9,15</sup> (Table 1<sup>16</sup>).

### Radiologic Work-up of Hematuria

Radiologic evaluation will almost always be accompanied by cystoscopy because many bleeding urinary tract lesions arise in the bladder and lower urinary tract, and no imaging technique is completely satisfactory for ruling out disease at these sites.<sup>2</sup> A complete history, physical examination, urinalysis, and appropriate blood tests should precede or accompany the imaging examinations. At the time of cystoscopy, bilateral retrograde pyelography (in which iodinated contrast is injected through catheters placed in the ureters during cystoscopy and plain radiographs are obtained) is often employed to evaluate the upper tracts for pathology.<sup>6</sup>

There is no universal agreement about the optimal imaging work-up of hematuria. Traditionally, excretory urography (intravenous pyelogram) was the standard,<sup>6-8</sup> but the establishment of this practice preceded the development of high-quality ultrasonography,<sup>17</sup> CT, and magnetic resonance imaging (MRI). Recently, multidetector CT scans, in which each rotation of the radiography beam

produces multiple sets of images instead of just one set, have become routine. They provide cross-sectional images and can be reformatted to demonstrate the urinary tract in a manner similar to traditional intravenous pyelograms. Similarly, MRI can be used to detect urinary tract abnormalities, but has limited use because of its expense and the lack of data supporting its use.

### Increasing Role of CT Urography

Conventional abdominal CT of the entire urinary tract can be augmented by delayed thin-section images of the contrast-opacified collecting systems, ureters, and bladder<sup>18</sup>; the combined examination is known as CT urography. The intravenous-pyelogram-like portions of the examination may be obtained by exposing film (or direct digital) images when intravenous contrast, previously administered for the CT, has opacified the urinary tract. Intravenous-pyelogram-like images may alternatively be produced by reformatting delayed CT images in the coronal or sagittal planes. Presumably, the pyelogram portion of this examination could be comparable with a standard intravenous pyelogram examination, making CT more sensitive and specific (statistically and pathologically) than ultrasonography or nephrotomography with regard to focal renal parenchymal abnormalities.

Nephrotomography is an older technique, used before the invention of CT, which images 1-cm coronal sections of the kidneys as part of an intravenous pyelogram. Thus, a distinction should be made between a routine CT of the abdomen and pelvis that may not be optimized for the urinary tract and a dedicated CT urogram that is tailored to evaluate the urinary tract for sources of hematuria. The latter study typically employs oral water instead of oral positive contrast media, as this will greatly aid the reconstruction of images.

In the CT urogram, all patients receive water, primarily to hydrate the kidneys and distend the collecting system and ureter. Next, a noncontrast helical CT of the kidneys is obtained to evaluate renal calculi. This is followed by the injection of iodinated

## Radiologic Evaluation of Hematuria

**Table 1. ACR Appropriateness Criteria Scale for Hematuria**

<i>Radiologic examination procedure</i>	<i>Appropriateness rating</i>	<i>Comments</i>
Multidetector CT urography	8	This is becoming the method of choice for hematuria, supplanting intravenous pyelography, even though their appropriateness ratings are the same
Radiography, intravenous urography (intravenous pyelogram, excretory urography)	8	—
Ultrasonography, kidney and bladder, transabdominal	6	May miss ureteral and urothelial lesions; abdominal radiography, retrograde pyelography, and cystoscopy are useful adjuncts
Radiography, retrograde urography	5	—
MRI urography	4	—
CT, abdomen and pelvis	4	CT may follow intravenous pyelogram or ultrasonography if initial findings are ambiguous
Kidney, angiography	4	Rarely, vascular malformations may cause hematuria and require angiography for diagnosis
Radiography, abdomen, KUB	2	It is assumed that a plain film of the abdomen will be part of the indicated intravenous pyelogram; if an intravenous pyelogram is not performed, KUB may be performed with ultrasonography
MRI, abdomen, and pelvis	2	—
Urinary tract scintigraphy	2	—
Virtual cystoscopy	2	—

NOTE: Appropriateness scale has a range of 1 to 9, with 1 = least appropriate and 9 = most appropriate; the criteria in this table apply to all patients except those with generalized renal parenchymal disease or young females with hemorrhagic cystitis.

ACR = American College of Radiology; CT = computed tomography; KUB = kidneys, ureters, and bladder; MRI = magnetic resonance imaging.

Adapted with permission from American College of Radiology. ACR appropriateness criteria. Topic: hematuria. Variant: all patients except those with generalized renal parenchymal disease or young females with hemorrhagic cystitis. <http://acsearch.acr.org/procedureslist.aspx?tid=30681&vid=3019558>. Accessed March 26, 2008.

contrast media with the acquisition of a high-resolution (1- to 2-mm sections) nephrographic phase and a high-resolution delayed (five to 10 minutes) phase to evaluate for tumors and filling defects. The latter can be reconstructed to evaluate the urinary tract and bladder. Some investigators employ a hybrid of CT urography and intravenous-pyelogram-like delayed images to form one complete study, which is also known as CT urography and has shown equal or

superior sensitivity to intravenous pyelogram for causes of hematuria.<sup>19,20</sup>

Virtual cystoscopy—the acquisition of high-resolution CT images reconstructed to allow virtual “fly-throughs” of the bladder—can be used to evaluate the bladder for causes of hematuria.<sup>21</sup> Virtual cystoscopy is inaccurate for small lesions and lesions located near the ureteric orifices. The urethra cannot be evaluated, so virtual cystoscopy cannot replace actual cystoscopy.

### Role of Ultrasonography and MRI

There is some support for the use of ultrasonography as the initial imaging study for selected patients with hematuria.<sup>17,22,23</sup> Ultrasonography is believed to have moderately high sensitivity with respect to the wide range of abnormalities that may be encountered, including urinary tract neoplasms of all sorts, stone disease, inflammatory processes, congenital abnormalities, vascular lesions, and obstruction from a wide variety of lesions.<sup>5,22-24</sup> Nevertheless, it appears that ultrasonography has similar rates of detection compared with excretory urography (intravenous pyelogram) for diagnosing clinically important lesions.<sup>25</sup> Ultrasonography and urography tend to miss different sorts of lesions, and ultrasonography is not likely to detect nonobstructing ureteral stones or small urothelial abnormalities, whereas urography with nephrotomography may miss small exophytic anterior and posterior renal masses and small bladder lesions.<sup>26,27</sup>

The choice of examination may be affected by clinical circumstances (e.g., a positive urinary cytologic analysis may make urography crucial, whereas serious risk factors for contrast reactions may make ultrasonography more appropriate). When ultrasonography is negative and the source of hematuria remains obscure, urography should be added; if urography is negative, CT may be ordered.<sup>8,27,28</sup> When ultrasonography is used as the primary screening modality, the yield from imaging may be increased by adding a plain film of the abdomen. Ultrasonography has a particularly important role in children and pregnant women with hematuria, in whom ionizing radiation must be avoided.

MRI urography currently serves as an alternative imaging technique for children and pregnant women and for patients with a contraindication to iodinated contrast media.<sup>29</sup> It has the potential to be useful in the search for important abnormalities (e.g., urothelial cancers, stones, renal tumors) that cause hematuria. MRI urography has not been widely adopted in clinical practice, is expensive, and has not been evaluated for

**Table 2. Causes of Hematuria**

#### Causes of hematuria in adults\*

- Calculi
- Infection
- Cancer (bladder, kidney, prostate, urothelial)
- Obstruction
- Bleeding diathesis
- Anticoagulation
- Antibiotics (rifampin [Rifadin])
- Diabetes
- Hypertension
- Sickle cell anemia
- Chronic renal disease
- Congenital
- Vascular malformations and aneurysms

#### Artifactual causes of hematuria

- Food (e.g., beets, berries, rhubarb)
- Food coloring
- Medications
- Menstrual blood

\*—The first five causes listed in the table are the most common.

Information from reference 30.

effectiveness, so it cannot be recommended as an initial examination.

### Final Comments

Most adults with hematuria require urinary tract imaging. Intravenous pyelogram and CT urography represent the leading techniques, with an increasing trend for the latter. Ultrasonography and MRI have secondary roles in selected populations. Carefully selected patients may require no further work-up if their clinical history can reasonably determine the cause of their hematuria (Table 2<sup>30</sup>).

The author recognizes the Urologic Imaging Expert Panel of the American College of Radiology that contributed to this version of the Appropriateness Criteria for Hematuria: Edward I. Bluth, MD; William H. Bush, Jr., MD; Davide D. Casalino, MD; Isaac Francis, MD; S. Zafar H. Jafri, MD; Akira Kawashima, MD, PhD; Robert Older, MD; Nicholas Papanicolaou, MD; Parvati Ramchandani, MD; Arthur T. Rosenfield, MD; Carl Sandler, MD; Arthur J. Segal, MD; Clare Tempany, MD; and Martin Resnick, MD.

Figure 1 courtesy of Akira Kawashima, MD, PhD, Department of Radiology, Mayo Clinic, Rochester, Minn.

## The Author

PETER L. CHOYKE, MD, is chief of the molecular imaging program at the National Cancer Institute in Bethesda, Md. Dr. Choyke received his medical degree from Jefferson Medical College, Philadelphia, Pa. He completed a residency in diagnostic radiology at Yale University, New Haven, Conn.

Address correspondence to Peter L. Choyke, MD, National Cancer Institute, Building 10, Room 1B40, Bethesda, MD 20892-1088 (e-mail: pchoyke@nih.gov). Reprints are not available from the author.

Author disclosure: Nothing to disclose.

## REFERENCES

- Grossfeld GD, Wolf JS Jr, Litwan MS, et al. Asymptomatic microscopic hematuria in adults: summary of the AUA best practice policy recommendations. *Am Fam Physician*. 2001;63(6):1145-1154.
- McDonald MM, Swagerty D, Wetzel L. Assessment of microscopic hematuria in adults. *Am Fam Physician*. 2006;73(10):1748-1754.
- Froom P, Ribak J, Benbassat J. Significance of microhaematuria in young adults. *Br Med J (Clin Res Ed)*. 1984;288(6410):20-22.
- Bender CB. Evaluation of the urologic patient. In: Campbell MF, Walsh PC. *Campbell's Urology*. 6th ed. Philadelphia, Pa: Saunders; 1992:307-317.
- Messing EM, Young TB, Hunt VB, Emoto SE, Wehbie JM. The significance of asymptomatic microhematuria in men 50 or more years old: findings of a home screening study using urinary dipsticks. *J Urol*. 1987;137(5):919-922.
- Abuelo JG. The diagnosis of hematuria. *Arch Intern Med*. 1983;143(5):967-970.
- Benson GS, Brewer ED. Hematuria: algorithms for diagnosis. II. Hematuria in the adult and hematuria secondary to trauma. *JAMA*. 1981;246(9):993-995.
- Copley JB. Isolated asymptomatic hematuria in the adult. *Am J Med Sci*. 1986;291(2):101-111.
- Pulmonary hypertension and edema. In: Fraser RG. *Diagnosis of Diseases of the Chest*. 3rd ed. Philadelphia, Pa.: Saunders; 1990:1823-1968.
- Abuelo JG. Evaluation of hematuria. *Urology*. 1983;21(3):215-225.
- Fairley K. Urinalysis. In: Schrier RW, Gottschalk CW. *Diseases of the Kidney*. 4th ed. Boston, Mass.: Little Brown; 1988:359-383.
- Sutton JM. Evaluation of hematuria in adults. *JAMA*. 1990;263(18):2475-2480.
- Avidor Y, Nadu A, Matzkin H. Clinical significance of gross hematuria and its evaluation in patients receiving anticoagulant and aspirin treatment. *Urology*. 2000;55(1):22-24.
- Golin AL, Howard RS. Asymptomatic microscopic hematuria. *J Urol*. 1980;124(3):389-391.
- Datta SN, Allen GM, Evans R, Vaughton KC, Lucas MG. Urinary tract ultrasonography in the evaluation of haematuria—a report of over 1,000 cases. *Ann R Coll Surg Engl*. 2002;84(3):203-205.
- American College of Radiology. ACR appropriateness criteria. Topic: hematuria. Variant: all patients except those with generalized renal parenchymal disease or young females with hemorrhagic cystitis. <http://acsearch.acr.org/procedureslist.aspx?tid=30681&vid=3019558>. Accessed March 26, 2008.
- Chisholm RA, Millet B, Sherwood T, Wright EP, Doyle PT. The investigation of painless haematuria—a comparison of intravenous urography and DMSA scintigraphy. *Clin Radiol*. 1988;39(5):494-495.
- McTavish JD, Jinzaki M, Zou KH, Nawfel RD, Silverman SG. Multi-detector row CT urography: comparison of strategies for depicting the normal urinary collecting system. *Radiology*. 2002;225(3):783-790.
- Joffe SA, Servaes S, Okon S, Horowitz M. Multi-detector row CT urography in the evaluation of hematuria. *Radiographics*. 2003;23(6):1441-1455.
- McNicholas MM, Raptopoulos VD, Schwartz RK, et al. Excretory phase CT urography for opacification of the urinary collecting system. *AJR Am J Roentgenol*. 1998;170(5):1261-1267.
- Nambirajan T, Sohaib SA, Muller-Pollard C, Reznik R, Chingwundoh FI. Virtual cystoscopy from computed tomography: a pilot study. *BJU Int*. 2004;94(6):828-831.
- Corwin HL, Silverstein MD. The diagnosis of neoplasia in patients with asymptomatic microscopic hematuria: a decision analysis [published correction appears in *J Urol*. 1988;140(6):1558]. *J Urol*. 1988;139(5):1002-1006.
- Murakami S, Igarashi T, Hara S, Shimazaki J. Strategies for asymptomatic microscopic hematuria: a prospective study of 1,034 patients. *J Urol*. 1990;144(1):99-101.
- Mariani AJ, Mariani MC, Macchioni C, Stams UK, Hariharan A, Moriera A. The significance of adult hematuria: 1,000 hematuria evaluations including a risk-benefit and cost-effectiveness analysis. *J Urol*. 1989;141(2):350-355.
- Mohr DN, Offord KP, Owen RA, Metlon LJ III. Asymptomatic microhematuria and urologic disease. A population-based study. *JAMA*. 1986;256(2):224-229.
- Aslaksen A, Gadeholt G, Göthlin JH. Ultrasonography versus intravenous urography in the evaluation of patients with microscopic haematuria. *Br J Urol*. 1990;66(2):144-147.
- Amendola MA, Bree RL, Pollack HM, et al. Small renal cell carcinomas: resolving a diagnostic dilemma. *Radiology*. 1988;166(3):637-641.
- Glen DA, Gilbert FJ, Bayliss AP. Renal carcinoma missed by urography. *Br J Urol*. 1989;63(5):457-459.
- Kawashima A, Glockner JF, King BF Jr. CT urography and MR urography. *Radiol Clin North Am*. 2003;41(5):945-961.
- Gerber GS, Brendler CB. Evaluation of the urologic patient: history, physical examination, and urinalysis. In: Wein AJ, ed. *Campbell-Walsh Urology*. 9th ed. Philadelphia, Pa.: Saunders; 2007:81-110.