

Diagnosis and Treatment of Acute Uncomplicated Cystitis

RICHARD COLGAN, MD, and MOZELLA WILLIAMS, MD
University of Maryland School of Medicine, Baltimore, Maryland

Urinary tract infections are the most common bacterial infections in women. Most urinary tract infections are acute uncomplicated cystitis. Identifiers of acute uncomplicated cystitis are frequency and dysuria in an immunocompetent woman of childbearing age who has no comorbidities or urologic abnormalities. Physical examination is typically normal or positive for suprapubic tenderness. A urinalysis, but not urine culture, is recommended in making the diagnosis. Guidelines recommend three options for first-line treatment of acute uncomplicated cystitis: fosfomycin, nitrofurantoin, and trimethoprim/sulfamethoxazole (in regions where the prevalence of *Escherichia coli* resistance does not exceed 20 percent). Beta-lactam antibiotics, amoxicillin/clavulanate, cefaclor, cefdinir, and cefpodoxime are not recommended for initial treatment because of concerns about resistance. Urine cultures are recommended in women with suspected pyelonephritis, women with symptoms that do not resolve or that recur within two to four weeks after completing treatment, and women who present with atypical symptoms. (*Am Fam Physician*. 2011;84(7):771-776. Copyright © 2011 American Academy of Family Physicians.)



ILLUSTRATION BY JOHN W. KARAFILOU

► **Patient information:** A handout on treating a bladder infection (cystitis), written by the authors of this article, is provided on page 778.

Urinary tract infections (UTIs) are the most common bacterial infections in women, with one-half of all women experiencing at least one UTI in their lifetime.¹ Most UTIs in women are acute uncomplicated cystitis caused by *Escherichia coli* (86 percent), *Staphylococcus saprophyticus* (4 percent), *Klebsiella* species (3 percent), *Proteus* species (3 percent), *Enterobacter* species (1.4 percent), *Citrobacter* species (0.8 percent), or *Enterococcus* species (0.5 percent).² Although acute uncomplicated cystitis may not be thought of as a serious condition, patients' quality of life is often significantly affected. Acute uncomplicated cystitis results in an estimated six days of discomfort leading to approximately 7 million office visits per year with associated costs of \$1.6 billion.^{3,4} In one study of women with acute uncomplicated cystitis, nearly one-half of participants reported that their symptoms caused them

to miss work or school.³ Additionally, up to one-half of those with acute uncomplicated cystitis also reported avoiding sexual activity for an average of one week.

Diagnosis

The history is the most important tool for diagnosing acute uncomplicated cystitis, and it should be supported by a focused physical examination and urinalysis. It also is important to rule out a more serious complicated UTI. By definition, the diagnosis of acute uncomplicated cystitis implies an uncomplicated UTI in a premenopausal, nonpregnant woman with no known urologic abnormalities or comorbidities (*Table 1*⁵).

Classic lower urinary tract symptoms include dysuria, frequent voiding of small volumes, and urinary urgency. Sometimes hematuria can occur; suprapubic discomfort is less common. The pretest probability of UTI in women is 5 percent; however,

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
The combination of new-onset frequency and dysuria, with the absence of vaginal discharge, is diagnostic for a urinary tract infection.	C	6
A urine culture is recommended for women with suspected acute pyelonephritis, women with symptoms that do not resolve or that recur within two to four weeks after the completion of treatment, and women who present with atypical symptoms.	C	11
First-line treatment options for acute uncomplicated cystitis include nitrofurantoin (macrocrystals; 100 mg twice per day for five days), trimethoprim/sulfamethoxazole (Bactrim, Septra; 160/800 mg twice per day for three days in regions where the uropathogen resistance is less than 20 percent), and fosfomycin (Monurol; a single 3-g dose).	C	16

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort.xml>.

when a woman presents with the acute onset of even one of the classic symptoms of acute uncomplicated cystitis, the probability of infection rises 10-fold to 50 percent.⁶ Therefore, presentation with one or more symptoms may be viewed as a valuable diagnostic

test in itself. In addition, the likelihood of acute uncomplicated cystitis is less if the patient reports vaginal discharge or irritation, both of which are more likely in women with vaginitis or cervicitis. The new onset of frequency and dysuria, with the absence of vaginal discharge or irritation, has a positive predictive value of 90 percent for UTI.⁶ A prospective study of 796 sexually active young women identified risk factors to help diagnose UTI, including recent sexual intercourse, diaphragm use with spermicide, and recurrent UTIs.⁷

Table 1. Characteristics of Patients with Uncomplicated and Complicated Urinary Tract Infections

Uncomplicated

- Immunocompetent
- No comorbidities
- No known urologic abnormalities
- Nonpregnant
- Premenopausal

Complicated*

- History of childhood urinary tract infections
- Immunocompromised
- Preadolescent or postmenopausal
- Pregnant
- Underlying metabolic disorder (e.g., diabetes mellitus)
- Urologic abnormalities (e.g., stones, stents, indwelling catheters, neurogenic bladder, polycystic kidney disease)

*—Urinary tract infections in men are usually complicated.

Information from reference 5.

Self-Diagnosis and Diagnosis by Telephone

For many patients, access to care can be difficult. Two recent studies have shown that some women who self-diagnose a UTI may be treated safely with telephone management. Women who have had acute uncomplicated cystitis previously are usually accurate in determining when they are having another episode. In one study of 172 women with a history of recurrent UTI, 88 women self-diagnosed a UTI based on symptoms, and self-treated with antibiotics.⁸ Laboratory evaluation showed that 84 percent of the urine samples showed a uropathogen, 11 percent showed sterile pyuria, and only 5 percent were negative for pyuria and bacteriuria. Another small, randomized

controlled trial compared outcomes of acute uncomplicated cystitis in healthy women managed by telephone versus in the office.⁹ There were no differences in symptom score or satisfaction. The authors concluded that the short-term outcomes of managing suspected UTIs by telephone were comparable with those managed by usual office care.

Physical Examination and Diagnostic Testing

The physical examination of patients with acute uncomplicated cystitis is typically normal, except in the 10 to 20 percent of women with suprapubic tenderness.¹⁰ Acute pyelonephritis should be suspected if the patient is ill-appearing and seems uncomfortable, particularly if she has concomitant fever, tachycardia, or costovertebral angle tenderness.

The convenience and cost-effectiveness of urine dipstick testing makes it a common diagnostic tool, and it is an appropriate alternative to urinalysis and urine microscopy to diagnose acute uncomplicated cystitis.¹¹ Nitrites and leukocyte esterase are the most accurate indicators of acute uncomplicated cystitis in symptomatic women.¹¹ To avoid contamination, the convention is to use a midstream, clean-catch urine specimen to diagnose UTI; however, at least two studies have shown no significant difference in number of contaminated or unreliable results between specimens collected with and without preparatory cleansing.^{12,13} Urine cultures are recommended only for patients with suspected acute pyelonephritis; patients with symptoms that do not resolve or that recur within two to four weeks after the completion of treatment; and patients who present with atypical symptoms.¹¹ A colony count greater than or equal to 10^3 colony-forming units per mL of a uropathogen is diagnostic of acute uncomplicated cystitis.¹⁴ However, studies have shown that more than 10^2 colony-forming-units per mL in women with typical symptoms of a UTI represent a positive culture.¹⁵ Routine posttreatment urinalysis or urine cultures in asymptomatic patients are not necessary.

Further studies beyond urinalysis and urine cultures are rarely needed to diagnose

acute uncomplicated cystitis. Patients who present with atypical symptoms of acute uncomplicated cystitis and those who do not respond to appropriate antimicrobial therapy may need imaging studies, such as computed tomography or ultrasonography, to rule out complications and other disorders.

International Clinical Practice Guidelines

In 2010, a panel of international experts updated the 1999 Infectious Diseases Society of America (IDSA) guidelines on the treatment of acute uncomplicated cystitis and pyelonephritis in women. The panel reviewed the literature, including the Cochrane Database of Systematic Reviews, and provided an evidence-based guideline for women with uncomplicated bacterial cystitis and pyelonephritis.^{16,17} The IDSA collaborated with the European Society of Clinical Microbiology and Infectious Diseases, and invited representation from diverse geographic areas and a wide variety of specialties, including urology, obstetrics and gynecology, emergency medicine, family medicine, internal medicine, and infectious diseases. Levels-of-evidence ratings were assigned to recommendations on the use of antimicrobials for the treatment of uncomplicated UTIs.

Treatment

No single agent is considered best for treating acute uncomplicated cystitis according to the 2010 guidelines, and the choice between recommended agents should be individualized¹⁶ (*Table 2*^{18,19}). Choosing an antibiotic depends on the agent's effectiveness, risks of adverse effects, resistance rates, and propensity to cause collateral damage (i.e., ecologic adverse effects of antibiotic therapy that may allow drug-resistant organisms to proliferate, and the colonization or infection with multidrug-resistant organisms). Additionally, physicians should consider cost, availability, and specific patient factors, such as allergy history. On average, patients will begin noting symptom relief within 36 hours of beginning treatment.²

Nitrites and leukocyte esterase on urine dipstick testing are the most accurate indicators of acute uncomplicated cystitis.

Table 2. Antimicrobial Agents for the Management of Acute Uncomplicated Cystitis

Tier	Drug	Dosage	Cost of generic (brand)	Pregnancy category
First	Fosfomycin (Monurol)	3-g single dose	NA (\$51)*	B
	Nitrofurantoin (macrocrystals)	100 mg twice per day for five days	\$55 (\$64)*	B
	Trimethoprim/sulfamethoxazole (Bactrim, Septra)	160/800 mg twice per day for three days	\$17 (\$34)*†	C
Second	Ciprofloxacin (Cipro)	250 mg twice per day for three days	\$26 (\$30)†‡	C
	Ciprofloxacin, extended release (Cipro XR)	500 mg per day for three days	\$57 (\$76)*	C
	Levofloxacin (Levaquin)	250 mg per day for three days	NA (\$86)*	C
	Ofloxacin	200 mg per day for three days or 400-mg single dose	\$14 (NA)‡ \$10 (NA)‡	C
Third§	Amoxicillin/clavulanate (Augmentin)	500/125 mg twice per day for seven days	\$32 (\$98)*	B
	Cefdinir (Omnicef)	300 mg twice per day for 10 days	\$40 (\$119)*	B
	Cefpodoxime	100 mg twice per day for seven days	\$71 (NA)‡	B

NA = not available.

*—Estimated retail price of one course of treatment based on information obtained at <http://www.drugstore.com> (accessed May 11, 2011).

†—May be available at discounted prices (\$10 or less for one month's treatment) at one or more national retail chains.

‡—Estimated cost to the pharmacist based on average wholesale prices in Red Book. Montvale, N.J.: Medical Economics Data; 2010. Cost to the patient will be higher, depending on prescription filling fee.

§—Not generally recommended because of relatively high rates of resistance. Third-tier options include beta-lactam antibiotics.

Information from references 18 and 19.

There are several first-line agents recommended by the IDSA for the treatment of acute uncomplicated cystitis (Figure 1).¹⁶ New evidence supports the use of nitrofurantoin (macrocrystals) and fosfomycin (Monurol) as first-line therapy.¹⁶ The following antimicrobials represent the first tier: (1) nitrofurantoin at a dosage of 100 mg twice per day for five days; (2) trimethoprim/sulfamethoxazole (Bactrim, Septra) at a dosage of one double-strength tablet (160/800 mg) twice per day for three days in regions where the prevalence of resistance of community uropathogens does not exceed 20 percent; and (3) fosfomycin at a single dose of 3 g. Note that the duration of therapy for nitrofurantoin has been reduced to five days compared with the previous IDSA guidelines of seven days, based on research showing

effectiveness with a shorter duration of therapy.²⁰ Fosfomycin may be less effective and is not widely available in the United States.

Fluoroquinolones (i.e., ofloxacin, ciprofloxacin [Cipro], and levofloxacin [Levaquin]) are considered second-tier antimicrobials, and are appropriate in some settings, such as in patients with allergy to the recommended agents. Although fluoroquinolones are effective, they have the propensity for collateral damage, and should be considered for patients with more serious infections than acute uncomplicated cystitis. Certain antimicrobials (i.e., beta-lactam antibiotics, amoxicillin/clavulanate [Augmentin], cefdinir [Omnicef], cefaclor, and cefpodoxime) may be appropriate alternatives if recommended agents cannot be used because of known resistance or patient intolerance. Despite wide use of

cranberry products for treating UTIs, there is no evidence to support their use in symptomatic patients.²¹


Antimicrobial Resistance

Beta-lactam antibiotics are not recommended as first-line therapy for acute uncomplicated cystitis because of widespread *E. coli* resistance rates above 20 percent. Fluoroquinolone resistance usually is found to be below 10 percent in North America and Europe, but with a trend toward increasing resistance

over the past several years.¹⁶ To preserve the effectiveness of fluoroquinolones, they are not recommended as a first-tier option. Fosfomycin and nitrofurantoin have retained high rates of in vitro activity in most areas.¹⁶

Because results of urine cultures are not routinely reported when treating acute uncomplicated cystitis, local resistance rates may not be available. Defaulting to the annual antimicrobial sensitivity data from a local hospital may provide resistance rates based on a population that

Choosing an Antimicrobial Agent for Empiric Treatment of Acute Uncomplicated Cystitis



*—The choice among these agents should be individualized and based on patient allergy and compliance history, local practice patterns, local community resistance prevalence, availability, cost, and patient and physician threshold for failure.

Figure 1. Algorithm for choosing an antimicrobial agent for empiric treatment of acute uncomplicated cystitis.

Adapted with permission from Gupta K, Hooton TM, Naber KG, et al. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin Infect Dis.* 2011;52(5):e104.

Cystitis

does not reflect women with simple acute uncomplicated cystitis (e.g., sicker patients, inpatients, patients of all ages, male patients). Several studies have been published that may help predict the likelihood of *E. coli* resistance to trimethoprim/sulfamethoxazole in patients with acute uncomplicated cystitis. Use of trimethoprim/sulfamethoxazole in the preceding three to six months has been found to be an independent risk factor for resistance in women with acute uncomplicated cystitis.^{22,23} In addition, two U.S. studies demonstrated that travel outside the United States in the preceding three to six months was independently associated with trimethoprim/sulfamethoxazole resistance.^{24,25}

The authors thank Kalpana Gupta, MD, for her review of the manuscript.

The Authors

RICHARD COLGAN, MD, is an associate professor and director of medical student education in the Department of Family and Community Medicine at the University of Maryland School of Medicine in Baltimore.

MOZELLA WILLIAMS, MD, is an assistant professor and assistant director of medical student education in the Department of Family and Community Medicine at the University of Maryland School of Medicine.

Address correspondence to Richard Colgan, MD, University of Maryland School of Medicine, 29 South Paca St., Baltimore, MD 21201 (e-mail: rcolgan@som.umaryland.edu). Reprints are not available from the authors.

Author disclosure: No relevant financial affiliations to disclose.

REFERENCES

1. Nicolle LE. Epidemiology of urinary tract infection. *Infect Med*. 2001;18:153-162.
2. Gupta K, Scholes D, Stamm WE. Increasing prevalence of antimicrobial resistance among uropathogens causing acute uncomplicated cystitis in women. *JAMA*. 1999;281(8):736-738.
3. Colgan R, Keating K, Dougouih M. Survey of symptom burden in women with uncomplicated urinary tract infections. *Clin Drug Investig*. 2004;24(1):55-60.
4. Foxman B. Epidemiology of urinary tract infections: incidence, morbidity, and economic costs. *Am J Med*. 2002;113(suppl 1A):5S-13S.
5. Nicolle L; AMMI Canada Guidelines Committee. Complicated urinary tract infection in adults. *Can J Infect Dis Med Microbiol*. 2005;16(6):349-360.
6. Bent S, Nallamothu BK, Simel DL, Fihn SD, Saint S. Does this woman have an acute uncomplicated urinary tract infection? *JAMA*. 2002;287(20):2701-2710.
7. Hooton TM, Scholes D, Hughes JP, et al. A prospective study of risk factors for symptomatic urinary tract infection in young women. *N Engl J Med*. 1996;335(7):468-474.
8. Gupta K, Hooton TM, Roberts PL, Stamm WE. Patient-initiated treatment of uncomplicated recurrent urinary tract infections in young women. *Ann Intern Med*. 2001;135(1):9-16.
9. Barry HC, Hickner J, Ebell MH, Ettenhofer T. A randomized controlled trial of telephone management of suspected urinary tract infections in women. *J Fam Pract*. 2001;50(7):589-594.
10. Stamm WE. Urinary tract infections. In: Root RK, Waldvogel F, Corey L, Stamm WE. *Clinical Infectious Diseases: A Practical Approach*. New York, NY: Oxford University Press; 1999:649-656.
11. Colgan R, Hyner S, Chu S. Uncomplicated urinary tract infections in adults. In: Grabe M, Bishop MC, Bjerklund-Johansen, et al., eds. *Guidelines on Urological Infections*. Arnhem, The Netherlands: European Association of Urology; 2009:11-38.
12. Bradbury SM. Collection of urine specimens in general practice: to clean or not to clean? *J R Coll Gen Pract*. 1988;38(313):363-365.
13. Lifshitz E, Kramer L. Outpatient urine culture: does collection technique matter? *Arch Intern Med*. 2000;160(16):2537-2540.
14. Stamm WE. Criteria for the diagnosis of urinary tract infection and for the assessment of therapeutic effectiveness. *Infection*. 1992;20(suppl 3):S151-S154.
15. Kunin CM. Guidelines for urinary tract infections. Rationale for a separate strata for patients with "low-count" bacteriuria. *Infection*. 1994;22(suppl 1):S38-S40.
16. Gupta K, Hooton TM, Naber KG, et al. International clinical practice guidelines for the treatment of acute uncomplicated cystitis and pyelonephritis in women: a 2010 update by the Infectious Diseases Society of America and the European Society for Microbiology and Infectious Diseases. *Clin Infect Dis*. 2011;52(5):e103-e120.
17. Zalmanovici Trestioreanu A, Green H, Paul M, Yaphe J, Leibovici L. Antimicrobial agents for treating uncomplicated urinary tract infection in women. *Cochrane Database Syst Rev*. 2010;(10):CD007182.
18. Mehnert-Kay SA. Diagnosis and management of uncomplicated urinary tract infections. *Am Fam Physician*. 2005;72(3):451-456.
19. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin No. 91: treatment of urinary tract infections in nonpregnant women. *Obstet Gynecol*. 2008;111(3):785-794.
20. Gupta K, Hooton TM, Roberts PL, Stamm WE. Short-course nitrofurantoin for the treatment of acute uncomplicated cystitis in women. *Arch Intern Med*. 2007;167(20):2207-2212.
21. Jepson RG, Mihaljevic L, Craig J. Cranberries for treating urinary tract infections. *Cochrane Database Syst Rev*. 2000;(2):CD001322.
22. Brown PD, Freeman A, Foxman B. Prevalence and predictors of trimethoprim-sulfamethoxazole resistance among uropathogenic *Escherichia coli* isolates in Michigan. *Clin Infect Dis*. 2002;34(8):1061-1066.
23. Metlay JP, Strom BL, Asch DA. Prior antimicrobial drug exposure: a risk factor for trimethoprim-sulfamethoxazole-resistant urinary tract infections. *J Antimicrob Chemother*. 2003;51(4):963-970.
24. Burman WJ, Breese PE, Murray BE, et al. Conventional and molecular epidemiology of trimethoprim-sulfamethoxazole resistance among urinary *Escherichia coli* isolates. *Am J Med*. 2003;115(5):358-364.
25. Colgan R, Johnson JR, Kuskowski M, Gupta K. Risk factors for trimethoprim-sulfamethoxazole resistance in patients with acute uncomplicated cystitis. *Antimicrob Agents Chemother*. 2008;52(3):846-851.