

## Clinical Preventive Services Recommendations for Adolescents and Young Adults

	Ages 11-13	Ages 14-17	Ages 18-26
Universal Screening and Counseling (Individuals at average risk)	Body Mass Index Depression Tobacco Use Skin Cancer Prevention (fair skin)	Body Mass Index Depression Tobacco Use Skin Cancer Prevention (fair skin) Intimate Partner Violence (females)	Body Mass Index Depression Tobacco Use Skin Cancer Prevention (fair skin) Intimate Partner Violence (females) Blood Pressure HIV Alcohol Misuse Cervical Cancer (ages ≥ 21, q3 years)
	If sexually active: - Gonorrhea/Chlamydia (females) - Contraception - Prevention of sexually transmitted infections	If sexually active: - Gonorrhea/Chlamydia (females) - Contraception - Prevention of sexually transmitted infections	If sexually active: - Gonorrhea/Chlamydia (females) - Contraception
Selective Screening and Counseling (Individuals at increased risk)	HIV Syphilis Hepatitis B	HIV Syphilis Hepatitis B	HIV Syphilis Hepatitis B Cholesterol Prevention of sexually transmitted infections
Immunizations	Meningococcal Influenza (annual) Tdap HPV	Meningococcal Influenza (annual)	Influenza (annual) Td (q 10 years)
Discussion topics	Other substance use Physical activity Healthy eating Screen time School performance Dental hygiene General safety (e.g., seat belts, bike helmets)	Other substance use Physical activity Healthy eating Screen time School performance Dental hygiene General safety	Other substance use Physical activity Healthy eating Screen time Reproductive life plan Dental hygiene General safety

sources: Centers for Disease Control and Prevention immunization schedules. cdc.gov/vaccines/schedules/index.html | AAFP Clinical Recommendations for Prevention Services. aafp.org/patient-care.html \*Recommendations apply to non-pregnant adolescents and young adults. This work was supported by an unrestricted grant from the Merck Foundation.