

Tobacco Use Prevention and Cessation Counseling

2014 Medicare Benefits

Health Status	Symptomatic Patient	Asymptomatic Patient
Qualifying Medicare Patient	<p>A patient who:</p> <ol style="list-style-type: none"> 1. Uses tobacco and <ol style="list-style-type: none"> a. Has been diagnosed with a disease or an adverse health effect that has been found by the U.S. Surgeon General to be linked to tobacco use; or b. Is taking a therapeutic agent for which the metabolism or dosing is affected by tobacco use (based on U.S. Food and Drug Administration [FDA]-approved information); and 2. Is competent and alert at the time that counseling is provided; and 3. Receives counseling from a qualified physician or other Medicare-recognized health care professional. 	<p>A patient who:</p> <ol style="list-style-type: none"> 1. Uses tobacco but does not have symptoms of tobacco-related disease; 2. Is competent and alert at the time that counseling is provided; and 3. Receives counseling from a qualified physician or other Medicare-recognized health care professional.
CPT Codes	<p>99406: Smoking and tobacco-use cessation counseling visit; intermediate, greater than 3 minutes, up to 10 minutes</p> <p>99407: Smoking and tobacco-use cessation counseling visit; intensive, greater than 10 minutes</p>	<p>G0436: Smoking and tobacco cessation counseling visit for the asymptomatic patient; intermediate, greater than 3 minutes, up to 10 minutes</p> <p>G0437: Smoking and tobacco cessation counseling visit for the asymptomatic patient; intensive, greater than 10 minutes</p>
ICD-9 Diagnosis Codes	<p>305.1: Nondependent tobacco-use disorder</p> <p>AND</p> <p>ICD-9 code of condition that is adversely affected by tobacco use, or condition for which treatment is adversely affected by tobacco use</p>	<p>305.1: Nondependent tobacco-use disorder</p> <p>OR</p> <p>V15.82: Personal history of tobacco use</p>

Health Status	Symptomatic Patient	Asymptomatic Patient
ICD-10 Diagnosis Codes	F17.200: Nicotine dependence, unspecified, uncomplicated F17.201: Nicotine dependence, unspecified, in remission F17.210: Nicotine dependence, cigarettes, uncomplicated F17.211: Nicotine dependence, cigarettes, in remission F17.220: Nicotine dependence, chewing tobacco, uncomplicated F17.221: Nicotine dependence, chewing tobacco, in remission F17.290: Nicotine dependence, other tobacco product, uncomplicated F17.291: Nicotine dependence, other tobacco product, in remission	F17.200: Nicotine dependence, unspecified, uncomplicated F17.201: Nicotine dependence, unspecified, in remission F17.210: Nicotine dependence, cigarettes, uncomplicated F17.211: Nicotine dependence, cigarettes, in remission F17.220: Nicotine dependence, chewing tobacco, uncomplicated F17.221: Nicotine dependence, chewing tobacco, in remission F17.290: Nicotine dependence, other tobacco product, uncomplicated F17.291: Nicotine dependence, other tobacco product, in remission OR Z87.891: Personal history of nicotine dependence
Patient's Out-of-Pocket Cost	Eligible beneficiaries are covered under Medicare Part B. Both the coinsurance and deductible apply.	Eligible beneficiaries are covered under Medicare Part B. Both the coinsurance and deductible are waived.
Frequency	2 cessation attempts per 12-month period; maximum of 4 intermediate or intensive sessions per attempt (i.e., up to 8 sessions per 12-month period)	2 cessation attempts per 12-month period; maximum of 4 intermediate or intensive sessions per attempt (i.e., up to 8 sessions per 12-month period)

Developed by American Academy of Family Physicians (AAFP) Practice Advancement staff. Copyright © 2014 American Academy of Family Physicians. Physicians may photocopy or adapt for use in their practices; all other rights reserved. To seek permissions, contact copyrights@aafp.org.