



Body System: Cardiovascular		
Session Topic: Adult and Elderly Hypertension		
Educational Format		Faculty Expertise Required
REQUIRED	Interactive Lecture	Expertise in the field of study. Experience teaching in the field of study is desired. Preferred experience with audience response systems (ARS). Utilizing polling questions and engaging the learners in Q&A during the final 15 minutes of the session are required.
OPTIONAL	Problem-Based Learning (PBL)	Expertise teaching highly interactive, small group learning environments. Case-based, with experience developing and teaching case scenarios for simulation labs preferred. Other workshop-oriented designs may be accommodated. A typical PBL room is set for 50-100 participants, with 7-8 each per round table. <u>Please describe your interest and plan for teaching a PBL on your proposal form.</u>
Professional Practice Gap	Learning Objective(s) that will close the gap and meet the need	Outcome Being Measured
<ul style="list-style-type: none"> Family physicians may need additional training on effective counseling methods for patients to encourage healthy behavior changes to reduce their risk for developing hypertension or prehypertension. Family physicians need additional training on guideline recommendations for diagnosing and appropriately treating patients with hypertension using a stepwise approach to achieve stable tight control of blood pressure. Family physicians should recognize barriers to care that may prevent some patients from making appropriate health decisions, and they should understand ways to address them and/or offer alternative options. Family physicians should also be aware that blacks and Hispanics may have higher risk of developing 	<ol style="list-style-type: none"> Evaluate current management of hypertension in adult patients, as compared to current JNC 8 guidelines. Counsel patients on how to make healthy behavior changes to reduce their risk for developing hypertension or prehypertension. Address barriers to care among patients in your practice, especially elderly and minority patients, and identify or develop tools to help address hypertension. Prepare treatment regimens of antihypertensive medications and tools with an emphasis on patient adherence. 	Learners will submit written commitment to change statements on the session evaluation, indicating how they plan to implement presented practice recommendations.



<p>hypertension, and that hypertension in these populations could be more severe, requiring enhanced physician vigilance and aggressive management.</p> <ul style="list-style-type: none"> • Family physicians need to be able to apply current evidence-based guidelines to the management of elderly patients with hypertension. • While patients who have poorly controlled hypertension do not necessarily have “resistant hypertension,” family physicians should still be prepared to offer treatment regimens of antihypertensive medications and tools to improve patient adherence. • New JNC-8 guidelines have been released & need to be translated into practice 		
ACGME Core Competencies Addressed (select all that apply)		
X	Medical Knowledge	Patient Care
	Interpersonal and Communication Skills	Practice-Based Learning and Improvement
	Professionalism	Systems-Based Practice
Faculty Instructional Goals		
<p>Faculty play a vital role in assisting the AAFP to achieve its mission by providing high-quality, innovative education for physicians, residents and medical students that will encompass the art, science, evidence and socio-economics of family medicine and to support the pursuit of lifelong learning. By achieving the instructional goals provided, faculty will facilitate the application of new knowledge and skills gained by learners to practice, so that they may optimize care provided to their patients.</p> <ul style="list-style-type: none"> • Provide up to 3 evidence-based recommended practice changes that can be immediately implemented, at the conclusion of the session; including SORT taxonomy & reference citations • Facilitate learner engagement during the session • Address related practice barriers to foster optimal patient management • Provide recommended journal resources and tools, during the session, from the American Family Physician (AFP), Family Practice Management (FPM), and Familydoctor.org patient resources; those listed in the <u>References</u> section below are a good place to start <ul style="list-style-type: none"> ○ Visit http://www.aafp.org/journals for additional resources ○ Visit http://familydoctor.org for patient education and resources 		



- Provide specific examples of pharmacologic treatment for prehypertension/hypertension in difficult/challenging situations
- Provide specific strategies to manage/treat elderly patients, focusing on adherence and participation in rehabilitation programs
- Provides specific examples illustrating key differences in new guidelines compared to the laster iteration.

Needs Assessment

Hypertension is the most common condition for which patients make physician office visits. According to the CDC's 2010 *National Ambulatory Medical Care Survey* (NAMCS), hypertension was the top-ranked medical diagnosis by physicians at office visits; essential hypertension was the most commonly diagnosed condition seen at an office visit.¹

In addition, the prevalence of hypertension is significantly higher in blacks than in whites—39% compared to 29%, and uncontrolled hypertension may cause up to one-quarter of all deaths among black adults. Blacks also experience earlier onset of hypertension, their disease is more aggressive, and it is more difficult to treat and severe, especially when considering organ damage (e.g., renal failure).^{2,3} The prevalence of hypertension in Hispanics is comparable to or lower than that of whites, but the rates have been increasing, and Hispanics who have hypertension are less likely to have their blood pressure treated and controlled than are whites or blacks.³ The causes of these disparities are not completely understood, though differences in access to care, clinician management, hypertension severity and patient adherence to treatment all may play roles.² Family physicians should be aware of these disparities, as well as their own potential bias when treating minority patients, and pay close attention to any signs or symptoms of hypertension.⁴

Data from a recent American Academy of Family Physicians (AAFP) CME Needs Assessment survey indicate that family physicians have significant knowledge gaps with regard to managing adult hypertension, patient-adherence/shared decision making, and cardiovascular pharmacology.⁵ More specifically, CME outcomes data from 2012-2014 AAFP Assembly: *Adult Hypertension* sessions suggest that physicians need more continuing medical education with regard to integration of recommendations from current guidelines into practice; health coaching and lifestyle modifications; monitoring therapy adherence; and evidence-based prescribing practices.^{6,7}

Elderly patients have significant morbidity and mortality from coronary heart disease (CHD), with 84 percent of persons 65 years or older dying from this disease; however, medical therapies for hypertension have shown to have a positive effect on cardiovascular morbidity and mortality rates in older patients.⁸ This is further validated by Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 8) and summarized by key points of the report summarized below:^{9,10}

- In the general population ages 60 and older, pharmacologic treatment to lower BP should be initiated at a systolic blood pressure (SBP) of 150 mmHg or higher or a diastolic blood pressure (DBP) of 90 mmHg or higher. Patients should be treated to a goal SBP lower than 150 mmHg and a goal DBP lower than 90 mmHg. If treatment results in lower achieved SBP and is not associated with adverse effects, treatment does not need to be adjusted.



- In the general population younger than age 60, initiate pharmacologic treatment at a DBP of 90 mmHg or higher or an SBP of 140 mmHg or higher and treat to goals below these respective thresholds.
- In the population ages 18 years or older with diabetes or CKD, initiate pharmacologic treatment at an SBP of 140 mmHg or higher or a DBP of 90 mmHg or higher and treat to goals below these respective thresholds.
- In the general nonblack population, including those with diabetes, initial treatment should include a thiazide-type diuretic, calcium channel blocker (CCB), angiotensin-converting enzyme (ACE) inhibitor or angiotensin receptor blocker (ARB).
- In the general black population, including those with diabetes, initial treatment should include a thiazide-type diuretic or a CCB.
- In the population ages 18 or older with CKD and hypertension, initial (or add-on) treatment should include an ACE inhibitor or an ARB to improve kidney outcomes. This applies to all patients in this population regardless of race or diabetes status.
- Finally, the main objective of hypertension treatment is to attain and maintain goal BP. If goal BP is not reached within a month of initiating treatment, increase the dose of the initial drug or add a second drug from one of these four classes. The clinician should continue to assess BP and adjust the treatment regimen until goal BP is reached. If goal BP cannot be reached with two drugs, add and titrate a third drug from the list provided.

Although most physicians routinely check blood pressure as part of regular office visits, hypertension remains a challenging condition for both patients and physicians. The American Heart Association (AHA) reports that interventions targeting weight loss, physical activity and dietary modification can result in behavior change that, if sustained over time, can reduce morbidity and mortality associated with CVD.¹¹ However, because such practices take time to develop and adhere to, some patients may still require the use of medications and treatments, which family physicians can prescribe and monitor, to control risk factors (such as gender, age, family history and race/ethnicity) that cannot be controlled, or in cases in which hypertension is present with comorbidities such as hyperlipidemia and diabetes.

Most patients with hypertension have no clear etiology and are classified as having essential hypertension; however, 5 to 10 percent have secondary hypertension and may require specific imaging modalities to identify the secondary cause.¹² The most common cause of secondary hypertension is renal vascular hypertension, and is potentially curable if identified and treated properly.¹³ Data from the 2012 AAFP CME Needs Assessment Survey indicates that family physicians require additional education and training in the selection and use of imaging modalities.⁵ Additionally, some patients are diagnosed with “resistant hypertension,” in which high blood pressure persists despite patient adherence to medications, and other patients are at increased risk of cardiovascular diseases due to chronic conditions – such as diabetes – for which blood pressure control is a critical component of their care.¹² As a result, family physicians must be prepared to treat all different forms of hypertension in multiple types of patient encounters. According to the ACCF/AHA/PCPI Hypertension Performance Measurement Set, only 65% of patients with hypertension receive the recommended quality care outlined in current evidence-based guidelines.¹⁴ This PCPI measurement set outlines the desired hypertension outcomes as a reduction in mortality, the reduction in cardiovascular and renal morbidity, the reduction of hospitalization, improved adoption of lifestyle modifications, the reduction in patient harm, and improved patient understanding of/adherence to the prescribed treatment plan. Family physicians integrate the AMA PCPI Hypertension Measures into practice to improve overall patient care for



patients with hypertension. Adherence to evidence-based guidelines alone is insufficient in providing optimal patient care. In addition to being up to date on current evidence-based guidelines, family physicians also require tools and resources for evaluating hypertension.¹⁵

In order for family physicians to understand what treatment options are available – and be able to refer patients to a specialist, if necessary – they must have knowledge and understanding of the various elements associated with cardiovascular conditions (not just hypertension), their comorbidities and their risks. Consistency in treatment, follow-up and adherence to guidelines offered by a family physician may be a patient’s best opportunity to decrease the incidence of cardiovascular conditions and improve his or her quality of life. Providing family physicians with appropriate education and training on how to screen, diagnose, assess and treat patients with various cardiovascular diseases will help to decrease or eliminate practice gaps and lead to improved patient care.

Physicians can improve blood pressure control by using medical assistants and health education specialists in a team-based approach to patient care; however, roles, responsibilities, evaluation and communication plans should be carefully planned and established to ensure optimal effectiveness and buy-in.¹⁶⁻¹⁹

Resources: Evidence-Based Practice Recommendations/Guidelines/Performance Measures

- Screening for high blood pressure²⁰
- A tool for evaluating hypertension¹⁵
- Secondary prevention of coronary heart disease in elderly patients⁸
- Evaluation and management of the patient with difficult-to-control or resistant hypertension¹²
- Radiologic evaluation of suspected renovascular hypertension¹³
- AAFP/U.S. Preventive Services Task Force. Screening for High Blood Pressure: Reaffirmation Recommendation Statement.²¹
- AMA/ACCF/AHA PCPI Hypertension Performance Measurement Set¹⁴
- Guidelines for the primary prevention of stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association²²
- USPSTF: Behavioral Counseling Interventions to Promote a Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults²³
- 2014 evidence-based guideline for the management of high blood pressure in adults: Report from the panel members appointed to the eighth joint national committee (jnc 8)⁹
- How to document and code for hypertensive diseases in ICD-10²⁴
- Health coaching for patients with chronic illness²⁵
- Engaging Patients in Collaborative Care Plans²⁶
- Encouraging patients to change unhealthy behaviors with motivational interviewing²⁷
- Medication adherence: we didn't ask and they didn't tell²⁸
- Adding health education specialists to your practice¹⁹
- FamilyDoctor.org. High Blood Pressure | Overview (patient resource)²⁹



References

1. Centers for Disease Control and Prevention (CDC). National Ambulatory Medical Care Survey. In: Ambulatory and Hospital Care Statistics Branch, ed2010.
2. Fiscella K, Holt K. Racial Disparity in Hypertension Control: Tallying the Death Toll. *The Annals of Family Medicine*. November 1, 2008 2008;6(6):497-502.
3. Morenoff JD, House JS, Hansen BB, Williams DR, Kaplan GA, Hunte HE. Understanding social disparities in hypertension prevalence, awareness, treatment, and control: The role of neighborhood context. *Social Science & Medicine*. 2007;65(9):1853-1866.
4. Blair IV, Steiner JF, Fairclough DL, et al. Clinicians' Implicit Ethnic/Racial Bias and Perceptions of Care Among Black and Latino Patients. *The Annals of Family Medicine*. January 1, 2013 2013;11(1):43-52.
5. AAFP. 2012 CME Needs Assessment: Clinical Topics. American Academy of Family Physicians; 2012.
6. American Academy of Family Physicians (AAFP). 2012 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2012.
7. American Academy of Family Physicians (AAFP). 2013 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2013.
8. Hanna IR, Wenger NK. Secondary prevention of coronary heart disease in elderly patients. *American family physician*. Jun 15 2005;71(12):2289-2296.
9. James PA, Oparil S, Carter BL, et al. 2014 evidence-based guideline for the management of high blood pressure in adults: Report from the panel members appointed to the eighth joint national committee (jnc 8). *JAMA : the journal of the American Medical Association*. 2014;311(5):507-520.
10. Borgmeyer C. 2014 Hypertension Guideline Stands to Simplify Treatment, Says Expert: AAFP to Begin Rigorous Process of Reviewing for Possible Endorsement *AAFP News Now*. 2014. <http://www.aafp.org/news/health-of-the-public/20131218hypertensiongdln.html>. Accessed August 2014.
11. Artinian NT, Fletcher GF, Mozaffarian D, et al. Interventions to promote physical activity and dietary lifestyle changes for cardiovascular risk factor reduction in adults: a scientific statement from the American Heart Association. *Circulation*. Jul 27 2010;122(4):406-441.
12. Viera AJ, Hinderliter AL. Evaluation and management of the patient with difficult-to-control or resistant hypertension. *American family physician*. May 15 2009;79(10):863-869.
13. Hartman RP, Kawashima A. Radiologic evaluation of suspected renovascular hypertension. *American family physician*. Aug 1 2009;80(3):273-279.
14. Association AM, Cardiology ACo, Association AH. Hypertension Performance Measurement Set. *PCPI Approved Quality Measures 2011*; <http://www.ama-assn.org/apps/listserv/x-check/qmeasure.cgi?submit=PCPI>. Accessed May, 2013.
15. Ebell MH. A tool for evaluating hypertension. *Family practice management*. Mar 2004;11(3):79-81.
16. Margolius D, Wong J, Goldman ML, Rouse-Iniguez J, Bodenheimer T. Delegating responsibility from clinicians to nonprofessional personnel: the example of hypertension



- control. *Journal of the American Board of Family Medicine : JABFM*. Mar-Apr 2012;25(2):209-215.
17. Martinez M, Bigney J, Jernigan J. A feedback tool to improve physician-medical assistant communication. *Family practice management*. May-Jun 2014;21(3):5-9.
 18. Naughton D, Adelman AM, Bricker P, Miller-Day M, Gabbay R. Envisioning new roles for medical assistants: strategies from patient-centered medical homes. *Family practice management*. Mar-Apr 2013;20(2):7-12.
 19. Chambliss ML, Lineberry S, Evans WM, Bibeau DL. Adding health education specialists to your practice. *Family practice management*. Mar-Apr 2014;21(2):10-15.
 20. Lin KW, Moutsiakis D. Screening for high blood pressure. *Am Fam Physician*. Jun 15 2009;79(12):1093-1094.
 21. American Academy of Family Physicians (AAFP). Clinical Preventive Services Recommendations. 2012; <http://www.aafp.org/online/en/home/clinical/exam.html>. Accessed May, 2013.
 22. Meschia JF, Bushnell C, Boden-Albala B, et al. Guidelines for the primary prevention of stroke: a statement for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke; a journal of cerebral circulation*. Dec 2014;45(12):3754-3832.
 23. Moyer VA, on behalf of the USPSTF. Behavioral Counseling Interventions to Promote a Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults: U.S. Preventive Services Task Force Recommendation Statement. *Annals of internal medicine*. Jun 26 2012.
 24. Beckman KD. How to document and code for hypertensive diseases in ICD-10. *Family practice management*. Mar-Apr 2014;21(2):5-9.
 25. Bennett HD, Coleman EA, Parry C, Bodenheimer T, Chen EH. Health coaching for patients with chronic illness. *Family practice management*. Sep-Oct 2010;17(5):24-29.
 26. Mauksch L, Safford B. Engaging Patients in Collaborative Care Plans. *Family practice management*. 2013;20(3):35-39.
 27. Stewart EE, Fox CH. Encouraging patients to change unhealthy behaviors with motivational interviewing. *Family practice management*. May-Jun 2011;18(3):21-25.
 28. Brown M, Sinsky CA. Medication adherence: we didn't ask and they didn't tell. *Family practice management*. Mar-Apr 2013;20(2):25-30.
 29. FamilyDoctor.org. High Blood Pressure | Overview. 1996; <http://familydoctor.org/familydoctor/en/diseases-conditions/high-blood-pressure.html>. Accessed August, 2013.