



Body System: Cardiovascular			
Session Topic: Heart Failure			
Educational Format		Faculty Expertise Required	
REQUIRED	Interactive Lecture	Expertise in the field of study. Experience teaching in the field of study is desired. Preferred experience with audience response systems (ARS). Utilizing polling questions and engaging the learners in Q&A during the final 15 minutes of the session are required.	
OPTIONAL	Problem-Based Learning (PBL)	Expertise teaching highly interactive, small group learning environments. Case-based, with experience developing and teaching case scenarios for simulation labs preferred. Other workshop-oriented designs may be accommodated. A typical PBL room is set for 50-100 participants, with 7-8 each per round table. <u>Please describe your interest and plan for teaching a PBL on your proposal form.</u>	
Professional Practice Gap		Learning Objective(s) that will close the gap and meet the need	Outcome Being Measured
<ul style="list-style-type: none"> Family physicians have knowledge gaps with regard to evaluating and managing heart failure, analyzing electrocardiography, performing cardiovascular physical examinations, effective use of cardiac rehabilitation, and cardiovascular pharmacology. Family physicians have knowledge gaps with regard to utilization of staging criteria for classifying at risk patients, selecting appropriate and effective treatment medications, and consistent adherence to current practice guidelines. Diagnostic errors associated with heart failure in common in primary care. Data to suggest that physicians following evidence-based clinical practice guidelines are less likely to underuse beta blockers in congestive 		<ol style="list-style-type: none"> Use current ACC/AHA guidelines to evaluate and classify patients with suspected heart failure. Use evidence-based criteria to identify appropriate diagnostic imaging and laboratory tests to identify causes or precipitating factors. Establish guideline-directed medical therapy (GDMT), as indicated by the initial evaluation. Develop collaborative care plans with patients that emphasize making healthy lifestyle changes and adherence to prescribed therapies. 	Learners will submit written commitment to change statements on the session evaluation, indicating how they plan to implement presented practice recommendations.



heart failure treatment • Physicians require training and strategies to improve documentation and decision making, improve the effectiveness of health coaching, and improve medication adherence.		
---	--	--

ACGME Core Competencies Addressed (select all that apply)

X	Medical Knowledge	Patient Care
X	Interpersonal and Communication Skills	Practice-Based Learning and Improvement
	Professionalism	Systems-Based Practice

Faculty Instructional Goals

Faculty play a vital role in assisting the AAFP to achieve its mission by providing high-quality, innovative education for physicians, residents and medical students that will encompass the art, science, evidence and socio-economics of family medicine and to support the pursuit of lifelong learning. By achieving the instructional goals provided, faculty will facilitate the application of new knowledge and skills gained by learners to practice, so that they may optimize care provided to their patients.

- Provide up to 3 evidence-based recommended practice changes that can be immediately implemented, at the conclusion of the session; including SORT taxonomy & reference citations
- Facilitate learner engagement during the session
- Address related practice barriers to foster optimal patient management
- Provide recommended journal resources and tools, during the session, from the American Family Physician (AFP), Family Practice Management (FPM), and Familydoctor.org patient resources; those listed in the References section below are a good place to start
 - Visit <http://www.aafp.org/journals> for additional resources
 - Visit <http://familydoctor.org> for patient education and resources
- Provide strategies for using current ACC/AHA guidelines to evaluate and classify patients with suspected heart failure.
- Provide recommendations on the use of evidence-based criteria to identify appropriate diagnostic imaging and laboratory tests to identify causes or precipitating factors.
- Provide recommendations for establishing guideline-directed medical therapy (GDMT), as indicated by the initial evaluation, with an emphasis on consistent guideline adherence.
- Provide strategies and resources to develop collaborative care plans with patients that emphasize making healthy lifestyle changes and adherence to prescribed therapies.
- Provide recommendations regarding guidelines for Medicare reimbursement.
- Provide recommendations to maximize office efficiency and guideline adherence to the diagnosis and management of
- Provide an overview of newly available treatments, including efficacy, safety, contraindications, and cost/benefit relative to existing treatments.
- Provide instructions regarding the incorporation and use of the PCMH/ACO/Primary Care Core Measure Set into practice.



Needs Assessment

Nearly 5.1 million people in the United States have heart failure (HF), over 23 million worldwide, and was the underlying cause of death in 57,757 people in the U.S. in 2010.¹⁻³ Congestive heart failure (non-hypertensive) is the leading condition resulting in hospital readmissions among Medicare patients (aged 65 and older), the seventh leading cause for hospital readmissions among Medicaid patients (aged 18-64), and the eighth leading cause among uninsured patients (aged 18-64 years).⁴ Recent studies from the Agency for Healthcare Research and Quality (AHRQ), demonstrate that home visiting programs and multidisciplinary clinic interventions for adult patients with heart failure reduced all-cause readmissions and mortality over three to six months. (Strength of recommendation [SOR]: A, based on consistent, good-quality patient-oriented evidence.) Structured telephone support reduced heart failure-specific readmissions and mortality over three to six months. (SOR: A, based on consistent, good-quality patient-oriented evidence.) However, structured telephone support did not reduce all-cause readmissions over a similar period. (SOR: B, based on inconsistent or limited-quality patient-oriented evidence.)⁵

Multicomponent interventions such as home visiting programs and multidisciplinary clinic interventions for heart failure reduced all-cause readmissions and mortality over three to six months. Key components of these interventions included heart failure education emphasizing self-care, heart failure pharmacotherapy emphasizing adherence, face-to-face contact after hospital discharge, mechanisms for post-discharge medication adjustment, and streamlined mechanisms to contact care delivery personnel (e.g., a patient hotline). These higher-intensity interventions were delivered by teams of clinicians.

Data from a recent American Academy of Family Physicians (AAFP) CME Needs Assessment survey indicate that family physicians have knowledge gaps with regard to evaluating and managing heart failure, analyzing electrocardiography, performing cardiovascular physical examinations, effective use of cardiac rehabilitation, and cardiovascular pharmacology.⁶ More specifically, CME outcomes data from 2012-2015 AAFP FMX (formerly Assembly) *Heart Failure* sessions suggest that physicians have knowledge and practice gaps with regard to utilization of staging criteria for classifying at risk patients (e.g. NYHA & ACC); selecting appropriate and effective treatment medications (e.g. extended-release vs. immediate release Metoprolol, appropriate use of beta blockers, appropriate use of statins, etc.); better awareness of drug interactions; effective lifestyle modification counseling; and consistent adherence to current practice guidelines.⁷⁻⁹

Diagnostic errors associated with heart failure in common in primary care; particularly, process breakdowns frequently involved the patient-practitioner clinical encounter (78.9%) but are also related to referrals (19.5%), patient-related factors (16.3%), follow-up and tracking of diagnostic information (14.7%), and performance and interpretation of diagnostic tests (13.6%).¹⁰ CME outcomes data from the American Academy of Family Physicians (AAFP) Advanced Cardiac Testing CME Program, as well as data from a national study of the National Research Network (NRN), indicate that cardiac tests ordered by family physicians vary regionally, and that physician-patient communication gaps, as well as coordination of care gaps exists and are barriers to optimal patient care. It is well established that interpretation of ECGs is within the scope of family medicine, and that the diagnosis and management of cardiovascular disorders is



routinely taught in family medicine residency programs.¹¹ However, studies suggest that family practice residents have considerable deficiencies in ECG interpretation skills.¹² Documentation of diagnostic imaging is also inconsistent among many primary care providers; therefore, family physicians should be familiar with ACR practice guidelines for communication of diagnostic imaging findings.^{13,14} Family physicians can also improve clinical decision making by utilizing clinical decision support tools and documenting decision making.^{15,16} Physicians require training and strategies to improve documentation and decision making, improve the effectiveness of health coaching, and improve medication adherence.¹⁵⁻²¹

The goal for using any cardiac imaging or stress test should be to maximize usefulness without unnecessary cost or harm to the patient, however, some studies suggest that 30% to 50% of cardiovascular imaging tests are partially or totally inappropriate.²² Physicians must always consider the safe and effective use of diagnostic imaging, as risks of diagnostic imaging include cancer from radiation exposure and nephrogenic systemic fibrosis, and the overuse of health care services leads to poor quality and high costs.²³⁻²⁶ Physicians should consult Choosing Wisely[®] recommendations when considering cardiac imaging tests:²⁷⁻³³

Physicians may improve their care of patients with HF by engaging in continuing medical education that provides practical integration of current evidence-based guidelines and recommendations into their standards of care, including, but not limited to the following:³⁴⁻³⁶

- The initial evaluation of patients with suspected heart failure should include a history and physical examination, laboratory assessment, chest radiography, and electrocardiography. Echocardiography can confirm the diagnosis.
- A displaced cardiac apex, a third heart sound, and chest radiography findings of pulmonary venous congestion or interstitial edema are good predictors to rule in the diagnosis of heart failure.
- Systolic heart failure can be effectively ruled out with a normal B-type natriuretic peptide or N-terminal pro-B-type natriuretic peptide level.
- Systolic heart failure can be effectively ruled out when the Framingham criteria are not met.
- Treatment of stage A heart failure should focus on reducing modifiable risk factors, including management of hypertension and hyperlipidemia.
- To prevent symptomatic heart failure, ACE inhibitors and beta blockers should be used in all patients with stage B or C heart failure who have a reduced ejection fraction.
- Patients with stage C heart failure and fluid retention should be treated with diuretics in addition to ACE inhibitors and beta blockers.
- ACA/AHA guidelines for lifestyle management to reduce cardiovascular risk through counseling, management, and prevention

There is data to suggest that physicians following evidence-based clinical practice guidelines are less likely to underuse beta blockers in congestive heart failure treatment.²⁵ This is significant, as recent research suggests that beta blocker therapy reduces mortality in patients with heart failure with reduced ejection fraction.³⁷ These recommendations are provided only as assistance for physicians making clinical decisions regarding the care of their patients. As such, they cannot substitute for the individual judgment brought to each clinical situation by the patient's family



physician. As with all clinical reference resources, they reflect the best understanding of the science of medicine at the time of publication, but they should be used with the clear understanding that continued research may result in new knowledge and recommendations. These recommendations are only one element in the complex process of improving the health of America. To be effective, the recommendations must be implemented. As such, physicians require continuing medical education to assist them with making decisions about specific clinical considerations.

The American Academy of Family Physicians Academy has participated in the Core Measures Collaborative (the Collaborative) convened by America's Health Insurance Plans (AHIP) since August 2014. The Collaborative is a multi-stakeholder effort working to define core measure sets of various specialties promoting alignment and harmonization of measure use and collection across both public and private payers.

Participants in the Collaborative included Centers for Medicare and Medicaid Services (CMS), the National Quality Forum (NQF), private payers, provider organizations, employers, and patient and consumer groups. This effort exists to decrease physician burden by reducing variability in measure selection, specifications and implementation—making quality measurement more useful and meaningful for consumers, employers, as well as public and private clinicians.

With significant AAFP input, a PCMH/ACO/Primary Care Core Measure Set has been developed for primary care. The goal of this set is to decrease burden and allow for more congruence between payer reporting programs.³⁸

Resources: Evidence-Based Practice Recommendations/Guidelines/Performance Measures

- Diagnosis and evaluation of heart failure³⁴
- 2013 ACCF/AHA guideline for the management of heart failure³⁵
- AHA/ACC guideline on lifestyle management to reduce cardiovascular risk³⁶
- NICE Acute heart failure: diagnosing and managing acute heart failure in adults³⁹
- Appropriate and safe use of diagnostic imaging²³
- ACR practice guideline for communication of diagnostic imaging findings¹⁴
- Clinical decision support: using technology to identify patients' unmet needs¹⁵
- Thinking on paper: documenting decision making¹⁶
- Medication adherence: we didn't ask and they didn't tell¹⁷
- Health Coaching: Teaching Patients to Fish¹⁸
- An organized approach to chronic disease care¹⁹
- The integrated summary: a documentation tool to improve patient care.²⁰
- Engaging Patients in Collaborative Care Plans²¹
- Heart Failure | Overview (patient education)⁴⁰

References



1. Go AS, Mozaffarian D, Roger VL, et al. Executive summary: heart disease and stroke statistics--2014 update: a report from the American Heart Association. *Circulation*. Jan 21 2014;129(3):399-410.
2. Bui AL, Horwich TB, Fonarow GC. Epidemiology and risk profile of heart failure. *Nature reviews. Cardiology*. Jan 2011;8(1):30-41.
3. Centers for Disease Control and Prevention. Heart Failure Fact Sheet. 2013; http://www.cdc.gov/dhdsdp/data_statistics/fact_sheets/fs_heart_failure.htm. Accessed August, 2014.
4. Hines AL, Barrett ML, Jiang HJ, Steiner CA. Conditions With the Largest Number of Adult Hospital Readmissions by Payer, 2011: Statistical Brief #172. In: AHRQ, ed. *Healthcare Cost and Utilization Project (HCUP) Statistical Briefs*. Rockville MD 2014.
5. Guirguis-Blake J. Transitional Care Interventions to Prevent Readmissions for Patients with Heart Failure. *American family physician*. Mar 1 2016;93(5):401-403.
6. AAFP. 2012 CME Needs Assessment: Clinical Topics. American Academy of Family Physicians; 2012.
7. American Academy of Family Physicians (AAFP). 2012 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2012.
8. American Academy of Family Physicians (AAFP). 2013 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2013.
9. American Academy of Family Physicians (AAFP). AAFP FMX CME Outcomes Report. Leawood KS: AAFP; 2015.
10. Singh H, Giardina TD, Meyer AN, Forjuoh SN, Reis MD, Thomas EJ. Types and origins of diagnostic errors in primary care settings. *JAMA internal medicine*. Mar 25 2013;173(6):418-425.
11. American Academy of Family Physicians (AAFP). AAFP Policies: Electrocardiograms, Family Physician Interpretation of. 2001; <http://www.aafp.org/about/policies/all/electrocardiograms.html>. Accessed June, 2013.
12. Boltri JM, Hash RB, Vogel RL. Are family practice residents able to interpret electrocardiograms? *Advances in health sciences education : theory and practice*. 2003;8(2):149-153.
13. Weiner SN. Radiology by nonradiologists: is report documentation adequate? *The American journal of managed care*. Dec 2005;11(12):781-785.
14. American College of Radiology (ACR). ACR practice guideline for communication of diagnostic imaging findings. 2010; <http://www.guidelines.gov/content.aspx?id=32541>. Accessed 6/14/2013.
15. McLeod W, Eidus R, Stewart EE. Clinical decision support: using technology to identify patients' unmet needs. *Family practice management*. Mar-Apr 2012;19(2):22-28.
16. Edsall RL, Moore KJ. Thinking on paper: documenting decision making. *Family practice management*. Jul-Aug 2010;17(4):10-15.
17. Brown M, Sinsky CA. Medication adherence: we didn't ask and they didn't tell. *Family practice management*. Mar-Apr 2013;20(2):25-30.
18. Ghorob A. Health Coaching: Teaching Patients to Fish. *Family practice management*. 2013;20(3):40-42.
19. Lyon RK, Slawson J. An organized approach to chronic disease care. *Family practice management*. May-Jun 2011;18(3):27-31.



20. Stelman MA. The integrated summary: a documentation tool to improve patient care. *Family practice management*. Apr 2003;10(4):33-39.
21. Mauksch L, Safford B. Engaging Patients in Collaborative Care Plans. *Family practice management*. 2013;20(3):35-39.
22. Picano E. The risks of inappropriateness in cardiac imaging. *International journal of environmental research and public health*. May 2009;6(5):1649-1664.
23. Crownover BK, Bepko JL. Appropriate and safe use of diagnostic imaging. *American family physician*. Apr 1 2013;87(7):494-501.
24. Korenstein D, Falk R, Howell EA, Bishop T, Keyhani S. Overuse of health care services in the United States: an understudied problem. *Archives of internal medicine*. Jan 23 2012;172(2):171-178.
25. Kale MS, Bishop TF, Federman AD, Keyhani S. Trends in the overuse of ambulatory health care services in the United States. *JAMA internal medicine*. Jan 28 2013;173(2):142-148.
26. Kale MS, Bishop TF, Federman AD, Keyhani S. "Top 5" lists top \$5 billion. *Archives of internal medicine*. Nov 14 2011;171(20):1856-1858.
27. American Board of Internal Medicine (ABIM). Choosing Wisely: EKGs and exercise stress tests - When you need them for heart disease - and when you don't. 2012; <http://www.choosingwisely.org/doctor-patient-lists/ekgs-and-exercise-stress-tests/>. Accessed June, 2013.
28. Rao VM, Levin DC. The Overuse of Diagnostic Imaging and the Choosing Wisely Initiative. *Annals of internal medicine*. 2012;157(8):574-576.
29. American Board of Internal Medicine (ABIM). Choosing Wisely: Lists. 2013; <http://www.choosingwisely.org/doctor-patient-lists/>. Accessed July, 2013.
30. American Academy of Family Physicians (AAFP). Don't use coronary CT angiography in high-risk emergency department patients presenting with acute chest pain. note: Risk defined by the Thrombolysis In Myocardial Infarction risk score for unstable angina/acute coronary syndromes. *Choosing Wisely* 2014; <http://www.aafp.org/afp/recommendations/viewRecommendation.htm?recommendationId=15>. Accessed August, 2014.
31. American College of Cardiology. Don't perform stress cardiac imaging or advanced noninvasive imaging in the initial evaluation of patients without cardiac symptoms unless high-risk markers are present. *Choosing Wisely* 2014; <http://www.aafp.org/afp/recommendations/viewRecommendation.htm?recommendationId=3>. Accessed August, 2014.
32. American Society of Nuclear Cardiology. Don't perform cardiac imaging as a preoperative assessment in patients scheduled to undergo low- or intermediate-risk noncardiac surgery. *Choosing Wisely* 2014; <http://www.aafp.org/afp/recommendations/viewRecommendation.htm?recommendationId=166>. Accessed August, 2014.
33. American Society of Nuclear Cardiology. Five Things Physicians and Patients Should Question. *Choosing Wisely* 2012; <http://www.choosingwisely.org/doctor-patient-lists/american-society-of-nuclear-cardiology/>. Accessed August, 2014.
34. King M, Kingery J, Casey B. Diagnosis and evaluation of heart failure. *American family physician*. Jun 15 2012;85(12):1161-1168.



35. Writing Committee M, Yancy CW, Jessup M, et al. 2013 ACCF/AHA guideline for the management of heart failure: a report of the American College of Cardiology Foundation/American Heart Association Task Force on practice guidelines. *Circulation*. Oct 15 2013;128(16):e240-327.
36. Eckel RH, Jakicic JM, Ard JD, et al. 2013 AHA/ACC guideline on lifestyle management to reduce cardiovascular risk: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Journal of the American College of Cardiology*. Jul 1 2014;63(25 Pt B):2960-2984.
37. Nielsen PB, Larsen TB, Gorst-Rasmussen A, Skjoth F, Lip GY. beta-Blockers in Atrial Fibrillation Patients With or Without Heart Failure: Association With Mortality in a Nationwide Cohort Study. *Circulation. Heart failure*. Feb 2016;9(2):e002597.
38. American Academy of Family Physicians (AAFP). PCMH/ACO/Primary Care Core Measure Set. 2016; <http://www.aafp.org/practice-management/improvement/measures.html>. Accessed May, 2016.
39. National Guideline Clearinghouse. Acute heart failure: diagnosing and managing acute heart failure in adults. <http://www.guideline.gov/content.aspx?id=48752&search=heart+failure>. Accessed 6/4/2015.
40. FamilyDoctor.org. Heart Failure | Overview. 2000; <http://familydoctor.org/familydoctor/en/diseases-conditions/heart-failure.html>. Accessed August, 2014.