



Body System: Neurologic		
Session Topic: Dementia and Alzheimer's Disease		
Educational Format		Faculty Expertise Required
REQUIRED	Interactive Lecture	Expertise in the field of study. Experience teaching in the field of study is desired. Preferred experience with audience response systems (ARS). Utilizing polling questions and engaging the learners in Q&A during the final 15 minutes of the session are required.
OPTIONAL	Problem-Based Learning (PBL)	Expertise teaching highly interactive, small group learning environments. Case-based, with experience developing and teaching case scenarios for simulation labs preferred. Other workshop-oriented designs may be accommodated. A typical PBL room is set for 50-100 participants, with 7-8 each per round table. <u>Please describe your interest and plan for teaching a PBL on your proposal form.</u>
Professional Practice Gap	Learning Objective(s) that will close the gap and meet the need	Outcome Being Measured
<ul style="list-style-type: none"> • Knowledge and competence gap related to screening and evaluation of symptomatic patients for cognitive decline. • Many primary care physicians view the treatment of dementia as time-consuming and not fitting well into the usual primary care setting. • Knowledge gap about resources and strategies to maintain or improve cognitive health in the prevention of dementia and Alzheimer's disease. • Knowledge and competence gap related to providing support and counseling to caregivers of dementia and Alzheimer's patients. • Knowledge and competence gap related to the consistent implementation and adherence to evidence-based guidelines for the management of dementia and Alzheimer's patients. 	<ol style="list-style-type: none"> 1. Use evidence-based guidelines to screen and evaluate patients who are symptomatic for cognitive decline for dementia. 2. Identify tools and resources available to the care team, caregivers, and patients about strategies to maintain or improve cognitive health. 3. Use evidence-based guidelines to formulate pharmacologic and non-pharmacologic therapies to help slow the progression of Alzheimer's. 4. Counsel patients and their family members on how to cope with neurologic disorders that result in the loss of cognitive functioning, such as Alzheimer's disease. 	Learners will submit written commitment to change statements on the session evaluation, indicating how they plan to implement presented practice recommendations.



2017 AAFP FMX Needs Assessment

<ul style="list-style-type: none"> New research showing a possible link between proton pump inhibitors (PPIs) and risk of dementia in older adults. 		
ACGME Core Competencies Addressed (select all that apply)		
X	Medical Knowledge	Patient Care
X	Interpersonal and Communication Skills	Practice-Based Learning and Improvement
	Professionalism	Systems-Based Practice
Faculty Instructional Goals		
<p>Faculty play a vital role in assisting the AAFP to achieve its mission by providing high-quality, innovative education for physicians, residents and medical students that will encompass the art, science, evidence and socio-economics of family medicine and to support the pursuit of lifelong learning. By achieving the instructional goals provided, faculty will facilitate the application of new knowledge and skills gained by learners to practice, so that they may optimize care provided to their patients.</p> <ul style="list-style-type: none"> Provide up to 3 evidence-based recommended practice changes that can be immediately implemented, at the conclusion of the session; including SORT taxonomy & reference citations Facilitate learner engagement during the session Address related practice barriers to foster optimal patient management Provide recommended journal resources and tools, during the session, from the American Family Physician (AFP), Family Practice Management (FPM), and Familydoctor.org patient resources; those listed in the <u>References</u> section below are a good place to start <ul style="list-style-type: none"> Visit http://www.aafp.org/journals for additional resources Visit http://familydoctor.org for patient education and resources Provide specific examples illustrating the application of evidence-based guidelines, as it applies to the screening, diagnosis, and management of dementia and Alzheimer’s. This should include recommendations regarding screening for depression. Provide specific examples of tools and strategies to help make the office visit more efficient, and to appropriately code for reimbursement Provide learners with practical strategies to utilize community-based resources as a way to support caregivers of dementia and Alzheimer’s patients Provide specific, evidence-based information about effective pharmacologic and non-pharmacologic therapies to lessen symptoms of Alzheimer’s Provide recommendations regarding the ADNI Alzheimer’s Disease Neuroimaging Initiative study. Provide recommendations regarding guidelines for Medicare reimbursement. Provide recommendations to maximize office efficiency and guideline adherence to the diagnosis and management of dementia and Alzheimer’s. Provide an overview of newly available treatments, including efficacy, safety, contraindications, and cost/benefit relative to existing treatments. 		

Needs Assessment



Dementia is one of the most common afflictions of the elderly, but it is important for patients and their families to understand that it is not a “normal” part of aging. Normal aging does not result in the loss of large numbers of neurons in the brain, which eventually leads to impaired cognition and functioning. While there are numerous disorders that can cause dementia, the most common of which is Alzheimer’s disease (AD), family physicians can rule out any other treatable conditions (such as depression or vitamin B₁₂ deficiency) that can cause similar symptoms.¹⁻³ Alzheimer’s disease (AD) is the most common neurological disorder and most common cause of dementia for people over the age of 65; it is currently estimated to affect more than 5 million Americans.⁴ Alzheimer’s disease causes nearly 75,000 deaths per year; it is currently the sixth-leading cause of death in the U.S. It is estimated that 41.8% of assisted living residents, 15.5% of residents of nursing homes, and 7.5% of patients under the care of home hospices have AD.⁵ The Alzheimer’s Association indicate that the most important risk factors that increase the risk of Alzheimer’s is age, family history, and heredity.⁴

Data from a recent American Academy of Family Physicians (AAFP) CME Needs Assessment survey indicate that family physicians have a statistically significant and meaningful knowledge and skill gap related to the management of dementia and Alzheimer disease.⁶ More specifically, CME outcomes data from 2012-2014 AAFP Assembly: *Dementia and Alzheimer Diseases* sessions suggests that physicians have knowledge and competence gaps with regard to cognitive evaluation of symptomatic patients; early testing for asymptomatic patients; appropriate use of diagnostic testing; support of caregivers; pharmacologic and non-pharmacologic therapies, including knowing when to initiate &/or stop medications; and screening for depression.⁷⁻⁹

Physicians often feel challenged in caring for dementia patients due to barriers such as lack of time in the office visit, reimbursement issues, lack of access to and poor communication with specialists, access to community services, and lack of interdisciplinary teams.^{10,11} The American Academy of Neurology (AAN), American Geriatric Society (AGS), and the American Medical Directors Association (AMDA) in the AMA PCPI Dementia Performance Measure Set has also identified gaps in physician performance and patient care:¹²

- On average, older patients with dementia received the recommended quality of care only about 35 percent of the time
- Some studies suggest that caregivers in general only adhere to evidence-based recommendations for the care of patients with dementia about 40 percent of the time
- Adherence to clinical practice guidelines was moderate to low at VA medical centers with regard to cognitive and depression screening, reporting of elder abuse, discussing care needs and decision making issues with patients’ family and implementing caregiver support practices

Additionally, the AAFP National Research Network conducted a study of active members to investigate the practice patterns, beliefs, concerns, and perceived need of family physicians that treat patients with dementia. This study identified a number of factors affecting the diagnosis of dementia and the delivery of dementia care included patient behavior challenges (aggressiveness, restlessness, paranoia, wandering); comorbidities (falls, delirium, adverse medication reactions, urinary incontinence); caregiver challenges (fatigue, planning for patient’s institutional placement, anger); and structural barriers (clinician time, time required for screening, limited



treatment options). Tools needed to provide enhanced dementia care included better assessment tools, community resources, and diagnostic and screening tools.¹³

Prevention is a key aspect of patient-centered care. Some primary care physicians report barriers to talking with patients about cognitive health such as lack of time, patient negative reactions to recommendations, and many physicians do not believe that strategies for cognitive health have sufficient evidence in their efficacy.¹⁴ As physicians transform their practices to provide patient-centered care, family physicians must remain up to date on current initiatives to maintain or improve cognitive health. On May 15, 2012, the U.S. Department of Health and Human Services released the National Alzheimer's Plan, with a specific goal of finding effective ways to prevent and treat Alzheimer's diseases and related dementias by 2025.¹⁵ Additionally, the Healthy Brain Initiative from the Division of Population Health, National Center for Chronic Prevention and Health Promotion provides resources for providers, caregivers, and patients for cognitive health.^{14,16} Physicians should also be aware of studies suggesting a link between dementia and benzodiazepine use, and understand evidence-based recommendations for counseling patients about risk reduction.¹⁷

The American Academy of Family Physicians (AAFP) and the U.S. Preventive Services Task Force (USPSTF) make no recommendation for or against routine screening for dementia in asymptomatic older adults; however, utilizing the MMSE for screening is recommended whenever cognitive impairment or deterioration is suspected in order to help make diagnostic and treatment decisions.^{18,19} In June 2014 the USPSTF updated its recommendation statement regarding screening for cognitive impairment as, "USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for cognitive impairment."^{20,21} As such, physicians should be familiar with clinical considerations, and be prepared to make appropriate individual adjustments.

The National Institute on Aging and the Alzheimer's Association has released guidelines for clinical diagnosis of dementia and mild cognitive impairment. Family physicians should use current evidence-based guidelines to evaluate patients with suspected dementia. Evidence-based recommendations for practice are outlined in the October 15, 2011 issue of *American Family Physician*.³ A summary of key recommendations are as follows:

- For patients with suspected dementia, quick screening tests, such as a verbal fluency test and the Mini-Cognitive Assessment Instrument, should be performed to determine whether further evaluation is warranted.
- Formal cognitive testing should be performed in patients with abnormal results on initial dementia screening.
- Routine blood work (i.e., complete blood count; complete metabolic panel; and measurement of thyroid-stimulating hormone, vitamin B12, folate, and calcium levels) should be performed in patients with suspected dementia.
- Additional testing (e.g., neuroimaging, cerebrospinal fluid analysis, human immunodeficiency virus testing, Lyme titer, and rapid plasma reagin test) can be performed in patients with suspected dementia and specific risk factors or symptoms.
- A second visit should include a Mini-Mental State Examination, Geriatric Depression Scale, and verbal fluency and clock drawing tests, if not previously completed



Physicians should also be familiar with the Montreal Cognitive Assessment (MoCA), and be able to determine whether the MoCA or the MMSE is the appropriate assessment, under the given situation.^{3,22,23}

A study from the AAFP National Research Network indicates that if screening and diagnostic testing for ECI/dementia as demonstrated in this project is going to be implemented in primary care offices, there must be:²⁴

- ways to implement this process without disrupting ordinary practice flow, including alternatives for its implementation that can be shown to work;
- reimbursable to the extent to which it will be worth the effort financially for the practice to implement systematically; and
- (Probably most importantly) effective treatments for ECI and dementia or demonstration of other benefits from early recognition of the problem.

Recent studies indicate that positron emission tomography (PET) imaging in the evaluation of patients with cognitive impairment or dementia is uncertain.^{25,26}

Many primary care physicians view the treatment of dementia as time-consuming and not fitting well into the usual primary care setting. Group medical visits for the treatment of dementia can lead to better compliance, improved outcomes, and provide needed support for caregivers.¹⁰

Researchers are constantly looking for pharmacologic and non-pharmacologic treatments to help with both cognitive and behavioral symptoms, such as antidepressants for agitation and psychosis in patients with dementia.²⁷ A Cochrane review recommends that selective serotonin reuptake inhibitors only be used if non-pharmacologic interventions are unsuccessful.²⁷ Still other recent Cochrane reviews suggest that exercise improves cognitive function and the ability to perform activities of daily living in patients with dementia.²⁸ Physicians must be provided up to date on effective and ineffective therapies.⁴

The AAFP and the American College of Physicians (ACP) have released guidelines for the current pharmacologic treatment of dementia.^{29,30} The key recommendations are summarized as follows:

- Clinicians should base the decision to initiate a trial of therapy with a cholinesterase inhibitor or memantine on individualized assessment. (Grade: weak recommendation, moderate-quality evidence.)
- Clinicians should base the choice of pharmacologic agents on tolerability, adverse effect profile, ease of use, and cost of medication. The evidence is insufficient to compare the effectiveness of different pharmacologic agents for the treatment of dementia. (Grade: weak recommendation, low-quality evidence.)
- There is an urgent need for further research on the clinical effectiveness of pharmacologic management of dementia.

In 2012, 15.4 million caregivers provided more than 17.5 billion hours of unpaid care, valued at \$216 billion, to family members with Alzheimer's.³¹ Family and caretaker involvement is critical in the treatment of patients with dementia and Alzheimer's, so family physicians are uniquely positioned to initiate discussions with families about the importance of treatment and monitoring



the patient's behavior and activities of daily living (ADLs). Some family members and/or caregivers may also investigate specific treatments (such as integrative therapies) and inquire about them with family physicians, in which case family physicians should be prepared to address questions and provide accurate and appropriate information.

Family physicians are also in a unique position to provide not just information to caregivers, but also to help refer them to community resources (such as support groups) that can be helpful. Some studies also suggest that family physicians are unaware of community-based social services for patients with dementia or Alzheimer's disease; therefore, family physicians should be aware of these services and be able to provide guidance to caregivers.³² Family physicians should be provided education and strategies to assist them with the integration of behavioral health specialist into practice.³³ Physicians can also make their visits with patients in nursing homes or other long-term care facilities more efficient by using a standardized clinician documentation tool.³⁴ The stress and burden of caring for someone who has dementia or Alzheimer's can cause physical and psychological problems for caregivers; family physicians should be aware of this and discuss any problems, and should be aware of existing resources and be prepared to refer patients and/or family members to such resources.

Family physicians required education and strategies to consistently implement evidence-based guidelines for the management of dementia and Alzheimer's in their patient population.

Physicians may improve their care of patients with Alzheimer's Disease by engaging in continuing medical education that provides practical integration of current evidence-based guidelines and recommendations into their standards of care, including, but not limited to the following:³⁵

- Acetylcholinesterase inhibitors are modestly effective in patients with mild to moderate Alzheimer disease, although limited by their adverse effects.
- Combination therapy with an acetylcholinesterase inhibitor and memantine (Namenda) should be considered in patients with moderate to severe Alzheimer disease.
- Atypical antipsychotic agents can improve some behavioral manifestations of Alzheimer disease but are associated with increased mortality in older patients.
- Nonsteroidal anti-inflammatory drugs, vitamin E, testosterone, estrogen, statins, and insulin sensitizers are not recommended for the treatment of Alzheimer disease.
- Physicians should consider discontinuing treatment for Alzheimer disease in patients who continue to decline despite maximal therapy.

These recommendations are provided only as assistance for physicians making clinical decisions regarding the care of their patients. As such, they cannot substitute for the individual judgment brought to each clinical situation by the patient's family physician. As with all clinical reference resources, they reflect the best understanding of the science of medicine at the time of publication, but they should be used with the clear understanding that continued research may result in new knowledge and recommendations. These recommendations are only one element in the complex process of improving the health of America. To be effective, the recommendations must be implemented. As such, physicians require continuing medical education to assist them with making decisions about specific clinical considerations.



Physicians should also be kept up to date on current research showing a possible link between proton pump inhibitors (PPIs) and risk of dementia in older adults.^{36,37}

Finally, the Alzheimer's Disease Neuroimaging Initiative (ADNI)(adcs.org) is asking for help from family physicians throughout the United States.³⁸ The largest and most comprehensive Alzheimer's disease research effort in history, ADNI is moving into its second phase -- ADNI2(www.nia.nih.gov) -- but finds itself hampered by recruitment challenges. According to principal investigator Michael Weiner, M.D., a professor of medicine, radiology, psychiatry and neurology at the University of California, San Francisco, ADNI2 is seeking 400 volunteers between the ages of 55 and 90 who have been diagnosed with early mild cognitive impairment, late mild cognitive impairment or early/mild Alzheimer's disease. The plan sets forth five goals, including the development of effective prevention and treatment approaches for Alzheimer's disease and related dementias by 2025.

Resources: Evidence-Based Practice Recommendations/Guidelines/Performance Measures

- Initial evaluation of the patient with suspected dementia²
- Evaluation of suspected dementia³
- (AMA) PCPI Approved Quality Measures: Dementia¹²
- (AMDA) Dementia in the long term care setting³⁹
- Treating Dementia With Shared Group Visits¹⁰
- Treatment of Alzheimer disease³⁵
- CDC Healthy Aging: Healthy Brain Initiative¹⁶
- (AAFP) Clinical Preventive Services Recommendations¹⁹
- Integrating a behavioral health specialist into your practice³³
- A nursing home documentation tool for more efficient visits³⁴
- Alzheimer's Association recommendations for operationalizing the detection of cognitive impairment during the Medicare Annual Wellness Visit in a primary care setting⁴⁰
- AAFP and ACP Guideline on dementia treatment³⁰
- ACR Appropriateness Criteria - dementia and movement disorders⁴¹
- FamilyDoctor.org. Alzheimer's Disease | Overview (patient resource)⁴²
- FamilyDoctor.org. Dementia | Overview (patient resource)⁴³

References

1. National Institute of Neurological Disease and Stroke (NINDS). Dementia: Hope Through Research. 2010; http://www.ninds.nih.gov/disorders/dementias/detail_dementia.htm. Accessed June, 2013.
2. Adelman AM, Daly MP. Initial evaluation of the patient with suspected dementia. *American family physician*. May 1 2005;71(9):1745-1750.
3. Simmons BB, Hartmann B, DeJoseph D. Evaluation of suspected dementia. *American family physician*. Oct 15 2011;84(8):895-902.
4. Alzheimer's Association. Alzheimer's Disease. 2013; http://www.alz.org/alzheimers_disease_facts_and_figures.asp. Accessed June, 2013.



5. Centers for Disease Control and Prevention. FastStats: Alzheimer's Disease. 2010; <http://www.cdc.gov/nchs/fastats/alzheimr.htm>. Accessed June, 2013.
6. AAFP. 2012 CME Needs Assessment: Clinical Topics. American Academy of Family Physicians; 2012.
7. American Academy of Family Physicians (AAFP). 2012 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2012.
8. American Academy of Family Physicians (AAFP). 2013 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2013.
9. American Academy of Family Physicians (AAFP). AAFP Assembly CME Outcomes Report. Leawood KS: AAFP; 2014.
10. Khandelwal C, Prentice A, Fisher J, Parrott R, Sloane PD. Treating dementia with shared group visits. *Family practice management*. May-Jun 2015;22(3):16-21.
11. Hinton L, Franz CE, Reddy G, Flores Y, Kravitz RL, Barker JC. Practice constraints, behavioral problems, and dementia care: primary care physicians' perspectives. *Journal of general internal medicine*. Nov 2007;22(11):1487-1492.
12. American Medical Association (AMA). PCPI Approved Quality Measures. 2013; <http://www.ama-assn.org/apps/listserv/x-check/qmeasure.cgi>. Accessed May, 2013.
13. Stewart TV, Loskutova N, Galliher JM, et al. Practice patterns, beliefs, and perceived barriers to care regarding dementia: a report from the American Academy of Family Physicians (AAFP) national research network. *Journal of the American Board of Family Medicine : JABFM*. Mar-Apr 2014;27(2):275-283.
14. Hochhalter AK, Bryant LL, Hunter R, et al. Multisite qualitative study of primary care physicians' and midlevel providers' self-reported practices and perceptions about maintaining cognitive health. *Preventing chronic disease*. Nov 2012;9:E169.
15. National Institute of Neurological Disease and Stroke (NINDS). NINDS Alzheimer's Disease Information Page. 2013; <http://www.ninds.nih.gov/disorders/alzheimersdisease/alzheimersdisease.htm>. Accessed June, 2013.
16. Centers for Disease Control and Prevention. Healthy Aging: Healthy Brain Initiative. 2011; <http://www.cdc.gov/aging/healthybrain/>. Accessed June, 2013.
17. Endres J, Graber MA, Dachs R. Benzodiazepines and Alzheimer disease. *American family physician*. Feb 1 2015;91(3):191-192.
18. U.S. Preventive Services Task Force (USPSTF). USPSTF A-Z Topic Guide. 2012; <http://www.uspreventiveservicestaskforce.org/uspsttopics.htm>. Accessed June, 2013.
19. American Academy of Family Physicians (AAFP). Clinical Preventive Services Recommendations. 2012; <http://www.aafp.org/online/en/home/clinical/exam.html>. Accessed May, 2013.
20. Fan T, Rossi C. Putting Prevention into Practice: AN EVIDENCE-BASED APPROACH - Screening for Cognitive Impairment in Older Adults. *American family physician*. 2015;92(2):125-126. <http://www.aafp.org/afp/2015/0715/p125.html>. Accessed Aug 2015.
21. Moyer VA, Force USPST. Screening for cognitive impairment in older adults: U.S. Preventive Services Task Force recommendation statement. *Annals of internal medicine*. Jun 3 2014;160(11):791-797.
22. Davis DH, Creavin ST, Yip JL, Noel-Storr AH, Brayne C, Cullum S. Montreal Cognitive Assessment for the diagnosis of Alzheimer's disease and other dementias. *Cochrane Database Syst Rev*. 2015;10:CD010775.



23. Nasreddine ZS, Phillips NA, Bedirian V, et al. The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc.* Apr 2005;53(4):695-699.
24. American Academy of Family Physicians (AAFP). Dementia - Screening and Diagnosis of Early Cognitive Impairment. *NRN Studies* 2008; <http://www.aafp.org/patient-care/nrn/studies/all/dementia-screening.html>. Accessed Apr, 2016.
25. Jansen WJ, Ossenkoppele R, Knol DL, et al. Prevalence of cerebral amyloid pathology in persons without dementia: a meta-analysis. *JAMA : the journal of the American Medical Association.* May 19 2015;313(19):1924-1938.
26. Ossenkoppele R, Jansen WJ, Rabinovici GD, et al. Prevalence of amyloid PET positivity in dementia syndromes: a meta-analysis. *JAMA : the journal of the American Medical Association.* May 19 2015;313(19):1939-1949.
27. Bui Q. Antidepressants for agitation and psychosis in patients with dementia. *American family physician.* Jan 1 2012;85(1):20-22.
28. Raleigh MF, Shaha D. Exercise programs for older patients with dementia. *American family physician.* Sep 1 2014;90(5):287.
29. Qaseem A, Snow V, Cross JT, Jr., et al. Current pharmacologic treatment of dementia: a clinical practice guideline from the American College of Physicians and the American Academy of Family Physicians. *Annals of internal medicine.* Mar 4 2008;148(5):370-378.
30. Graham L. AAFP and ACP release guideline on dementia treatment. *American family physician.* Apr 15 2008;77(8):1173-1175.
31. Thies W, Bleiler L. 2011 Alzheimer's disease facts and figures. *Alzheimer's & dementia : the journal of the Alzheimer's Association.* Mar 2011;7(2):208-244.
32. Yaffe MJ, Orzeck P, Barylak L. Family physicians' perspectives on care of dementia patients and family caregivers. *Canadian family physician Medecin de famille canadien.* Jul 2008;54(7):1008-1015.
33. Reitz R, Fifield P, Whistler P. Integrating a behavioral health specialist into your practice. *Family practice management.* Jan-Feb 2011;18(1):18-21.
34. Kalish VB, Burns OB, Unwin BK. A nursing home documentation tool for more efficient visits. *Family practice management.* Mar-Apr 2012;19(2):19-21.
35. Winslow BT, Onysko MK, Stob CM, Hazlewood KA. Treatment of Alzheimer disease. *American family physician.* Jun 15 2011;83(12):1403-1412.
36. Gomm W, von Holt K, Thome F, et al. Association of Proton Pump Inhibitors With Risk of Dementia: A Pharmacoepidemiological Claims Data Analysis. *JAMA neurology.* Apr 1 2016;73(4):410-416.
37. Kuller LH. Do Proton Pump Inhibitors Increase the Risk of Dementia? *JAMA neurology.* 2016.
38. Brown M. Largest Alzheimer's Disease Study Seeking Recruitment Help from FPs. *AAFP News.* 2012 <http://www.aafp.org/news/health-of-the-public/20120613alzheimersstudy.html>. Accessed Aug 2015.
39. National Guideline Clearinghouse. Dementia in the long term care setting. 2012; <http://www.guideline.gov/content.aspx?id=45525&search=dementia>. Accessed 7/31/2014.
40. Cordell CB, Borson S, Boustani M, et al. Alzheimer's Association recommendations for operationalizing the detection of cognitive impairment during the Medicare Annual



- Wellness Visit in a primary care setting. *Alzheimer's & dementia : the journal of the Alzheimer's Association*. Mar 2013;9(2):141-150.
41. Wipplod FJ II, Cornelius RS, Broderick DF, et al. ACR Appropriateness Criteria - dementia and movement disorders. 2010;
<http://www.guideline.gov/content.aspx?id=32612&search=dementia>. Accessed 6/6/2013.
 42. FamilyDoctor.org. Alzheimer's Disease | Overview. 2012;
<http://familydoctor.org/familydoctor/en/diseases-conditions/alzheimers-disease.html>. Accessed August, 2013.
 43. FamilyDoctor.org. Dementia | Overview. 2000;
<http://familydoctor.org/familydoctor/en/diseases-conditions/dementia.html>. Accessed August, 2013.