



Body System: Nephrologic			
Session Topic: Fluid and Electrolyte Abnormalities			
Educational Format		Faculty Expertise Required	
REQUIRED	Interactive Lecture	Expertise in the field of study. Experience teaching in the field of study is desired. Preferred experience with audience response systems (ARS). Utilizing polling questions and engaging the learners in Q&A during the final 15 minutes of the session are required.	
OPTIONAL	Problem-Based Learning (PBL)	Expertise teaching highly interactive, small group learning environments. Case-based, with experience developing and teaching case scenarios for simulation labs preferred. Other workshop-oriented designs may be accommodated. A typical PBL room is set for 50-100 participants, with 7-8 each per round table. <u>Please describe your interest and plan for teaching a PBL on your proposal form.</u>	
Professional Practice Gap		Learning Objective(s) that will close the gap and meet the need	Outcome Being Measured
<ul style="list-style-type: none"> • Knowledge gaps about the association between hypercalcemia and hyperparathyroidism and malignancy and be able to select the appropriate treatment based on etiology. • Knowledge gaps in being able to identify and treat the underlying etiology of hyponatremia and prescribe appropriate treatment for hyponatremia based on symptoms, degree, and presence of any hypotension. • Knowledge gaps differentiating the etiology of hypernatremia and prescribe appropriate treatment with caution due to the dangers of rapid correction. • Knowledge gaps treating vitamin D deficiency based on serum 25-hydroxyvitamin D levels. • Knowledge gaps in diagnosing hyper- and 		<ol style="list-style-type: none"> 1. Identify the underlying etiology of hypercalcemia and select the appropriate treatment. 2. Identify and treat the underlying etiology of hyponatremia and prescribe appropriate treatment for hyponatremia based on symptoms, degree, and presence of any hypotension. 3. Differentiate the etiology of hypernatremia and prescribe appropriate treatment with caution due to the dangers of rapid correction. 4. Identify the underlying etiology of hyper- and hypokalemia and select appropriate treatment. 	Learners will submit written commitment to change statements on the session evaluation, indicating how they plan to implement presented practice recommendations.



hypokalemia — and the underlying causes — and institute appropriate management.			
<ul style="list-style-type: none"> • Knowledge and performance gaps in evaluating patients at highest risk of drug-induced nephrotoxicity, and adjusting medications as indicated 			
ACGME Core Competencies Addressed (select all that apply)			
X	Medical Knowledge		Patient Care
	Interpersonal and Communication Skills		Practice-Based Learning and Improvement
	Professionalism	X	Systems-Based Practice
Faculty Instructional Goals			
<p>Faculty play a vital role in assisting the AAFP to achieve its mission by providing high-quality, innovative education for physicians, residents and medical students that will encompass the art, science, evidence and socio-economics of family medicine and to support the pursuit of lifelong learning. By achieving the instructional goals provided, faculty will facilitate the application of new knowledge and skills gained by learners to practice, so that they may optimize care provided to their patients.</p> <ul style="list-style-type: none"> • Provide up to 3 evidence-based recommended practice changes that can be immediately implemented, at the conclusion of the session; including SORT taxonomy & reference citations • Facilitate learner engagement during the session • Address related practice barriers to foster optimal patient management • Provide recommended journal resources and tools, during the session, from the American Family Physician (AFP), Family Practice Management (FPM), and Familydoctor.org patient resources; those listed in the <u>References</u> section below are a good place to start <ul style="list-style-type: none"> ○ Visit http://www.aafp.org/journals for additional resources ○ Visit http://familydoctor.org for patient education and resources • Provide recommendations for identifying the underlying etiology of hypercalcemia and select the appropriate treatment. • Provide recommendations for identifying and treating the underlying etiology of hyponatremia and prescribe appropriate treatment for hyponatremia based on symptoms, degree, and presence of any hypotension. • Provide recommendations for differentiating the etiology of hypernatremia and prescribe appropriate treatment with caution due the dangers of rapid correction. • Provide recommendations for identifying the underlying etiology of hyper- and hypokalemia and select appropriate treatment. 			

Needs Assessment

*Note: for purposes of this education, the focus will be on: *hypercalcemia, vitamin D deficiency and hypocalcemia, hyper/hyp-natremia, and hyper/hypo-kalemia.*



Fluid and electrolyte abnormalities, such as hyper/hypo-calcemia, hyper/hypo-natremia, and hyper/hypo-kalemia are commonly encountered by family physicians in inpatient and outpatient settings.^{1,2} Electrolyte disorders are especially common among older, community dwelling patients, and is especially associated with diabetes mellitus and diuretics.³ Approximately one in 500 patients who are treated in a general medicine clinic have undiagnosed primary hyperparathyroidism, the leading cause of hypercalcemia. The principal challenges in the management of hypercalcemia are distinguishing primary hyperparathyroidism from conditions that will not respond to parathyroidectomy and knowing when it is appropriate to refer the patient for surgery.⁴ Primary hyperparathyroidism, the most common cause of hypercalcemia in outpatients, is often discovered incidentally during evaluation of serum electrolyte levels. Most patients are asymptomatic, and it is therefore important to be aware that subtle and nonspecific symptoms may be present.⁵ Elderly hospitalized patients with ionized hypocalcemia and hypophosphatemia, with or without an elevated parathyroid hormone level, are most likely deficient in vitamin D.² The prevalence of severe hyponatremia (serum sodium level less than 125 mEq per L) was 4.5%, 0.8%, and 10.3%, respectively. It is estimated that hyponatremia occurs in 4% to 7% of the ambulatory population, with rates of 18.8% in nursing homes.¹ The prevalence of hyperkalemia in hospitalized patients is between 1 and 10 percent.⁶

Physicians should be knowledgeable about electrolyte abnormalities and be able to make a diagnosis through physical exam, history taking, or laboratory tests when applicable. However, data from the 2012 American Academy of Family Physicians (AAFP) CME Needs Assessment Survey indicates that family physicians have gaps in the medical skill necessary to provide optimal patient management of electrolyte abnormalities, dehydration, and oral rehydration.⁷ More specifically, CME outcomes data from 2013 AAFP Assembly: *Fluid and Electrolyte Abnormalities* sessions suggest that physicians have knowledge gaps with regard to understanding different presentations of fluid imbalances; evaluation after diagnosis; ordering appropriate lab work; knowing when to check for imbalances; being aware of calculation tools; and when to refer.⁸

In order to address identified knowledge gaps related to the diagnosis and management of fluid and electrolyte abnormalities, family physicians should engage in continuing education that facilitates the use of evidence-based recommendations and guidelines. Physicians are encouraged to consider the following evidence-based recommendations:^{1,2,6,9,10}

- Patients with hyperkalemia who have electrocardiographic (ECG) changes, a rapid rate of rise of serum potassium, decreased renal function, or significant acidosis should be urgently treated.
- Patients with hyperkalemia and characteristic ECG changes should be given intravenous calcium gluconate.
- Acutely lower potassium by giving intravenous insulin with glucose, a beta2 agonist by nebulizer, or both.
- Total body potassium should usually be lowered with sodium polystyrene sulfonate (Kayexalate).
- In patients with severe symptomatic hyponatremia, the rate of sodium correction should be 6 to 12 mEq per L in the first 24 hours and 18 mEq per L or less in 48 hours.
- Consensus guidelines based on systematic reviews



- A bolus of 100 to 150 mL of hypertonic 3% saline can be given to correct severe hyponatremia.
- Consensus guidelines based on small studies
- Vaptans appear to be safe for the treatment of severe hypervolemic and euvolemic hyponatremia but should not be used routinely.
- Consensus guidelines based on observational studies
- Chronic hypernatremia should be corrected at a rate of 0.5 mEq per L per hour, with a maximum change of 8 to 10 mEq per L in a 24-hour period.
- In patients with severe vitamin D deficiency (serum levels of below 8 ng per mL with hypocalcemia), 50,000 IU of vitamin D should be given daily for one to three weeks, followed by weekly doses of 50,000 IU.
- In critically ill patients, albumin-adjusted calcium levels underestimate true or ionized hypocalcemia. Therefore, measured ionized calcium levels are recommended, particularly in patients who are being treated in an intensive care unit.
- If calcium supplementation alone fails to maintain normal serum levels, the patient is vitamin D deficient or resistant and may benefit from a trial of calcitriol (Rocatriol).
- If the vitamin D deficiency is severe, the patient will require 90 mmol per L in the first 24 hours: 6 mL of K₂PO₄ added to each liter of fluid and given at 200 mL per hour (1 mL of K₂PO₄ is equal to 4 mEq of potassium and 3.0 mmol per L or 93 mg of phosphate).
- Adequate fluid balance should be maintained in patients with acute kidney injury by using isotonic solutions (e.g., normal saline) instead of hyperoncotic solutions (e.g., dextrans, hydroxyethyl starch, albumin).
-

Physicians are encouraged to establish shared decision-making relationships with adult patients with electrolyte abnormalities, and to develop a patient-physician relationship that promotes family-centered shared decision-making. Optimal management requires close collaboration among primary care physicians, nephrologists, hospitalists, and other subspecialists participating in the care of the patient. Physicians can improve patient satisfaction with the referral process by using readily available strategies and tools such as, improving internal office communication, engaging patients in scheduling, facilitating the appointment, tracking referral results, analyzing data for improvement opportunities, and gathering patient feedback.^{11,12}

Resources: Evidence-Based Practice Recommendations/Guidelines/Performance Measures

- Diagnosis and management of sodium disorders: hyponatremia and hypernatremia¹
- Hyperkalemia⁶
- A Practical Approach to Hypercalcemia⁴
- Undiagnosed vitamin D deficiency in the hospitalized patient²
- Drug-induced nephrotoxicity¹³
- Parathyroid disorders⁵
- Management of hyponatremia⁹
- Diagnosis, evaluation, and treatment of hyponatremia: expert panel recommendations¹⁴
- Acute kidney injury: a guide to diagnosis and management¹⁰



- Engaging Patients in Collaborative Care Plans¹⁵
- Simple tools to increase patient satisfaction with the referral process¹¹
- Adding health education specialists to your practice¹⁶
- Envisioning new roles for medical assistants: strategies from patient-centered medical homes¹⁷
- The benefits of using care coordinators in primary care: a case study¹⁸
- The Use of Symptom Diaries in Outpatient Care¹⁹
- Health Coaching: Teaching Patients to Fish²⁰
- Medication adherence: we didn't ask and they didn't tell²¹
- Encouraging patients to change unhealthy behaviors with motivational interviewing²²
- FamilyDoctor.org. Hydration: Why It's So Important (patient education)²³
- FamilyDoctor.org. Artificial Hydration and Nutrition (patient education)²⁴
- FamilyDoctor.org. Athletes: The Importance of Good Hydration (patient education)²⁵
- FamilyDoctor.org. Caregiving: Caring for an Elderly Relative - Managing Medicines (patient education)²⁶
- FamilyDoctor.org. Urinary Incontinence | Overview (patient education)²⁷
- FamilyDoctor.org. Chronic Kidney Disease | Overview (patient education)²⁸

References

1. Braun MM, Barstow CH, Pyzocha NJ. Diagnosis and management of sodium disorders: hyponatremia and hypernatremia. *American family physician*. Mar 1 2015;91(5):299-307.
2. Lyman D. Undiagnosed vitamin D deficiency in the hospitalized patient. *American family physician*. Jan 15 2005;71(2):299-304.
3. Liamis G, Rodenburg EM, Hofman A, Zietse R, Stricker BH, Hoorn EJ. Electrolyte disorders in community subjects: prevalence and risk factors. *The American journal of medicine*. Mar 2013;126(3):256-263.
4. Carroll MF, Schade DS. A practical approach to hypercalcemia. *American family physician*. May 1 2003;67(9):1959-1966.
5. Michels TC, Kelly KM. Parathyroid disorders. *American family physician*. Aug 15 2013;88(4):249-257.
6. Hollander-Rodriguez JC, Calvert JF, Jr. Hyperkalemia. *American family physician*. Jan 15 2006;73(2):283-290.
7. AAFP. 2012 CME Needs Assessment: Clinical Topics. American Academy of Family Physicians; 2012.
8. American Academy of Family Physicians (AAFP). 2013 AAFP Scientific Assembly: CME Outcomes Report. Leawood KS: AAFP; 2013.
9. Goh KP. Management of hyponatremia. *American family physician*. May 15 2004;69(10):2387-2394.
10. Rahman M, Shad F, Smith MC. Acute kidney injury: a guide to diagnosis and management. *American family physician*. Oct 1 2012;86(7):631-639.
11. Jarve RK, Dool DW. Simple tools to increase patient satisfaction with the referral process. *Family practice management*. Nov-Dec 2011;18(6):9-14.



12. American Academy of Family Physicians (AAFP). FPM Toolbox: Referral Management. 2013; <http://www.aafp.org/fpm/toolBox/viewToolType.htm?toolTypeId=26>. Accessed July, 2014.
13. Naughton CA. Drug-induced nephrotoxicity. *American family physician*. Sep 15 2008;78(6):743-750.
14. Verbalis JG, Goldsmith SR, Greenberg A, et al. Diagnosis, evaluation, and treatment of hyponatremia: expert panel recommendations. *The American journal of medicine*. Oct 2013;126(10 Suppl 1):S1-42.
15. Mauksch L, Safford B. Engaging Patients in Collaborative Care Plans. *Family practice management*. 2013;20(3):35-39.
16. Chambliss ML, Lineberry S, Evans WM, Bibeau DL. Adding health education specialists to your practice. *Family practice management*. Mar-Apr 2014;21(2):10-15.
17. Naughton D, Adelman AM, Bricker P, Miller-Day M, Gabbay R. Envisioning new roles for medical assistants: strategies from patient-centered medical homes. *Family practice management*. Mar-Apr 2013;20(2):7-12.
18. Mullins A, Mooney J, Fowler R. The benefits of using care coordinators in primary care: a case study. *Family practice management*. Nov-Dec 2013;20(6):18-21.
19. Hodge B. The Use of Symptom Diaries in Outpatient Care. *Family practice management*. 2013;20(3):24-28.
20. Ghorob A. Health Coaching: Teaching Patients to Fish. *Family practice management*. 2013;20(3):40-42.
21. Brown M, Sinsky CA. Medication adherence: we didn't ask and they didn't tell. *Family practice management*. Mar-Apr 2013;20(2):25-30.
22. Stewart EE, Fox CH. Encouraging patients to change unhealthy behaviors with motivational interviewing. *Family practice management*. May-Jun 2011;18(3):21-25.
23. FamilyDoctor.org. Hydration: Why It's So Important. 2010; <http://familydoctor.org/familydoctor/en/prevention-wellness/food-nutrition/nutrients/hydration-why-its-so-important.html>. Accessed August, 2013.
24. FamilyDoctor.org. Artificial Hydration and Nutrition 2000; <http://familydoctor.org/familydoctor/en/healthcare-management/end-of-life-issues/artificial-hydration-and-nutrition.html>. Accessed September, 2014.
25. FamilyDoctor.org. Athletes: The Importance of Good Hydration 2010; <http://familydoctor.org/familydoctor/en/prevention-wellness/exercise-fitness/exercise-basics/athletes-the-importance-of-good-hydration.html>. Accessed September, 2014.
26. FamilyDoctor.org. Caregiving: Caring for an Elderly Relative - Managing Medicines. 2012; <http://familydoctor.org/familydoctor/en/seniors/caring-for-an-elderly-relative/managing-medicines.html>. Accessed August, 2013.
27. FamilyDoctor.org. Urinary Incontinence | Overview. 2000; <http://familydoctor.org/familydoctor/en/diseases-conditions/urinary-incontinence.html>. Accessed August, 2013.
28. FamilyDoctor.org. Chronic Kidney Disease | Overview. 2005; <http://familydoctor.org/familydoctor/en/diseases-conditions/chronic-kidney-disease.html>. Accessed August, 2013.