

# PHARMACOLOGIC PRODUCT GUIDE: FDA-APPROVED MEDICATIONS

NICOTINE REPLACEMENT THERAPY (NRT) FORMULATIONS					BUPROPION SR	VARENICLINE	
	GUM	LOZENGE	TRANSDERMAL PATCH	NASAL SPRAY	ORAL INHALER		
PRODUCT	<b>Nicorette<sup>1</sup>, Generic</b> OTC 2 mg, 4 mg original, fruit, mint, orange, cinnamon	<b>Commit<sup>1</sup>, Generic</b> OTC 2 mg, 4 mg mint, cherry, cappuccino, original (light mint)	<b>Nicoderm CQ<sup>1</sup>, Generic<sup>2</sup></b> OTC, Rx 7 mg, 14 mg, 21 mg (24-hour release)	<b>Nicotrol NS<sup>3</sup></b> Rx Metered spray 0.5 mg nicotine in 50 mL aqueous nicotine solution	<b>Nicotrol Inhaler<sup>3</sup></b> Rx 10 mg cartridge delivers 4 mg inhaled nicotine vapor	<b>Zyban<sup>1</sup>, Generic</b> Rx 150 mg sustained-release tablet	<b>Chantix<sup>3</sup></b> Rx 0.5 mg, 1 mg tablet
PRECAUTIONS	<ul style="list-style-type: none"> <li>Recent (<math>\leq 2</math> weeks) myocardial infarction</li> <li>Serious underlying arrhythmias</li> <li>Serious or worsening angina pectoris</li> <li>Temporomandibular joint disease</li> <li>Pregnancy<sup>4</sup> and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul>	<ul style="list-style-type: none"> <li>Recent (<math>\leq 2</math> weeks) myocardial infarction</li> <li>Serious underlying arrhythmias</li> <li>Serious or worsening angina pectoris</li> <li>Pregnancy<sup>4</sup> and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul>	<ul style="list-style-type: none"> <li>Recent (<math>\leq 2</math> weeks) myocardial infarction</li> <li>Serious underlying arrhythmias</li> <li>Serious or worsening angina pectoris</li> <li>Pregnancy<sup>4</sup> (Rx formulations, category D) and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul>	<ul style="list-style-type: none"> <li>Recent (<math>\leq 2</math> weeks) myocardial infarction</li> <li>Serious underlying arrhythmias</li> <li>Serious or worsening angina pectoris</li> <li>Underlying chronic nasal disorders (rhinitis, nasal polyps, sinusitis)</li> <li>Severe reactive airway disease</li> <li>Pregnancy<sup>4</sup> (category D) and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul>	<ul style="list-style-type: none"> <li>Recent (<math>\leq 2</math> weeks) myocardial infarction</li> <li>Serious underlying arrhythmias</li> <li>Serious or worsening angina pectoris</li> <li>Bronchospastic disease</li> <li>Pregnancy<sup>4</sup> (category D) and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul>	<ul style="list-style-type: none"> <li>Concomitant therapy with medications or medical conditions known to lower the seizure threshold</li> <li>Severe hepatic cirrhosis</li> <li>Pregnancy<sup>4</sup> (category C) and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul> <p><b>Contraindications:</b></p> <ul style="list-style-type: none"> <li>Seizure disorder</li> <li>Concomitant bupropion (e.g., Wellbutrin) therapy</li> <li>Current or prior diagnosis of bulimia or anorexia nervosa</li> <li>Simultaneous abrupt discontinuation of alcohol or sedatives (including benzodiazepines)</li> <li>MAO inhibitor therapy in previous 14 days</li> </ul>	<ul style="list-style-type: none"> <li>Severe renal impairment (dosage adjustment is necessary)</li> <li>Pregnancy<sup>4</sup> (category C) and breastfeeding</li> <li>Adolescents (&lt;18 years)</li> </ul> <p><b>Warnings:</b></p> <ul style="list-style-type: none"> <li>Neuropsychiatric symptoms (behavior changes, agitation, depressed mood, suicidal ideation or behavior)</li> <li>Safety and efficacy have not been established in patients with serious psychiatric illness</li> </ul>
DOSING	<p><math>\geq 25</math> cigarettes/day: 4 mg  <math>&lt; 25</math> cigarettes/day: 2 mg</p> <p>Week 1 – 6:            1 piece q 1 – 2 hours            Week 7 – 9:            1 piece q 2 – 4 hours            Week 10 – 12:            1 piece q 4 – 8 hours</p> <ul style="list-style-type: none"> <li>Maximum, 24 pieces/day</li> <li>Chew each piece slowly</li> <li>Park between cheek and gum when peppery or tingling sensation appears (~15 – 30 chews)</li> <li>Resume chewing when taste or tingle fades</li> <li>Repeat chew/park steps until most of the nicotine is gone (taste or tingle does not return; generally 30 min)</li> <li>Park in different areas of mouth</li> <li>No food or beverages 15 min before or during use</li> <li>Duration: up to 12 weeks</li> </ul>	<p>1st cigarette <math>\leq 30</math> minutes after waking: 4 mg            1st cigarette <math>&gt; 30</math> minutes after waking: 2 mg</p> <p>Week 1 – 6:            1 lozenge q 1 – 2 hours            Week 7 – 9:            1 lozenge q 2 – 4 hours            Week 10 – 12:            1 lozenge q 4 – 8 hours</p> <ul style="list-style-type: none"> <li>Maximum: 20 lozenges/day</li> <li>Allow to dissolve slowly (20 – 30 minutes)</li> <li>Nicotine release may cause a warm, tingling sensation</li> <li>Do not chew or swallow</li> <li>Occasionally rotate to different areas of the mouth</li> <li>No food or beverages 15 minutes before or during use</li> <li>Duration: up to 12 weeks</li> </ul>	<p><math>&gt; 10</math> cigarettes/day:            21 mg/day x 4 weeks (generic)            6 weeks (NicoDerm CQ)            14 mg/day x 2 weeks            7 mg/day x 2 weeks</p> <p><math>\leq 10</math> cigarettes/day:            14 mg/day x 6 weeks            7 mg/day x 2 weeks</p> <ul style="list-style-type: none"> <li>May wear patch for 16 hours if patient experiences sleep disturbances (remove at bedtime)</li> <li>Duration: 8 – 10 weeks</li> </ul>	<p>1 – 2 doses/hour            (8 – 40 doses/day)            One dose = 2 sprays (one in each nostril); each spray delivers 0.5 mg of nicotine to the nasal mucosa</p> <ul style="list-style-type: none"> <li>Maximum:              - 5 doses/hour              - 40 doses/day</li> <li>For best results, initially use at least 8 doses/day</li> <li>Patients should not sniff, swallow, or inhale through the nose as the spray is being administered</li> <li>Duration: 3 – 6 months</li> </ul>	<p>6 – 16 cartridges/day            Individualized dosing; initially use 1 cartridge q 1 – 2 hours</p> <ul style="list-style-type: none"> <li>Best effects with continuous puffing for 20 minutes</li> <li>Nicotine in cartridge is depleted after 20 minutes of active puffing</li> <li>Patient should inhale into back of throat or puff in short breaths</li> <li>Do NOT inhale into the lungs (like a cigarette) but “puff” as if lighting a pipe</li> <li>Open cartridge retains potency for 24 hours</li> <li>Duration: 3 – 6 months</li> </ul>	<p>150 mg po q AM x 3 days, then increase to 150 mg po bid</p> <ul style="list-style-type: none"> <li>Do not exceed 300 mg/day</li> <li>Patients should begin therapy 1-2 weeks prior to quit date</li> <li>Allow at least 8 hours between doses</li> <li>Avoid bedtime dosing to minimize insomnia</li> <li>Dose tapering is not necessary</li> <li>Can be used safely with NRT</li> <li>Duration: 7 – 12 weeks, with maintenance up to 6 months in selected patients</li> </ul>	<p>Days 1 – 3:            0.5 mg po q AM            Days 4 – 7:            0.5 mg po bid            Weeks 2 – 12:            1 mg po bid</p> <ul style="list-style-type: none"> <li>Patients should begin therapy 1 week prior to quit date</li> <li>Take dose after eating with a full glass of water</li> <li>Dose tapering is not necessary</li> <li>Nausea and insomnia are side effects that are usually temporary</li> <li>Duration: 12 weeks; an additional 12 week course may be used in selected patients</li> </ul>

NICOTINE REPLACEMENT THERAPY (NRT) FORMULATIONS						BUPROPION SR	VARENICLINE
	GUM	LOZENGE	TRANSDERMAL PATCH	NASAL SPRAY	ORAL INHALER		
ADVERSE EFFECTS	<ul style="list-style-type: none"> <li>■ Mouth/jaw soreness</li> <li>■ Hiccups</li> <li>■ Dyspepsia</li> <li>■ Hypersalivation</li> <li>■ Effects associated with incorrect chewing technique:               <ul style="list-style-type: none"> <li>- Lightheadedness</li> <li>- Nausea/vomiting</li> <li>- Throat and mouth irritation</li> </ul> </li> </ul>	<ul style="list-style-type: none"> <li>■ Nausea</li> <li>■ Hiccups</li> <li>■ Cough</li> <li>■ Heartburn</li> <li>■ Headache</li> <li>■ Flatulence</li> <li>■ Insomnia</li> </ul>	<ul style="list-style-type: none"> <li>■ Local skin reactions (erythema, pruritus, burning)</li> <li>■ Headache</li> <li>■ Sleep disturbances (insomnia or abnormal/vivid dreams) associated with nocturnal nicotine absorption</li> </ul>	<ul style="list-style-type: none"> <li>■ Nasal and/or throat irritation (hot, peppery, or burning sensation)</li> <li>■ Rhinitis</li> <li>■ Tearing</li> <li>■ Sneezing</li> <li>■ Cough</li> <li>■ Headache</li> </ul>	<ul style="list-style-type: none"> <li>■ Mouth and/or throat irritation</li> <li>■ Cough</li> <li>■ Rhinitis</li> <li>■ Dyspepsia</li> <li>■ Headache</li> <li>■ Hiccups</li> </ul>	<ul style="list-style-type: none"> <li>■ Insomnia</li> <li>■ Dry mouth</li> <li>■ Nervousness/difficulty concentrating</li> <li>■ Rash</li> <li>■ Constipation</li> <li>■ Seizures (risk is 1/1,000 [0.1%])</li> </ul>	<ul style="list-style-type: none"> <li>■ Nausea</li> <li>■ Sleep disturbances (insomnia, abnormal/vivid dreams)</li> <li>■ Constipation</li> <li>■ Flatulence</li> <li>■ Vomiting</li> <li>■ Neuropsychiatric symptoms (see PRECAUTIONS)</li> </ul>
ADVANTAGES	<ul style="list-style-type: none"> <li>■ Might satisfy oral cravings</li> <li>■ Might delay weight gain</li> <li>■ Patients can titrate therapy to manage withdrawal symptoms</li> </ul>	<ul style="list-style-type: none"> <li>■ Might satisfy oral cravings</li> <li>■ Might delay weight gain</li> <li>■ Patients can titrate therapy to manage withdrawal symptom</li> </ul>	<ul style="list-style-type: none"> <li>■ Provides consistent nicotine levels over 24 hours</li> <li>■ Once-a-day dosing associated with fewer compliance problems</li> </ul>	<ul style="list-style-type: none"> <li>■ Patients can titrate therapy to manage withdrawal symptoms</li> </ul>	<ul style="list-style-type: none"> <li>■ Patients can titrate therapy to manage withdrawal symptoms</li> <li>■ Mimics hand-to-mouth ritual of smoking</li> </ul>	<ul style="list-style-type: none"> <li>■ Easy to use; oral formulation might be associated with fewer compliance problems</li> <li>■ Can be used with NRT</li> <li>■ Might be beneficial in patients with depression</li> </ul>	<ul style="list-style-type: none"> <li>■ Easy to use; oral formulation might be associated with fewer compliance problems</li> <li>■ Offers a new mechanism of action for patients who have failed other agents</li> </ul>
DISADVANTAGES	<ul style="list-style-type: none"> <li>■ Gum chewing may not be socially acceptable</li> <li>■ Might be problematic for patients with significant dental work</li> <li>■ Patients must use proper chewing technique to minimize adverse effects</li> <li>■ Need for frequent dosing can compromise compliance</li> </ul>	<ul style="list-style-type: none"> <li>■ Gastrointestinal side effects (nausea, hiccups, heartburn) might be bothersome</li> <li>■ Need for frequent dosing can compromise compliance</li> </ul>	<ul style="list-style-type: none"> <li>■ Patients cannot titrate the dose</li> <li>■ Allergic reactions to adhesive might occur</li> <li>■ Patients with dermatologic conditions should not use the patch</li> </ul>	<ul style="list-style-type: none"> <li>■ Nasal/throat irritation may be bothersome</li> <li>■ Patients must wait 5 minutes before driving or operating heavy machinery</li> <li>■ Patients with chronic nasal disorders or severe reactive airway disease should not use the spray</li> <li>■ Need for frequent dosing can compromise compliance</li> </ul>	<ul style="list-style-type: none"> <li>■ Initial throat or mouth irritation can be bothersome</li> <li>■ Cartridges should not be stored in very warm conditions or used in very cold conditions</li> <li>■ Patients with underlying bronchospastic disease must use the inhaler with caution</li> <li>■ Need for frequent dosing can compromise compliance</li> </ul>	<ul style="list-style-type: none"> <li>■ Seizure risk is increased</li> <li>■ Several contraindications and precautions preclude use in some patients (see PRECAUTIONS)</li> </ul>	<ul style="list-style-type: none"> <li>■ May induce nausea in up to one third of patients</li> <li>■ Post-marketing surveillance data indicate potential for neuropsychiatric symptoms (see PRECAUTIONS)</li> </ul>
COST/DAY <sup>5</sup>	2 mg: \$3.28 – \$6.58 (9 pieces) 4 mg: \$4.31 – \$6.58 (9 pieces)	2 mg: \$3.66 – \$5.26 (9 pieces) 4 mg: \$3.66 – \$5.26 (9 pieces)	\$1.90 – \$3.89 (1 patch)	\$4.10 (8 doses)	\$7.32 (6 cartridges)	\$3.62 – \$7.78 (2 tablets)	\$4.90 – \$5.18 (2 tablets)

05/09

<sup>1</sup> Marketed by GlaxoSmithKline.

<sup>2</sup> Transdermal patch formulation previously marketed as Habitrol.

<sup>3</sup> Marketed by Pfizer.

<sup>4</sup> The U.S. Clinical Practice Guideline states that pregnant smokers should be encouraged to quit without medication based on insufficient evidence of effectiveness and hypothetical concerns with safety. Pregnant smokers should be offered cessation counseling interventions that exceed minimal advice to quit.

<sup>5</sup> Average wholesale price from Medi-Span Electronic Drug File. Indianapolis, IN: Wolters Kluwer Health, April 2009.

Abbreviations: Hx = history; MAO = monoamine oxidase; NRT = nicotine replacement therapy; OTC = (over-the-counter) non-prescription product; Rx = prescription product.

**For complete prescribing information, please refer to the manufacturers' package inserts.**

Rx for Change: Clinician-Assisted Tobacco Cessation. Copyright © 1999-2009. The Regents of the University of California. All rights reserved. Reprinted with permission.

**ASK AND ACT**  
 A TOBACCO CESSATION PROGRAM

*Ask and Act is supported in part by the Smoking Cessation Leadership Center.*