

# Integrating Tobacco Cessation Into Electronic Health Records

The U.S. Public Health Service Clinical Practice Guideline, *Treating Tobacco Use and Dependence*, calls for systems-level tobacco intervention efforts. Electronic health records (EHRs) allow for integration of these guidelines into the practice workflow, facilitating system-level changes to reduce tobacco use.


The Certification Commission for Healthcare Information Technology (CCHIT) recognizes the importance of embedding tobacco cessation intervention documentation in EHRs. The CCHIT Ambulatory EHR Certification Criteria on the roadmap for 2009 include:

- AM 06.07: The system shall be capable of documenting current and past tobacco use in a quantitative fashion. Discussion note: Any quantitative measure of amount of consumption, time of start, time of stop, or total duration would be acceptable.
- AM 06.08: The system shall be capable of documenting that tobacco cessation counseling was provided, including a date stamp.

The American Academy of Family Physicians (AAFP) and the American Academy of Pediatrics (AAP) have developed a joint statement advocating that EHRs include a template that prompts clinicians and/or their practice teams to collect information about tobacco use, secondhand smoke exposure, cessation interest and past quit attempts. The electronic health record should also include automatic prompts that remind clinicians to encourage quitting, advise about smoke-free environments, and connect patients and families to appropriate cessation resources and materials.

This template should be automated to appear when patients present with complaints such as cough, upper respiratory problems, diabetes, ear infections, hypertension, depression, anxiety and asthma, as well as for well-patient exams.

*Template recommendations are on the back of this document.*



AMERICAN ACADEMY OF  
FAMILY PHYSICIANS  
STRONG MEDICINE FOR AMERICA

**ASK AND ACT**  
A TOBACCO CESSATION PROGRAM

## What should be included in a tobacco cessation EHR template?

Including tobacco use status as a vital sign provides an opportunity for office staff to begin the process. Status can be documented as current, former or never. A complementary field should document secondhand smoke exposure: current, former or never.

*The template should include:*

### History:

Type of tobacco:

- Cigarettes    Pipe    Cigars    Smokeless

How many years? \_\_\_\_\_ Packs per day: \_\_\_\_\_

Brand: \_\_\_\_\_

Approx date of last quit attempt: \_\_\_\_\_

Medication used in previous quit attempt:

- Patch  
 Inhaler  
 Gum  
 Lozenge  
 Bupropion  
 Varenicline  
 None  
 Other \_\_\_\_\_

Readiness to Quit:

- Not interested in quitting  
 Thinking about quitting at some point  
 Ready to quit

### Payment for Counseling

Medicare currently pays for tobacco cessation counseling for patients who use tobacco and have a "disease or an adverse health effect that has been found by the U.S. Surgeon General to be linked to tobacco use, or who [are] taking a therapeutic agent whose metabolism or dosing is affected by tobacco use." For a list of CPT, & ICD-9 Codes related to tobacco cessation counseling, click on the Ask and Act practice toolkit at [www.askandact.org](http://www.askandact.org).

## Assessment and Plan

### Counseling:

Counseled for:

- Three minutes or less  
 3 to 10 minutes  
 10+ minutes

Counseled for secondhand smoke

Counseling notes: \_\_\_\_\_  
\_\_\_\_\_

Handouts provided:

- "Prescription:" Quit Smoking  
 Quitline Card  
 Quit Smoking Brochure  
 Secondhand Smoke Brochure  
 Stop Smoking Guide  
 Familydoctor.org information  
 Other: \_\_\_\_\_

### Pharmacotherapy:

Recommended OTC:

- NRT Gum  
 NRT Lozenge  
 NRT Patch

Medical Treatment:

- NRT Nasal Spray  
*Dosing: 1–2 doses/hour (8–40 doses/day); one dose = one spray in each nostril; each spray delivers 0.5 mg of nicotine*
- NRT Inhaler  
*Dosing: 6–16 cartridges/day; initially use 1 cartridge q 1–2 hours*
- Bupropion SR  
*Dosing: Begin -2 weeks prior to quit date; 150 mg po q AM x 3 days, then increase to 150 mg po bid. Contraindications: head injury, seizures.*
- Varenicline  
*Dosing: Begin 1 week prior to quit date; days 1–3: 0.5 mg po q AM; days 4–7: 0.5 mg po bid; weeks 2–12: 1 mg po bid. Screen for suicidal ideations*

### Follow Up Plan

- Follow up visit in 2 weeks  
 Staff to follow up in \_\_\_ weeks  
 Address at next visit