

Diagnosing the Cause of Chest Pain

WILLIAM E. CAYLEY, JR., M.D., *Eau Claire Family Medicine Residency, Eau Claire, Wisconsin*

Chest pain presents a diagnostic challenge in outpatient family medicine. Noncardiac causes are common, but it is important not to overlook serious conditions such as an acute coronary syndrome, pulmonary embolism, or pneumonia. In addition to a thorough history and physical examination, most patients should have a chest radiograph and an electrocardiogram. Patients with chest pain that is predictably exertional, with electrocardiogram abnormalities, or with cardiac risk factors should be evaluated further with measurement of troponin levels and cardiac stress testing. Risk of pulmonary embolism can be determined with a simple prediction rule, and a D-dimer assay can help determine whether further evaluation with helical computed tomography or venous ultrasound is needed. Fever, egophony, and dullness to percussion suggest pneumonia, which can be confirmed with chest radiograph. Although some patients with chest pain have heart failure, this is unlikely in the absence of dyspnea; a brain natriuretic peptide level measurement can clarify the diagnosis. Pain reproducible by palpation is more likely to be musculoskeletal than ischemic. Chest pain also may be associated with panic disorder, for which patients can be screened with a two-item questionnaire. Clinical prediction rules can help clarify many of these diagnoses. (*Am Fam Physician* 2005;72:2012-21. Copyright © 2005 American Academy of Family Physicians.)



Chest pain is the chief complaint in about 1 to 2 percent of outpatient visits,¹ and although the cause is often noncardiac, heart disease remains the leading cause of death in the United States.² Thus, distinguishing between serious and benign causes of chest pain is imperative, and diagnostic and prognostic questions are important in making this determination.

The epidemiology of chest pain differs markedly between outpatient and emergency settings. Cardiovascular conditions such as myocardial infarction (MI), angina, pulmonary embolism (PE), and heart failure are found in more than 50 percent of patients presenting to the emergency department with chest pain,³ but the most common causes of chest pain seen in outpatient primary care are musculoskeletal conditions, gastrointestinal disease, stable coronary artery disease (CAD), panic disorder or other psychiatric conditions, and pulmonary disease (*Table 1*).^{3,4} Unstable CAD rarely is the cause of chest pain in primary care, and around 15 percent of chest pain episodes never reach a

definitive diagnosis.^{3,4} Despite these figures, when evaluating chest pain in primary care it is important to consider serious conditions such as stable or unstable CAD, PE, and pneumonia, in addition to more common (but less serious) conditions such as chest wall pain, peptic ulcer disease, gastroesophageal reflux disease (GERD), and panic disorder.

Clinical Diagnosis

Chest pressure with dyspnea commonly leads physicians and other health care professionals to consider an acute coronary syndrome such as unstable angina or MI, but these symptoms also may represent chest wall pain or PE. Dyspnea is common in patients with heart failure, whereas dyspnea with fever is characteristic of pneumonia and bronchitis. The usual descriptions of peptic ulcer disease and GERD include epigastric discomfort and retrosternal burning, but often it is difficult to distinguish clearly between classic “heartburn” and classic “chest pressure.” Although it often is thought that symptoms of anxiety can help distinguish pulmonary diseases from other causes of chest pain, this is not a

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
Determining whether chest pain is anginal, atypical anginal, or non-anginal is recommended to help determine a patient's cardiac risk.	C	16
The Rouan decision rule is recommended to help predict which patients are at higher risk of MI.	C	17
A Wells score of less than 2 plus a normal D-Dimer assay should rule out PE.	A	20, 32, 33
In patients with an abnormal D-Dimer assay or a Wells score indicating moderate to high risk, helical CT and lower extremity venous ultrasound examination should be used to rule in or rule out PE.	A	33, 35
The Diehr diagnostic rule is recommended to predict the likelihood of pneumonia based on clinical findings.	A	11
Patients should be screened for panic disorder using two set questions.	C	14
Patients presenting with chest pain should have an ECG evaluation for ST segment elevation, Q waves, and conduction defects. Results should be compared with previous tracings.	C	7, 9
Serum troponin-level testing is recommended to aid in the diagnosis of MI and help predict the likelihood of death or recurrent MI within 30 days.	C	25, 28, 29
Patients with chest pain and a negative initial cardiac evaluation should have further testing with stress ECG, perfusion scanning, or angiography depending on their level of risk.	C	16
The Duke treadmill score is recommended to help predict long-term prognosis for patients undergoing stress ECG testing.	A	31

MI = myocardial infarction; PE = pulmonary embolism; CT = computed tomography; ECG = electrocardiography.

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, see page 1949 or <http://www.aafp.org/afpsort.xml>.

TABLE 1
Epidemiology of Chest Pain in Primary Care and Emergency Department Settings

<i>Diagnosis*</i>	<i>Percentage of patients presenting with chest pain</i>		
	<i>Primary care: United States⁴</i>	<i>Primary care: Europe³</i>	<i>Emergency department³</i>
Musculoskeletal condition	36	29	7
Gastrointestinal disease	19	10	3
Serious cardiovascular disease†	16	13	54
Stable coronary artery disease	10	8	13
Unstable coronary artery disease	1.5	—	13
Psychosocial or psychiatric disease	8	17	9
Pulmonary disease‡	5	20	12
Nonspecific chest pain	16	11	15

*—Diagnoses are listed in order of prevalence in United States.

†—Including infarction, unstable angina, pulmonary embolism, and heart failure.

‡—Including pneumonia, pneumothorax, and lung cancer.

Adapted with permission from Klinkman MS, Stevens D, Gorenflo DW. Episodes of care for chest pain: a preliminary report from MIRNET. *J Fam Pract* 1994;38:349, with additional information from reference 3.

TABLE 2
Accuracy of Chest Pain Diagnosis Using the History and Physical Examination

Diagnosis* (overall outpatient probability)	Clinical finding	LR+	LR-	Probability of diagnosis (%) if finding is:	
				Present	Absent
Myocardial infarction (2%) ⁴	Chest pain radiates to both arms ⁵	7.10	0.67	13	1
	Hypotension ⁶	3.80	0.96	7	2
	S ₃ gallop ⁷	3.20	0.88	6	2
	Diaphoresis ^{8,9}	2.00	0.64	4	1
	Pleuritic chest pain ⁷	0.17	1.20	<1	2
	Palpation of tender area reproduces chest pain ⁸	0.16	1.20	<1	2
Pneumonia (5%) ^{10,11}	Egophony ¹¹	8.60	0.96	31	5
	Dullness to percussion ¹⁰	4.30	0.79	18	4
	Fever ¹¹	2.10	0.71	10	4
Heart failure (2%) ¹²	Exertional dyspnea ¹³	1.20	0	2	<1
	Displaced apical impulse ¹³	17.00	0.35	26	1
Panic disorder (8%) ⁴	"Yes" on at least one item of Autonomic Nervous System Questionnaire ^{14†}	1.30	0.60	10	1
Chest wall pain (36%) ⁴	Palpation of tender area reproduces chest pain ¹⁵	12.00	0.78	87	30

LR+ = positive likelihood ratio; LR- = negative likelihood ratio.

*—Diagnoses are listed in order of clinical importance.

†—Screening questions: (1) "In the past six months, did you ever have a spell or an attack when all of a sudden you felt frightened, anxious, or very uneasy?" and (2) "In the past six months, did you ever have a spell or an attack when for no reason your heart suddenly began to race, you felt faint, or you could not catch your breath?"¹⁴

Information from references 4 through 15.

consistent finding and should not be relied upon. There is enough overlap among the clinical manifestations of different causes of chest pain to make "classic" symptoms unhelpful in differentiating among diagnoses and ruling out serious causes. However, there are several validated clinical

decision rules that combine key groups of symptoms.

HISTORY AND PHYSICAL EXAMINATION

It is important to obtain a clear history of the onset and evolution of chest pain, with particular attention to details such as location, quality, duration, and aggravating or alleviating factors. Certain key symptoms and clinical findings can help rule in or out specific diagnoses (Table 2).⁴⁻¹⁵

Determining whether pain is (1) substernal, (2) provoked by exertion, or (3) relieved by rest or nitroglycerin helps to clarify whether it is typical anginal pain (has all three characteristics), atypical anginal pain (has two characteristics), or nonanginal

The Author

WILLIAM E. CAYLEY, JR., M.D., M.DIV., is assistant professor at the Eau Claire (Wis.) Family Medicine Residency, University of Wisconsin, Eau Claire. He received his medical degree from the Medical College of Wisconsin, Milwaukee, and completed a residency at the Eau Claire Family Medicine Residency, Eau Claire.

Address correspondence to William E. Cayley, Jr., M.D., M.Div., Eau Claire Family Medicine Residency, 617 W. Clairemont, Eau Claire, WI 54701 (e-mail: bcayley@yahoo.com). Reprints are not available from the author.

TABLE 3
Rouan Decision Rule
for Myocardial Infarction

Clinical Characteristics

Age greater than 60 years
Diaphoresis
History of MI or angina
Male sex
Pain described as pressure
Pain radiating to arm, shoulder, neck, or jaw

Score*	Risk of MI (%)
0	Up to 0.6
1	Up to 3.4
2	Up to 4.8
3	Up to 12.0
4	Up to 26.0

MI = myocardial infarction.

*—One point for each clinical characteristic.

NOTE: At no level of risk can MI be completely ruled out.

Information from reference 17.

pain (has one characteristic). Anginal chest pain has a high risk for CAD in all age groups; atypical anginal chest pain carries intermediate risk for CAD in women older than 50 years and in all men; and nonanginal chest pain carries intermediate risk for CAD in women older than 60 years and men older than 40 years.¹⁶

The likelihood of MI is higher if there is pain radiating to both arms,⁵ hypotension,⁶ an S₃ gallop on physical examination,⁷ or diaphoresis.^{8,9} Other factors predicting MI include age greater than 60 years, male sex, and prior MI.¹⁷ MI is less likely if pain is sharp or pleuritic.⁷ If the pain is reproducible by palpation of a specific tender area, the likelihood of MI decreases⁸ but the likelihood of chest wall pain increases.¹⁵ A history of rheumatoid arthritis or osteoarthritis also increases the likelihood of chest wall pain.¹⁸ The Rouan decision rule reliably predicts which patients with chest pain and a normal or nonspecific electrocardiogram (ECG) are

at higher risk for MI (Table 3).¹⁷ However, because up to 3 percent of patients initially diagnosed with a noncardiac cause of chest pain suffer death or MI within 30 days of presentation, patients with cardiac risk factors such as male sex, greater age, diabetes, hyperlipidemia, prior CAD, or heart failure warrant close follow-up.¹⁹

There are no individual signs or symptoms that reliably diagnose PE, but the simplified Wells scoring system²⁰ (Table 4^{20,21}) is well validated for determining whether patients have low, moderate, or high likelihood of PE,²⁰⁻²² and this guides further evaluation.

Findings that suggest pneumonia include fever, egophony, and dullness to percussion, but their absence does not rule out the diagnosis.¹⁰ Although chest pain in patients with chronic obstructive pulmonary disease and at least four previous acute exacerbations of chronic bronchitis is more likely to be caused by a recurrent exacerbation of bronchitis or pneumonia,²³ these patients are also at greater risk for CAD or acute coronary syndrome. The Diehr diagnostic rule, developed in a large study¹¹ from 1984, uses seven clinical findings to predict the likelihood of pneumonia (Table 5¹¹).

Although heart failure alone is an uncommon cause of chest pain, it may accompany acute coronary syndrome, valvular disease, or MI. A displaced apical impulse and a history of MI also support this diagnosis. Almost all patients with heart failure have exertional dyspnea, so the absence of exertional dyspnea is helpful in ruling out this diagnosis.¹³

Two simple questions¹⁴ are a highly sensitive screen for panic disorder:

- “In the past six months, did you ever have a spell or an attack when all of a sudden you felt frightened, anxious, or very uneasy?”
- “In the past six months, did you ever have a spell or an attack when for no reason your heart suddenly began to race, you felt faint, or you couldn’t catch your breath?”¹⁴

A “yes” on either item is a positive screen,

The most common causes of chest pain seen in outpatient primary care are musculoskeletal conditions, gastrointestinal disease, stable coronary artery disease, panic disorder or other psychiatric conditions, and pulmonary disease.

TABLE 4

Wells Model for Clinical Diagnosis of Pulmonary Embolism

Clinical finding*	Points
Clinical signs and symptoms of DVT (i.e., objectively measured leg swelling or pain with palpation of deep leg veins)	3.0
PE as likely or more likely than an alternative diagnosis	3.0
Heart rate more than 100 beats per minute	1.5
Immobilization (i.e., bedrest except for bathroom access for at least three consecutive days) or surgery in the past four weeks	1.5
Previous objectively diagnosed DVT or PE	1.5
Hemoptysis	1.0
Malignancy (treatment for cancer that is ongoing, within the past six months, or palliative)	1.0

Total points	Risk of PE	LR+ ²¹	Probability of PE (%) ²¹
<2 points	Low	0.13	1 to 28
2 to 6 points	Moderate	1.82	28 to 40
>6 points	High	6.75	38 to 91

*—Findings are listed in order of clinical importance.

DVT = deep venous thrombosis; PE = pulmonary embolism; LR+ = positive likelihood ratio.

Adapted with permission from Wells PS, Anderson DR, Rodger M, Ginsberg JS, Kearon C, Gent M, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000;83:418, with additional information from reference 21.

TABLE 5

Diehr Rule for Diagnosing Pneumonia in Adults with Acute Cough

Finding	Points
Rhinorrhea	-2
Sore throat	-1
Myalgia	1
Night sweats	1
Sputum all day	1
Respiratory rate more than 25 breaths per minute	2
Temperature more than 100°F (37.8°C)	2

Total points	Probability of pneumonia (%; overall probability = 3)
-3	0.0
-2	0.7
-1	1.6
0	2.2
1	8.8
2	10.3
3	25.0
≥4	29.4

Adapted with permission from Diehr P, Wood RW, Bushyhead J, Krueger L, Wolcott B, Tompkins RK. Prediction of pneumonia in outpatients with acute cough—a statistical approach. *J Chronic Dis* 1984;37:220.

and a “no” on both items makes panic disorder unlikely. However, neither these questions nor a general clinical impression are specific enough to allow a definite diagnosis of anxiety-related noncardiac chest pain, and a positive screen should not preclude further cardiac testing in patients with cardiac risk factors.¹⁹

Gastrointestinal disease can cause chest pain, but the history and physical examination are relatively inaccurate for ruling in or ruling out serious gastrointestinal pathology,²⁴ and it is important first to rule out immediately life-threatening cardiovascular and pulmonary causes of chest pain.

Diagnostic Testing

Once the clinical examination has narrowed the differential diagnosis, diagnostic testing helps determine whether the patient has a serious condition (Table 6).^{4,7,12,25,26} Most adults with chest pain should have at least an ECG and a chest radiograph, unless the history and physical examination suggest

an obviously nonthreatening cause of chest discomfort.

ACUTE CORONARY SYNDROME AND CAD

Important diagnostic tests when evaluating for acute coronary syndrome include the 12-lead ECG, serum markers of myocardial damage, and cardiac testing with stress testing or nuclear imaging. ECG findings that most strongly suggest MI are ST segment elevation, Q waves, and a conduction defect, especially if such findings are new compared with a previous ECG. New T-wave inversion also increases the likelihood of MI.^{7,9} However, none of these findings is sensitive enough that its absence can exclude MI.

The most common markers of myocardial damage are creatine kinase, the MB isoenzyme of creatine kinase (CK-MB), troponin T, and troponin I. A CK-MB level greater than 6.0 ng per mL (6.0 mcg per L) within nine hours of presentation for emergency care modestly increases the likelihood of MI or death in the next 30 days.²⁷ Elevated

TABLE 6
Accuracy of Chest Pain Diagnosis Using Diagnostic and Prognostic Tests

Diagnosis* (overall outpatient probability)	Clinical finding	LR+	LR-	Probability of diagnosis (%) if finding is:	
				Present	Absent
Myocardial infarction (2%) ⁴	New ST elevation ⁷	16.0	0.52	25	1
	New Q wave ⁷	8.7	0.68	15	1
	New conduction defect ⁷	6.3	0.88	11	2
	Any ST segment elevation ⁷	11.0	0.45	18	1
	Any Q wave ⁷	3.9	0.60	7	1
	Any conduction defect ⁷	2.7	0.89	5	2
	New T-wave inversion ⁷	2.5	0.72	5	1
	Troponin T >2 ng per mL (2 mcg per L) at least eight hours from presentation ²⁵	5.2	0.05	10	<1
Heart failure (2%) ¹²	Troponin I >1 ng per mL (1 mcg per L) at least six hours from presentation ²⁵	18.0	0.01	27	<1
	Abnormal electrocardiogram ²⁶	38.0	0.36	44	1
	Abnormal BNP level (cutoff 80 pg per mL [1 ng per L]) ¹²	3.6	0.10	7	<1

LR+ = positive likelihood ratio; LR- = negative likelihood ratio; BNP = brain natriuretic peptide.

*—Diagnoses are listed in order of clinical importance.

Information from references 4, 7, 12, 25, and 26.

levels of either troponin T (i.e., higher than 2 ng per mL [2 mcg per L]) at least eight hours from presentation or troponin I (i.e., higher than 1 ng per mL [1 mcg per L]) at least six hours from presentation support the diagnosis of MI or acute coronary syndrome and increase the likelihood of death or recurrent MI within 30 days. A normal level of troponin T or troponin I between six and 72 hours after the onset of chest pain is strong evidence against MI and acute coronary syndrome, particularly if the ECG is normal or near normal.^{25,28} In one study²⁹ of 773 patients who each presented to an emergency department with chest pain and had a normal ECG, researchers found that only 0.3 percent of those with a normal troponin I at six hours and 1.1 percent of those with a normal troponin T at six hours experienced acute MI or death in the 30 days following presentation. Thus, individuals with chest pain who have a history that indicates low risk of cardiovascular disease, a normal or near-normal ECG, and normal troponin levels can safely be evaluated as outpatients.

Patients at low risk usually do not need further testing unless there are other risk factors

in their family or medical history that markedly increase their likelihood of CAD. Patients at intermediate risk for CAD who can exercise and have no left bundle branch block, preexcitation, or significant resting ST depression on their ECG can be evaluated with an exercise stress ECG. Patients with baseline ECG abnormalities should have perfusion imaging performed along with a stress ECG, and patients who cannot exercise may be evaluated with a pharmacologic stress or vasodilator test (e.g., dobutamine [Dobutrex], adenosine [Adenocard]). Patients at high risk for CAD generally should proceed directly to angiography, which allows definitive assessment of coronary artery anatomy for patients in whom other testing is nondiagnostic and for patients who could benefit from revascularization.³⁰

For patients undergoing stress ECG testing, the Duke treadmill score (Table 7³¹) provides helpful prognostic information. Among 1,466 patients with a normal resting ECG, and 939 patients with ST-T abnormalities on a resting ECG, low-, intermediate-, and high-risk Duke treadmill scores accurately predicted seven-year survival rates for all-cause mortality.³¹

PULMONARY EMBOLISM

D-dimer testing has become an important part of the evaluation for PE and deep venous thrombosis (DVT). Quantitative enzyme-linked immunosorbent antibody assay (ELISA) D-dimer assays are more sensitive and have been more thoroughly tested in clinical settings than whole-blood agglutination assays.³² A low clinical suspicion for PE (e.g., Wells score less than 2) plus a normal quantitative ELISA D-dimer assay safely rules out PE, with a negative predictive value greater than 99.5 percent.^{20,32,33} If further testing is needed, helical computed

tomography (CT), combined with clinical suspicion and other testing such as lower extremity venous ultrasound, can be used to rule in or rule out PE.^{33,34} A number of different sequential testing protocols have been proposed, all of which involve the same basic elements: (1) for patients with low clinical suspicion and a normal D-dimer, no further evaluation or treatment is needed unless symptoms change or progress; (2) for patients with low clinical suspicion and an abnormal D-dimer, or moderate to high clinical suspicion, helical CT and lower extremity venous ultrasound examination should be ordered; (3) for patients with moderate or

high clinical suspicion and an abnormal CT scan or venous ultrasound result, treatment should be given for PE or DVT regardless of D-dimer; and (4) for patients with an abnormal D-dimer plus a normal CT scan and a normal venous ultrasound result, serial ultrasound should be considered if clinical suspicion is low to moderate, and pulmonary angiography should be considered if clinical suspicion is high.^{33,35} Patients in whom PE initially is ruled out by such an approach and who do not receive treatment have a less than 1 percent risk for PE occurring over the subsequent three months.³³ An encounter form that takes this approach appears in the February 1, 2004, issue of *American Family Physician* and can be accessed online at <http://www.aafp.org/afp/20040201/pocform.html>.³⁶

PNEUMONIA AND HEART FAILURE

Chest radiograph generally is considered the reference standard for patients suspected of having pneumonia, and it is the standard against which clinical evaluations for pneumonia are compared.¹⁰ An abnormal ECG and cardiomegaly on chest radiograph increase the likelihood of heart failure among patients with chest pain,²⁶ and brain natriuretic peptide (also known as B-type natriuretic peptide) level has been found to be reliable for detecting heart failure in patients presenting with acute dyspnea. Brain natriuretic peptide level is particularly helpful for ruling in heart failure if it is more than 500 pg per mL (500 ng per L), and for ruling out heart failure if it is less than 100 pg per mL (100 ng per L).^{14,37}

CHEST WALL PAIN

Chest wall pain usually can be diagnosed by history and physical examination if other etiologies have been excluded. Measurement of the sedimentation rate generally is not helpful in making the diagnosis¹⁸; in unusual situations, radiography may be helpful.³⁸

Recommended Diagnostic Strategy

An algorithm illustrating the discussed diagnostic strategy is provided in *Figure 1*.^{4,5,7-12, 14-17,20-22,25,26,28,29,32-35} When a patient presents with new chest pain, a typical or an atypical

The Wells score combined with a D-dimer assay, helical computed tomography, and lower extremity ultrasound examination can safely diagnose pulmonary embolism in many patients.

TABLE 7
Duke Treadmill Score

Duke treadmill score:

$$\text{Exercise time (minutes)} - (5 \times \text{ST deviation [mm]}) - (4 \times \text{angina index}^*)$$


Score	Risk	Seven-year survival rate (%)	
		Normal resting ECG	ST-T abnormalities on resting ECG
>5	Low	95	91
-10 to 4	Medium	91	80
<-11	High	78	78

ECG = electrocardiogram.

*—No angina during testing = 0, typical angina = 1, test stopped because of angina = 2.

Information from reference 31.

Outpatient Diagnosis of Chest Pain



NOTE: This algorithm combines and simplifies diagnostic recommendations from multiple sources to provide an overview, and does not represent a validated decision rule.

Figure 1. Algorithm for the outpatient diagnosis of causes of chest pain. (ECG = electrocardiography; CT = computed tomography.)

Information from references 4, 5, 7 through 12, 14 through 17, 20 through 22, 25, 26, 28, 29, and 32 through 35.

anginal pattern, pain radiation or diaphoresis, cardiac risk factors, or ischemic ECG changes, serial measurement of troponins should be considered to determine whether hospitalization or outpatient evaluation with stress testing is warranted. If the probability of PE is low, based on the Wells score, a negative D-dimer result eliminates the need for further testing; an abnormal D-dimer or moderate to high probability of PE should prompt helical CT and venous ultrasound examination to guide further management. Fever, egophony, or dullness to percussion should prompt evaluation for pneumonia with chest radiograph. If life-threatening causes of chest pain are ruled out, then a history of spontaneous anxiety, palpitations, faintness, or dyspnea suggests panic disorder. A history of exertional dyspnea and a displaced apical impulse should prompt investigation for heart failure. Gastrointestinal symptoms should prompt further evaluation.

Data Sources: The PubMed database was searched using the following terms: chest pain, angina, acute myocardial infarction, coronary artery disease, heart failure, pulmonary embolism, chest wall pain, bronchitis, pneumonia, and peptic ulcer disease. Titles were reviewed to identify literature relevant to the outpatient diagnosis of chest pain. Additional searches were performed using the following databases: InfoPOEMs (<http://www.infopeoms.com>), Agency for Healthcare Research and Quality (<http://www.ahrq.gov>), Cochrane Collaboration (<http://www.cochrane.org>), Database of Abstracts of Reviews of Effects (<http://www.york.ac.uk/inst/crd/darehp.htm>), and Institute for Clinical Systems Improvement (<http://www.icsi.org>).

Author disclosure: Nothing to disclose.

REFERENCES

- Woodwell DA. National ambulatory medical care survey: 1998 summary. *Adv Data* 2000;19:1-26.
- National Center for Health Statistics. Health, United States, 2003: with chartbook on trends in the health of Americans. Hyattsville, Md.: Department of Health and Human Services, CDC, National Center for Health Statistics, 2003.
- Buntinx F, Knockaert D, Bruyninckx R, de Blaey N, Aerts M, Knottnerus JA, et al. Chest pain in general practice or in the hospital emergency department: is it the same? *Fam Pract* 2001;18:586-9.
- Klinkman MS, Stevens D, Gorenflo DW. Episodes of care for chest pain: a preliminary report from MIRNET. *J Fam Pract* 1994;38:345-52.
- Berger JP, Buclin R, Haller E, Van Melle G, Yersin B. Right arm involvement and pain extension can help to differentiate coronary diseases from chest pain of other origin: a prospective emergency ward study of 278 consecutive patients admitted for chest pain. *J Intern Med* 1990;227:165-72.
- Weaver WD, Eisenberg S, Martin JS, Litwin PE, Shaef-fer SM, Ho MT, et al. Myocardial Infarction Triage and Intervention Project—phase I: patient characteristics and feasibility of prehospital initiation of thrombolytic therapy. *J Am Coll Cardiol* 1990;15:925-31.
- Tierney WM, Fitzgerald J, McHenry R, Roth BJ, Psaty B, Stump DL, et al. Physicians' estimates of the probability of myocardial infarction in emergency room patients with chest pain. *Med Decis Making* 1986;6:12-7.
- Solomon CG, Lee TH, Cook EF, Weisberg MC, Brand DA, Rouan GW, et al. Comparison of clinical presentation of acute myocardial infarction in patients older than 65 years of age to younger patients: the Multicenter Chest Pain Study experience. *Am J Cardiol* 1989;63:772-6.
- Panju AA, Hemmelgarn BR, Guyatt GH, Simel DL. The rational clinical examination. Is this patient having a myocardial infarction? *JAMA* 1998;280:1256-63.
- Metlay JP, Kapoor WN, Fine MJ. Does this patient have community-acquired pneumonia? Diagnosing pneumonia by history and physical examination. *JAMA* 1997;278:1440-5.
- Diehr P, Wood RW, Bushyhead J, Krueger L, Wolcott B, Tompkins RK. Prediction of pneumonia in outpatients with acute cough—a statistical approach. *J Chronic Dis* 1984;37:215-25.
- Maisel AS, Krishnaswamy P, Nowak RM, McCord J, Hollander JE, Duc P, et al. Rapid measurement of B-type natriuretic peptide in the emergency diagnosis of heart failure. *N Engl J Med* 2002;347:161-7.
- Davie AP, Francis CM, Caruana FL, Sutherland GR, McMurray JJ. Assessing diagnosis in heart failure: which features are any use? *QJM* 1997;90:335-9.
- Stein MB, Roy-Byrne PP, McQuaid JR, Laffaye C, Russo J, McCahill ME, et al. Development of a brief diagnostic screen for panic disorder in primary care. *Psychosom Med* 1999;61:359-64.
- Wise CM, Semble EL, Dalton CB. Musculoskeletal chest wall syndromes in patients with non-cardiac chest pain: a study of 100 patients. *Arch Phys Med Rehabil* 1992;73:147-9.
- Gibbons RJ, Balady GJ, Bricker JT, Chaitman BR, Fletcher GF, Froelicher VF, et al. ACC/AHA 2002 guideline update for exercise testing: summary article: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1997 Exercise Testing Guidelines). *Circulation* 2002;106:1883-92.
- Rouan GW, Lee TH, Cook EF, Brand DA, Weisberg MC, Goldman L. Clinical characteristics and outcome of acute myocardial infarction in patients with initially normal or nonspecific electrocardiograms (a report from the Multicenter Chest Pain Study). *Am J Cardiol* 1989;64:1087-92.
- Disla E, Rhim HR, Reddy A, Karten I, Taranta A. Costochondritis. A prospective analysis in an emergency department setting. *Arch Intern Med* 1994;154:2466-9.
- Miller CD, Lindsell CJ, Khandelwal S, Chandra A, Pollack CV, Tiffany BR, et al. Is the initial diagnostic impression of "noncardiac chest pain" adequate to exclude cardiac disease [published correction appears in *Ann Emerg Med* 2005;45:87]? *Ann Emerg Med* 2004;44:565-74.

20. Wells PS, Anderson DR, Rodger M, Ginsberg JS, Kearon C, Gent M, et al. Derivation of a simple clinical model to categorize patients probability of pulmonary embolism: increasing the models utility with the SimpliRED D-dimer. *Thromb Haemost* 2000;83:416-20.
21. Tamariz LJ, Eng J, Segal JB, Krishnan JA, Bolger DT, Streiff MD, et al. Usefulness of clinical prediction rules for the diagnosis of venous thromboembolism: a systematic review. *Am J Med* 2004;117:676-84.
22. Chunilal SD, Eikelboom JW, Attia J, Miniati M, Panju AA, Simel DL, et al. Does this patient have pulmonary embolism? *JAMA* 2003;290:2849-58.
23. Ball P, Harris JM, Lowson D, Tillotson G, Wilson R. Acute infective exacerbations of chronic bronchitis. *QJM* 1995;88:61-8.
24. Meineche-Schmidt V, Jorgensen T. "Alarm symptoms" in patients with dyspepsia: a three-year prospective study from general practice. *Scand J Gastroenterol* 2002;37:999-1007.
25. Ebell MH, Flewelling D, Flynn CA. A systematic review of troponin T and I for diagnosing acute myocardial infarction. *J Fam Pract* 2000;49:550-6.
26. Talreja D, Gruver C, Sklenar J, Dent J, Kaul S. Efficient utilization of echocardiography for the assessment of left ventricular systolic function. *Am Heart J* 2000;139:394-8.
27. McCord J, Nowak RM, Hudson MP, McCullough PA, Tomlanovich MC, Jacobsen G, et al. The prognostic significance of serial myoglobin, troponin I, and creatine kinase-MB measurements in patients evaluated in the emergency department for acute coronary syndrome. *Ann Emerg Med* 2003;42:343-50.
28. Ebell MH, White LL, Weismantel D. A systematic review of troponin T and I values as a prognostic tool for patients with chest pain. *J Fam Pract* 2000;49:746-53.
29. Hamm CW, Goldmann BU, Heeschen C, Kreymann G, Berger J, Meinertz T. Emergency room triage of patients with acute chest pain by means of rapid testing for cardiac troponin T or troponin I. *N Engl J Med* 1997;337:1648-53.
30. Gibbons RJ, Abrams J, Chatterjee K, Daley J, Deedwania PC, Douglas JS, et al. ACC/AHA 2002 guideline update for the management of patients with chronic stable angina: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Committee to Update the 1999 Guidelines for the Management of Patients with Chronic Stable Angina). *J Am Coll Cardiol* 2003;41:159-68.
31. Kwok JM, Miller TD, Christian TF, Hodge DO, Gibbons RJ. Prognostic value of a treadmill exercise score in symptomatic patients with nonspecific ST-T abnormalities on resting ECG. *JAMA* 1999;282:1047-53.
32. Stein PD, Hull RD, Patel KC, Olson RE, Ghali WA, Brant R, et al. d-dimer for the exclusion of acute venous thrombosis and pulmonary embolism: a systematic review. *Ann Intern Med* 2004;140:589-602.
33. Perrier A, Roy PM, Aujesky D, Chagnon I, Howarth N, Gourdier AL, et al. Diagnosing pulmonary embolism in outpatients with clinical assessment, D-dimer measurement, venous ultrasound, and helical computed tomography: a multicenter management study. *Am J Med* 2004;116:291-9.
34. Musset D, Parent F, Meyer G, Maitre S, Girard P, Leroyer C, et al. Diagnostic strategy for patients with suspected pulmonary embolism: a prospective multicentre outcome study. *Lancet* 2002;360:1914-20.
35. Institute for Clinical Systems Improvement. Healthcare guidelines. Venous thromboembolism. Accessed online September 21, 2005, at: http://www.icsi.org/display_file.asp?FileId=187&title=Venous%20Thromboembolism%20.
36. Ebell M. Suspected pulmonary embolism: evidence-based diagnostic testing. *Am Fam Physician* 2004;69:601.
37. Cardarelli R, Lumicao TG Jr. B-type natriuretic peptide: a review of its diagnostic, prognostic, and therapeutic monitoring value in heart failure for primary care physicians. *J Am Board Fam Pract* 2003;16:327-33.
38. Meyer CA, White CS. Cartilaginous disorders of the chest. *Radiographics* 1998;18:1109-23.