

Treatment of the Common Cold in Children and Adults

JULIA FASHNER, MD; KEVIN ERICSON, MD; and SARAH WERNER, DO

St. Joseph Family Medicine Residency, Mishawaka, Indiana

The common cold, or upper respiratory tract infection, is one of the leading reasons for physician visits. Generally caused by viruses, the common cold is treated symptomatically. Antibiotics are not effective in children or adults. In children, there is a potential for harm and no benefits with over-the-counter cough and cold medications; therefore, they should not be used in children younger than four years. Other commonly used medications, such as inhaled corticosteroids, oral prednisolone, and *Echinacea*, also are ineffective in children. Products that improve symptoms in children include vapor rub, zinc sulfate, *Pelargonium sidoides* (geranium) extract, and buckwheat honey. Prophylactic probiotics, zinc sulfate, nasal saline irrigation, and the herbal preparation Chizukit reduce the incidence of colds in children. For adults, antihistamines, intranasal corticosteroids, codeine, nasal saline irrigation, *Echinacea angustifolia* preparations, and steam inhalation are ineffective at relieving cold symptoms. Pseudoephedrine, phenylephrine, inhaled ipratropium, and zinc (acetate or gluconate) modestly reduce the severity and duration of symptoms for adults. Nonsteroidal anti-inflammatory drugs and some herbal preparations, including *Echinacea purpurea*, improve symptoms in adults. Prophylactic use of garlic may decrease the frequency of colds in adults, but has no effect on duration of symptoms. Hand hygiene reduces the spread of viruses that cause cold illnesses. Prophylactic vitamin C modestly reduces cold symptom duration in adults and children. (*Am Fam Physician*. 2012;86(2):153-159. Copyright © 2012 American Academy of Family Physicians.)



ILLUSTRATION BY TODD BUCK

► **Patient information:** Handouts on treating the common cold, written by the authors of this article, are available at <http://www.aafp.org/afp/2012/0715/p153-s1.html> and <http://www.aafp.org/afp/2012/0715/p153-s2.html>. Access to the handouts is free and unrestricted. Let us know what you think about AFP putting handouts online only; e-mail the editors at afpcomment@aafp.org.

The common cold, or upper respiratory tract infection, usually is caused by one of several respiratory viruses, most commonly rhinovirus. These viruses, which concentrate in nasal secretions, are easily transmitted through sneezing, coughing, or nose blowing. Signs and symptoms of the common cold include fever, cough, rhinorrhea, nasal congestion, sore throat, headache, and myalgias.

Patients seek care for cold symptoms during all seasons of the year, with cough being the third most common and nasal congestion the 15th most common presenting symptom among all office visits.¹ The common cold is the third most common primary diagnosis in office visits.¹ Colds are self-limited, usually lasting up to 10 days; therefore, management is directed at symptom relief rather than treating the infection. Multiple remedies,

including complementary and alternative medicine products, over-the-counter products, and prescription drugs, have been used to prevent and treat cold symptoms.

When medications are requested, physicians play an important role in educating patients about the treatment choices. Many familiar prescription cough and cold medications were removed from the market in early 2011 because the U.S. Food and Drug Administration had not evaluated them for safety, effectiveness, or quality.² Physicians should caution patients about over-the-counter and complementary and alternative medicine products because manufacturers are not required to prove claims of therapeutic benefit.

Children

Cold and cough medications are among the top 20 substances leading to death in children

Table 1. Therapies Not Effective for the Common Cold in Children

<i>Therapy</i>	<i>Evidence</i>	<i>Findings</i>
Antibiotics	Cochrane review of four studies ⁷	No difference in persistence of symptoms for the common cold or acute purulent rhinitis compared with placebo
Carbocysteine	Cochrane review of three RCTs ¹³	No significant difference in cough, dyspnea, or overall general health compared with placebo
Dextromethorphan	One cohort study ¹²	Not superior to placebo in nocturnal cough or sleep quality in the child or parents
Diphenhydramine (Benadryl)	One cohort study ¹²	Not superior to placebo in nocturnal cough or sleep quality in the child or parents
<i>Echinacea purpurea</i>	Cochrane review of two RCTs ¹⁰	No difference in severity of symptoms, peak of symptom severity, number of days of fever, or parental report of severity score compared with placebo
Low-dose inhaled corticosteroids	Cochrane review of two studies ⁸	No decrease in the number of episodes requiring oral corticosteroids, emergency department visits, hospital admissions, the frequency of wheezing, or duration of episodes
Oral prednisolone	One RCT of a five-day course ⁹	No significant difference in duration of hospitalization, interval between admission and discharge, mean seven-day symptom score reported by a parent, or hospital readmission for wheezing within one month compared with placebo
OTC antihistamines	Cochrane review of two studies ¹¹	No more effective than placebo for cough
OTC antihistamine with decongestant	Cochrane review of two studies ¹¹	No more effective than placebo for cough
OTC antitussives	Cochrane review of three studies ¹¹	No more effective than placebo for cough
OTC antitussive and bronchodilator	Cochrane review of one study ¹¹	No more effective than placebo for cough
Vitamin C	Not studied in children ¹⁴	—

OTC = over-the-counter; RCT = randomized controlled trial.

Information from references 7 through 14.

younger than five years.³ In 2008, the U.S. Food and Drug Administration recommended that over-the-counter cough and cold medications be avoided in children younger than two years.⁴ After the removal of over-the-counter infant cough and cold medications from pharmacy shelves, the estimated number of emergency department visits for adverse events involving these medications was cut in half for children younger than two years.⁵ Manufacturers of these medications have voluntarily modified the product labels to state that they should not be used in children younger than four years.⁶

INEFFECTIVE INTERVENTIONS

Prescription and Over-the-Counter Products. Because viruses cause most colds, antibiotics are ineffective.⁷ Low-dose inhaled corticosteroids⁸ and oral prednisolone⁹ do not improve outcomes in children without asthma. *Echinacea* products also are ineffective for

treating cold symptoms in children.¹⁰ There is no evidence to support the use of most over-the-counter cough remedies in children.^{11,12} *Table 1* summarizes findings of studies on these medications.⁷⁻¹⁴

Fluids. Caregivers are often advised to increase a child's fluid intake. However, in two case series and a prevalence study, some children with respiratory infections but no signs of dehydration developed hyponatremia with increased fluids.¹⁵ Therefore, extra fluid intake is not advised in children because of potential harm.

EFFECTIVE INTERVENTIONS

Table 2 summarizes therapies that may be effective in children with the common cold.^{8,13,16-20}

Complementary and Alternative Medicine Products. Several of these therapies provide relief from cold symptoms. Vapor rub applied to the chest and neck has been shown to improve cough severity and quality of sleep for

the child and parents, but it has a strong smell that children may not tolerate.¹⁹ Studies regarding therapeutic use of zinc sulfate show a trend toward decreased duration of cold symptoms when it is taken within the first 24 hours of symptom onset.²⁰ Adverse effects, such as bad taste and nausea, are more common with zinc lozenges than with syrup or tablets.²⁰ *Pelargonium sidoides* (geranium) extract (Umcka Coldcare) may help resolve cough and sputum production in children with the common cold.¹⁸ Buckwheat honey is superior to placebo for reducing frequency of cough, reducing bothersome cough, and improving quality of sleep for the child.¹⁶ Honey should not be used in children younger than one year because of the risk of botulism.

Nasal Irrigation and Acetylcysteine. During acute illness, nasal irrigation with saline can help alleviate sore throat, thin nasal secretions, and improve nasal breathing and can reduce the need for nasal decongestants and mucolytics.¹⁷ A systematic review of six trials published in the 1990s found that acetylcysteine (commonly used

in Europe, but not in the United States, as a mucolytic) may decrease cough after six to seven days of therapy in children older than two years.¹³ The main adverse effect of acetylcysteine is vomiting.

Inhaled Corticosteroids. Some children with viral cold symptoms also develop wheezing. Although low-dose corticosteroids are ineffective in these children, one review of high-dose inhaled corticosteroids found a trend toward decreased frequency of wheezing episodes that require oral corticosteroids, the duration of episodes, and the number of physician visits.⁸

PROPHYLAXIS

Table 3 summarizes therapies that may be effective for cold prophylaxis in children.^{14,17,20-22}

Complementary and Alternative Medicine Products. Some of these products may help prevent colds if taken regularly. Probiotics, such as *Lactobacillus acidophilus* NCFM, alone or combined with *Bifidobacterium animalis*, taken by healthy children during the winter may

Table 2. Therapies That May Be Effective for the Common Cold in Children

Therapy	Age of children studied	Dosing	Duration of treatment
Acetylcysteine ¹³	0 to 18 years	Variable	Variable, up to 28 days
High-dose inhaled corticosteroids in children who are wheezing ⁸	One to five years	Budesonide (Pulmicort), 1,600 mcg by MDI with nebulizer or 3,200 mcg by MDI with nebulizer and face mask, if needed	Until asymptomatic for 24 hours
	One to five years	Beclomethasone, 2,250 mcg daily by MDI	Five days
	One to three years	Budesonide 1,600 mcg by MDI with nebulizer and face mask for first three days, then 800 mcg for another seven days	Total of 10 days
Honey (buckwheat) ¹⁶	Two to five years	2.5 mL	Once
	Six to 11 years	5 mL	Once
	12 to 18 years	10 mL	Once
Nasal irrigation with saline ¹⁷	Six to 10 years	3 to 9 mL per nostril	Up to three weeks
<i>Pelargonium sidoides</i> (geranium) extract (Umcka Coldcare) ¹⁸	One to 18 years	10 to 30 drops (depending on age)	Seven days
Vapor rub ¹⁹	Two to five years	5 mL	Once
	Six to 11 years	10 mL	Once
Zinc sulfate ²⁰	One to 10 years	Syrup, 15 mg per 5 mL	10 days

MDI = metered dose inhaler.

Information from references 8, 13, and 16 through 20.

Table 3. Therapies That May Be Effective for Common Cold Prophylaxis in Children

Therapy	Age of children studied	Dosing	Duration of treatment
Chizukit ²¹	One to three years	5 mL twice daily	12 weeks
	Four to five years	7.5 mL twice daily	12 weeks
Nasal irrigation with saline ¹⁷	Six to 10 years	3 to 9 mL per nostril three times daily	Nine weeks
Probiotics* ²²	Three to five years	1 g (1 × 10 ¹⁰ colony-forming units) mixed with 120 mL of 1% milk twice daily	Six months
Vitamin C ¹⁴	< 12 years	0.2 to 2 g daily	Two weeks to nine months
Zinc sulfate ²⁰	One to 10 years	Syrup, 15 mg per 5 mL daily	Seven months
	6.5 to 16 years	Tablet, 10 mg daily	Six days per week for five months

*—*Lactobacillus acidophilus* NCFM, alone or combined with *Bifidobacterium animalis*.

Information from references 14, 17, and 20 through 22.

reduce day care absences; the incidence of fever, cough, and rhinorrhea; and the use of antibiotics.²²

A Cochrane review showed a 13 percent decrease in cold symptoms in children who took 1 g of vitamin C daily before illness, although optimal duration of treatment to achieve these benefits is unknown.¹⁴ Zinc sulfate used prophylactically for at least five months reduces the incidence of viral colds, absences from school, and antibiotic use in children.²⁰

The herbal preparation Chizukit contains 50 mg per mL of *Echinacea*, 50 mg per mL of propolis, and 10 mg per mL of vitamin C.²¹ In a randomized, placebo-controlled trial of 430 children one to five years of age, Chizukit decreased the number of cold episodes, the number of days the child was ill, and the number of days the child missed school. It also decreased the need for antipyretics and antibiotics; physician visits; and episodes of otitis media, pneumonia, and tonsillitis. However, children may not comply with taking the product because of its unpleasant taste.²¹

Nasal Saline Irrigation. Nasal irrigation with saline as a preventive measure in children is better than standard treatment for multiple cold symptoms. Overall, the treatment decreases illness and nasal secretions, improving nasal breathing. These children also use fewer antipyretics, nasal decongestants, and mucolytics and have fewer school absences.¹⁷

Adults

INEFFECTIVE INTERVENTIONS

Table 4 summarizes studies of medications that are ineffective for the common cold in adults.^{7,11,14,23-28}

Antibiotics and Antihistamines. In adults, as in children, antibiotics do not decrease the duration or severity of illness, even when purulent rhinitis is present.⁷ Sedating and nonsedating antihistamines are ineffective for cough and other cold symptoms.^{11,23,29}

Opioids, Intranasal Corticosteroids, and Nasal Saline Irrigation. Despite widespread use, codeine is no more effective than placebo for reducing cough.^{11,24} The American College of Chest Physicians (ACCP) does not recommend other opioids for the treatment of cough.²⁴ Although intranasal corticosteroids reduce swelling and inflammation of the nasal mucosa, they have not been shown to significantly benefit patients with the common cold.^{26,27} Nasal irrigation with hypertonic or normal saline does not provide significant relief for cold symptoms in adults.²⁸

Complementary and Alternative Medicine Products. When used solely for treatment of symptoms after they appear, vitamin C does not consistently reduce their duration or severity.¹⁴ Herbal preparations containing *Echinacea angustifolia* are not beneficial.²⁵ Many physicians have recommended increased fluid intake and inhalation of heated, humidified air to thin secretions during a cold. No randomized trials have assessed the effect of increasing fluid intake in adults,³⁰ and a Cochrane review found inconsistent study results for steam inhalation.³¹

EFFECTIVE INTERVENTIONS

Decongestants With or Without Antihistamines. Oral or topical decongestants alone seem to be somewhat effective for short-term relief of cold symptoms, compared with placebo.³² Pseudoephedrine and phenylephrine decrease nasal edema to improve air intake.³² Although antihistamines do not work as monotherapy, combination medications containing a first-generation antihistamine and decongestant may be slightly beneficial in relieving general symptoms, nasal symptoms,²³ and cough.¹¹ Combination medications are recommended by the ACCP to treat acute cough.²⁹

Anticholinergics, Dextromethorphan, Guaiifenesin. Ipratropium (Atrovent) is the only orally inhaled

Table 4. Therapies Not Effective for the Common Cold in Adults

<i>Therapy</i>	<i>Evidence</i>	<i>Findings</i>
Antibiotics	Cochrane review of nine RCTs ⁷	No difference in symptoms or purulent rhinitis compared with placebo
Antihistamine monotherapy (sedating and nonsedating)	Cochrane review of three RCTs ¹¹ Cochrane review of 32 RCTs ²³	No more effective than placebo No more effective than placebo
Codeine	Cochrane review of two RCTs ¹¹ American College of Chest Physicians ²⁴	No more effective than placebo for cough Not recommended
<i>Echinacea angustifolia</i>	RCT with viral challenge ²⁵	No more effective than placebo for cold symptoms
Intranasal corticosteroids	Two RCTs ^{26,27}	No more effective than placebo
Nasal irrigation with hypertonic or normal saline	One RCT ²⁸	No more effective than observation
Vitamin C	Cochrane review of seven RCTs ¹⁴	No more effective than placebo for reducing duration or severity of cold symptoms

RCT = randomized controlled trial.

Information from references 7, 11, 14, and 23 through 28.

Table 5. CAM Products That May Be Effective for the Common Cold in Adults

<i>Preparation</i>	<i>Dosing</i>	<i>Duration of treatment</i>
Treatment		
<i>Andrographis paniculata</i> (Kalmcold) ^{35,36}	200 mg daily	Five days
<i>Echinacea purpurea</i> (solution of pressed juice of aerial parts and alcohol) ¹⁰	4 mL twice daily 20 drops every two hours on day 1, then 20 drops three times daily	Eight weeks 10 days
<i>Pelargonium sidoides</i> (geranium) extract (Umcka Coldcare) ^{18,37}	30 drops three times daily, alcohol root extract	10 days
Zinc acetate or gluconate ²⁰	Variable (lozenges contain between 4.5 and 23.7 mg of zinc)	As long as symptoms persist
Prophylaxis		
Garlic ³⁸	Supplement with 180 mg of allicin	12 weeks
Vitamin C ¹⁴	0.25 to 2 g daily	40 days to 28 weeks (generally around three months)

CAM = complementary and alternative medicine.

Information from references 10, 14, 18, 20, and 35 through 38.

anticholinergic recommended by the ACCP for cough caused by a common cold,²⁴ and one study showed that the nasal formulation decreases rhinorrhea and sneezing.³³ Studies of dextromethorphan and guaifenesin for cough are almost evenly split, with some demonstrating benefit and others not.^{11,24}

Nonsteroidal Anti-inflammatory Drugs. These medications effectively relieve pain from headache, myalgias, and arthralgias experienced during a cold; however, decreased sneezing is the only effect they have

on respiratory symptoms.³⁴ The ACCP has concluded that naproxen (Naprosyn) is beneficial in the treatment of acute cough.²⁴

Complementary and Alternative Medicine Products. Table 5 summarizes the herbal preparations that may be effective in adults.^{10,14,18,20,35-38} An herbal solution containing *P. sidoides* was shown to reduce the duration and severity of 10 different cold symptoms in a randomized controlled trial.³⁷ Another randomized controlled trial demonstrated the benefit of *Andrographis paniculata*

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	References
Antibiotics should not be used for the treatment of cold symptoms in children or adults.	A	7
Over-the-counter cough and cold medications should not be used in children younger than four years because of potential harms and lack of benefit.	B	4, 6, 11
Treatment with buckwheat honey, <i>Pelargonium sidoides</i> (geranium) extract (Umcka Coldcare), nasal saline irrigation, vapor rub, or zinc sulfate may decrease cold symptoms in children.	B	16-20
Codeine is not effective for cough in adults.	A	11, 24
Antihistamine monotherapy (sedating and nonsedating) does not improve cold symptoms in adults.	A	11, 23, 29
Decongestants, antihistamine/decongestant combinations, and intranasal ipratropium (Atrovent) may improve cold symptoms in adults.	B	11, 23, 32, 33
Nonsteroidal anti-inflammatory drugs reduce pain secondary to upper respiratory tract infection in adults.	A	34
<i>Andrographis paniculata</i> (Kalmcold) and <i>P. sidoides</i> may reduce severity and duration of cold symptoms in adults.	B	35-37

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort.xml>.

(Kalmcold) in improving symptom scores.³⁵ A systematic review also indicated that *A. paniculata*, alone or in combination with *Acanthopanax senticosus*, may be more effective for symptom relief than placebo.³⁶

Early use of *Echinacea purpurea* shortens duration and decreases severity of cold symptoms; preparations with the aerial parts versus the flowering parts are most effective.¹⁰ Although dosages and preparations of zinc are not standardized, a Cochrane review showed that starting zinc lozenges (acetate or gluconate) within the first 24 hours of symptom onset reduces the severity and duration of illness.²⁰ Adverse effects of zinc include bad taste and nausea.²⁰ Intranasal zinc should not be used because it may result in the permanent loss of smell.³⁹

PROPHYLAXIS

Few medications have been shown to be beneficial in preventing the common cold in adults (Table 5^{10,14,18,20,35-38}). The prophylactic use of vitamin C does not reduce the incidence of colds, but decreases illness duration by 8 percent.¹⁴ Limited, poor-quality studies of garlic show a decrease in the number of self-reported colds, but no decrease in days to recovery. Adverse effects from garlic included bad odor and skin rash.³⁸

Frequent hand washing can reduce the spread of respiratory viruses in all ages and can reduce transmission from children to other household members.⁴⁰ In a large meta-analysis, the benefits of antibacterial and nonantibacterial soaps were not significantly different.⁴¹

Benzalkonium chloride-based hand sanitizers that foam and leave a residue have a protective effect against colds. Alcohol hand sanitizers are less effective.⁴¹

Data Sources: A search of Essential Evidence Plus was completed using the key words cold and respiratory tract infections. This search included InfoPOEMs, Cochrane reviews, and practice guidelines. We also searched Dynamed and the U.S. Food and Drug Administration Web site for specific information regarding changes in recommendations for the use of cough and cold medications in children. Search dates: March 22, 2011, to April 6, 2011.

The Authors

JULIA FASHNER, MD, FAAFP, is an associate director at the St. Joseph Family Medicine Residency, Mishawaka, Ind.

KEVIN ERICSON, MD, FAAFP, is an associate director at the St. Joseph Family Medicine Residency.

SARAH WERNER, DO, is a third-year resident at the St. Joseph Family Medicine Residency.

Address correspondence to Julia Fashner, MD, FAAFP, St. Joseph Family Medicine Residency, 611 E. Douglas Rd., Ste. 412, Mishawaka, IN 46545 (e-mail: fashnerj@sjrmc.com). Reprints are not available from the authors.

Author disclosure: No relevant financial affiliations to disclose.

REFERENCES

1. Hsiao CJ, Cherry DK, Beatty PC, Rechtsteiner EA. National Ambulatory Medical Care Survey: 2007 summary. *Natl Health Stat Report*. 2010;(27):1-32.
2. FDA prompts removal of unapproved drugs from market [news release]. Silver Springs, Md.: U.S. Food and Drug Administration; March 2, 2011.

- <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/ucm245048.htm>. Accessed April 6, 2011.
- Bronstein AC, Spyker DA, Cantilena LR Jr, Green JL, Rumack BH, Giffin SL. 2009 annual report of the American Association of Poison Control Centers' National Poison Data System (NPDS): 27th annual report. *Clin Toxicol (Phila)*. 2010;48(10):979-1178.
 - FDA releases recommendations regarding use of over-the-counter cough and cold products [news release]. Silver Springs, Md.: U.S. Food and Drug Administration; January 17, 2008. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/2008/ucm116839.htm>. Accessed April 6, 2011.
 - Shehab N, Schaefer MK, Kegler SR, Budnitz DS. Adverse events from cough and cold medications after a market withdrawal of products labeled for infants. *Pediatrics*. 2010;126(6):1100-1107.
 - FDA statement following CHPA's announcement on nonprescription over-the-counter cough and cold medicines in children [news release]. Silver Springs, Md.: U.S. Food and Drug Administration; October 8, 2008. <http://www.fda.gov/NewsEvents/Newsroom/PressAnnouncements/2008/ucm116964.htm>. Accessed April 6, 2011.
 - Arroll B, Kenealy T. Antibiotics for the common cold and acute purulent rhinitis. *Cochrane Database Syst Rev*. 2005;(3):CD000247.
 - McKean M, Ducharme F. Inhaled steroids for episodic viral wheeze of childhood. *Cochrane Database Syst Rev*. 2000;(2):CD001107.
 - Panickar J, Lakhanpaul M, Lambert PC, et al. Oral prednisolone for preschool children with acute virus-induced wheezing. *N Engl J Med*. 2009;360(4):329-338.
 - Linde K, Barrett B, Wölkart K, Bauer R, Melchart D. *Echinacea* for preventing and treating the common cold. *Cochrane Database Syst Rev*. 2006;(1):CD000530.
 - Smith SM, Schroeder K, Fahey T. Over-the-counter medications for acute cough in children and adults in ambulatory settings. *Cochrane Database Syst Rev*. 2008;(1):CD001831.
 - Paul IM, Yoder KE, Crowell KR, et al. Effect of dextromethorphan, diphenhydramine, and placebo on nocturnal cough and sleep quality for coughing children and their parents. *Pediatrics*. 2004;114(1):e85-e90.
 - Duijvestijn YC, Mourdi N, Smucny J, Pons G, Chalumeau M. Acetylcysteine and carbocysteine for acute upper and lower respiratory tract infections in paediatric patients without chronic broncho-pulmonary disease. *Cochrane Database Syst Rev*. 2009;(1):CD003124.
 - Douglas RM, Hemilä H, Chalker E, Treacy B. Vitamin C for preventing and treating the common cold. *Cochrane Database Syst Rev*. 2007;(3):CD000980.
 - Guppy MP, Mickan SM, Del Mar CB. "Drink plenty of fluids": a systematic review of evidence for this recommendation in acute respiratory infections. *BMJ*. 2004;328(7438):499-500.
 - Paul IM, Beiler J, McMonagle A, Shaffer ML, Duda L, Berlin CM Jr. Effect of honey, dextromethorphan, and no treatment on nocturnal cough and sleep quality for coughing children and their parents. *Arch Pediatr Adolesc Med*. 2007;161(12):1140-1146.
 - Slapak I, Skoupá J, Strnad P, Hornik P. Efficacy of isotonic nasal wash (seawater) in the treatment and prevention of rhinitis in children. *Arch Otolaryngol Head Neck Surg*. 2008;134(1):67-74.
 - Timmer A, Günther J, Rücker G, Motschall E, Antes G, Kern WV. *Pelargonium sidoides* extract for acute respiratory tract infections. *Cochrane Database Syst Rev*. 2008;(3):CD006323.
 - Paul IM, Beiler JS, King TS, Clapp ER, Vallati J, Berlin CM Jr. Vapor rub, petrolatum, and no treatment for children with nocturnal cough and cold symptoms. *Pediatrics*. 2010;126(6):1092-1099.
 - Singh M, Das RR. Zinc for the common cold. *Cochrane Database Syst Rev*. 2011;(2):CD001364.
 - Cohen HA, Varsano I, Kahan E, Sarrell EM, Uziel Y. Effectiveness of an herbal preparation containing echinacea, propolis, and vitamin C in preventing respiratory tract infections in children: a randomized, double-blind, placebo-controlled, multicenter study. *Arch Pediatr Adolesc Med*. 2004;158(3):217-221.
 - Leyer GJ, Li S, Mubasher ME, Reifer C, Ouwehand AC. Probiotic effects on cold and influenza-like symptom incidence and duration in children. *Pediatrics*. 2009;124(2):e172-e179.
 - Sutter AI, Lemiengre M, Campbell H, Mackinnon HF. Antihistamines for the common cold. *Cochrane Database Syst Rev*. 2003;(3):CD001267.
 - Bolser DC. Cough suppressant and pharmacologic protussive therapy: ACCP evidence-based clinical practice guidelines. *Chest*. 2006;129(1 suppl):238S-249S.
 - Turner RB, Bauer R, Woelkart K, Hulsey TC, Gangemi JD. An evaluation of *Echinacea angustifolia* in experimental rhinovirus infections. *N Engl J Med*. 2005;353(4):341-348.
 - Puhakka T, Mäkelä MJ, Malmström K, et al. The common cold: effects of intranasal fluticasone propionate treatment. *J Allergy Clin Immunol*. 1998;101(6 pt 1):726-731.
 - Qvarnberg Y, Valtonen H, Laurikainen K. Intranasal beclomethasone dipropionate in the treatment of common cold. *Rhinology*. 2001;39(1):9-12.
 - Adam P, Stiffman M, Blake RL Jr. A clinical trial of hypertonic saline nasal spray in subjects with the common cold or rhinosinusitis. *Arch Fam Med*. 1998;7(1):39-43.
 - Pratter MR. Cough and the common cold: ACCP evidence-based clinical practice guidelines. *Chest*. 2006;129(1 suppl):72S-74S.
 - Guppy MP, Mickan SM, Del Mar CB, Thorning S, Rack A. Advising patients to increase fluid intake for treating acute respiratory infections. *Cochrane Database Syst Rev*. 2011;(2):CD004419.
 - Singh M, Singh M. Heated, humidified air for the common cold. *Cochrane Database Syst Rev*. 2011;(5):CD001728.
 - Taverner D, Latte J. Nasal decongestants for the common cold. *Cochrane Database Syst Rev*. 2007;(1):CD001953.
 - Hayden FG, Diamond L, Wood PB, Korts DC, Wecker MT. Effectiveness and safety of intranasal ipratropium bromide in common colds. A randomized, double-blind, placebo-controlled trial. *Ann Intern Med*. 1996;125(2):89-97.
 - Kim SY, Chang YJ, Cho HM, Hwang YW, Moon YS. Non-steroidal anti-inflammatory drugs for the common cold. *Cochrane Database Syst Rev*. 2009;(3):CD006362.
 - Saxena RC, Singh R, Kumar P, et al. A randomized double blind placebo controlled clinical evaluation of extract of *Andrographis paniculata* (KalmCold) in patients with uncomplicated upper respiratory tract infection. *Phytomedicine*. 2010;17(3-4):178-185.
 - Poolsup N, Suthisang C, Prathanturug S, Asawamekin A, Chanchareon U. *Andrographis paniculata* in the symptomatic treatment of uncomplicated upper respiratory tract infection: systematic review of randomized controlled trials. *J Clin Pharm Ther*. 2004;29(1):37-45.
 - Lizogub VG, Riley DS, Heger M. Efficacy of a *Pelargonium sidoides* preparation in patients with the common cold: a randomized, double blind, placebo-controlled clinical trial. *Explore (NY)*. 2007;3(6):573-584.
 - Lissiman E, Bhasale AL, Cohen M. Garlic for the common cold. *Cochrane Database Syst Rev*. 2009;(3):CD006206.
 - Institute for Clinical Systems Improvement. Health care guideline: diagnosis and treatment of respiratory illness in children and adults. January 2011. http://www.icsi.org/respiratory_illness_in_children_and_adults__guideline_/respiratory_illness_in_children_and_adults__guideline__13116.html. Accessed March 24, 2011.
 - Jefferson T, Del Mar C, Dooley L, et al. Physical interventions to interrupt or reduce the spread of respiratory viruses. *Cochrane Database Syst Rev*. 2010;(1):CD006207.
 - Aiello AE, Coulborn RM, Perez V, Larson EL. Effect of hand hygiene on infectious disease risk in the community setting: a meta-analysis. *Am J Public Health*. 2008;98(8):1372-1381.