

Trauma in Pregnancy: Assessment, Management, and Prevention

NEIL J. MURPHY, MD, *Alaska Native Medical Center, Anchorage, Alaska*

JEFFREY D. QUINLAN, MD, *Uniformed Services University of the Health Sciences, Bethesda, Maryland*

Trauma complicates one in 12 pregnancies, and is the leading nonobstetric cause of death among pregnant women. The most common traumatic injuries are motor vehicle crashes, assaults, falls, and intimate partner violence. Nine out of 10 traumatic injuries during pregnancy are classified as minor, yet 60% to 70% of fetal losses after trauma are a result of minor injuries. In minor trauma, four to 24 hours of tocodynamometric monitoring is recommended. Ultrasonography has low sensitivity, but high specificity, for placental abruption. The Kleihauer-Betke test should be performed after major trauma to determine the degree of fetomaternal hemorrhage, regardless of Rh status. To improve the effectiveness of cardiopulmonary resuscitation, clinicians should perform left lateral uterine displacement by tilting the whole maternal body 25 to 30 degrees. Unique aspects of advanced cardiac life support include early intubation, removal of all uterine and fetal monitors, and performance of perimortem cesarean delivery. Proper seat belt use reduces the risk of maternal and fetal injuries in motor vehicle crashes. The lap belt should be placed as low as possible under the protuberant portion of the abdomen and the shoulder belt positioned off to the side of the uterus, between the breasts and over the midportion of the clavicle. All women of childbearing age should be routinely screened for intimate partner violence. (*Am Fam Physician*. 2014;90(10):717-722. Copyright © 2014 American Academy of Family Physicians.)



More online
at <http://www.aafp.org/afp>.

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz Questions on page 696.

Author disclosure: No relevant financial affiliations.

► **Patient information:** A handout on this topic, written by the authors of this article, is available at <http://www.aafp.org/afp/2014/1115/p717-s1.html>.

Family physicians, regardless of their involvement in intrapartum care, often evaluate pregnant women after minor accidents, falls, or motor vehicle crashes. Trauma complicates one in 12 pregnancies, and is the leading nonobstetric cause of death among pregnant women.¹⁻³ Traumatic injuries to pregnant women are unintentional (motor vehicle crashes [48%], falls [25%], poisonings, and burns) or intentional (assaults/intimate partner violence [IPV; 17%], suicide [3.3%], homicide, and gunshot wounds [4%]).^{2,4} Injuries are classified as minor or major trauma. By convention, minor trauma does not involve the abdomen, rapid compression, deceleration, or shearing forces, and the patient does not report pain, vaginal bleeding, loss of fluid, or decreased fetal movement.⁵ Nine out of 10 traumatic injuries during pregnancy are classified as minor. However, of fetal losses associated with trauma, 60% to 70% follow minor injuries.⁴

Anatomic and physiologic changes of pregnancy influence the assessment, management, and prevention of trauma.^{3,6} Physiologic changes include a 30% to 50% increase in

blood volume and a 40% to 50% increase in respiratory rate.⁷ Anatomic changes include elevated diaphragm, delayed gastric emptying, and progressive uterine growth. During the first trimester, the thick-walled uterus is well protected from trauma by the pelvic girdle. In the second trimester, relatively abundant amniotic fluid volume protects the fetus. By the third trimester, however, the now thin-walled and prominent uterus is exposed to blunt and penetrating abdominal trauma. The placenta is an inelastic organ attached to an elastic organ (the uterus). Placental abruption may occur when trauma involving acceleration and deceleration deforms the uterus and shears the placenta off its implantation site.

Assessment

The most common pregnancy-related traumas are minor incidents, including motor vehicle crashes and mild blunt abdominal trauma. These situations require clinical judgment about the extent of maternal and fetal assessment. However, because insignificant trauma can result in fetal injury or demise, fetal monitoring recommendations for pregnant women with minor trauma

Trauma in Pregnancy

are similar to those for women with major trauma.⁸⁻¹⁰ In patients who have experienced IPV, an assessment for depression and suicidality should accompany assessment for immediate safety.¹¹

In major trauma, the primary assessment should focus on airway, breathing, and circulation. Once the primary assessment has been completed, a secondary assessment should include obstetric and nonobstetric injuries and fetal well-being.

A placental abruption may become apparent shortly after the injury. Fetal monitoring in women who experience trauma at greater than 20 weeks' gestation should be initiated as soon as the patient is stabilized. The patient should be monitored by tocodynamometry for a minimum of four hours.^{8,12} Eight or more contractions per hour were present in the first four hours of monitoring in 100% of patients with an ultimate diagnosis of placental abruption.^{8,9} If at least six contractions per hour are detected during the four-hour initial period, or the mechanism of injury presents a high degree of fetal risk,¹³ then 24 hours of monitoring is indicated.^{3,12} Before viability (23 to 24 weeks' gestation), the fetal heart should be monitored via intermittent Doppler auscultation, and after viability, continuous fetal monitoring should be performed.

Ultrasonography has poor sensitivity (24%) for detection of placental abruption. However, it is very specific (96%), resulting in a positive predictive value of 88% if abruption is seen and a negative predictive value of 53% if abruption is not seen.¹⁴

The Kleihauer-Betke test allows identification of fetal blood cells. It should be performed in pregnant women who sustain major trauma to detect fetomaternal transfusion, regardless of Rh status, to determine the degree of fetomaternal hemorrhage.⁸ Although the Kleihauer-Betke test screens for the degree of trauma in all patients, the result is used only to dictate Rh immune globulin therapy in Rh-negative patients.

Management

Figure 1 provides an algorithmic approach to the management of trauma during pregnancy.³

MINOR TRAUMA

Management of minor trauma is limited to care of lacerations or fractures, discussion of appropriate analgesics, counseling about the signs and symptoms of abruption, and ensuring appropriate follow-up. Tetanus

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	References
Pregnant women at greater than 20 weeks' gestation who have experienced trauma should be monitored for a minimum of four hours by tocodynamometry.	C	8, 12
The Kleihauer-Betke test should be performed in all pregnant women who sustain major trauma.	C	8, 12
Perimortem cesarean delivery after cardiac arrest may improve neonatal and maternal outcomes, and is not harmful.	C	16, 19, 20, 22
Pregnant women who are occupants in motor vehicles should wear lap and shoulder seat belts, and should not turn off air bags.	C	28, 29
All women of childbearing age should be screened for intimate partner violence.	B	31, 33

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.

toxoid should be administered if tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis (Tdap) vaccine has not already been administered during the current pregnancy. Appropriate referrals should be made to community-based advocacy groups for persons who have experienced IPV, and a plan should be made to ensure the safety of the patient and other vulnerable persons living in the household.¹¹

MAJOR TRAUMA

When feasible, patients who have experienced major trauma should be transported to a hospital that has the resources to perform a timely trauma evaluation.³ Because placental abruption has been reported to occur up to 24 hours following trauma, monitoring via tocodynamometry should be continued for a minimum of 24 hours if at least six uterine contractions have occurred or if there are nonreassuring fetal heart rate patterns, vaginal bleeding, significant uterine tenderness, serious maternal injury, or a positive Kleihauer-Betke test result.^{3,13} If none of these findings are present, the patient may be discharged home with precautions.³ Table 1 provides evaluation and discharge criteria for blunt trauma in pregnancy.^{9,12}

Between 24 and 34 weeks' gestation, if delivery appears imminent, 12 mg of betamethasone should be administered intramuscularly to promote fetal lung maturity, and repeated in 24 hours. All pregnant Rh-negative patients should receive Rh immune globulin therapy unless the injury is remote from the uterus (e.g., isolated distal extremity).¹²

CARDIOPULMONARY RESUSCITATION

Because of maternal physiologic changes, left lateral uterine displacement is required during cardiopulmonary

Management of Trauma in Pregnancy




Figure 1. Management of trauma in pregnancy. (bpm = beats per minute; FHR = fetal heart rate.)

Adapted with permission from Mendez-Figueroa H, Dahlke JD, Vrees RA, Rouse DJ. Trauma in pregnancy: an updated systematic review. *Am J Obstet Gynecol.* 2013;209(1):6. <http://www.sciencedirect.com/science/journal/00029378>.

resuscitation. This is performed by tilting the whole maternal body 25 to 30 degrees (*Figure 2*¹⁵), or by manual uterine displacement (*Figure 3*¹⁵). Manual uterine displacement may allow for more effective chest compressions because it avoids the need to facilitate a total body tilt.¹⁶ All standard medications and defibrillation regimens may be used in advanced cardiac life support (ACLS).¹⁷ Unique aspects of ACLS include early

intubation, removal of all uterine and fetal monitors, and performance of a perimortem cesarean delivery.¹⁸ Modifications of resuscitative efforts in pregnancy are listed in *eTable A*.

PERIMORTEM CESAREAN DELIVERY

Perimortem cesarean delivery may save the life of the fetus when performed after 23 to 24 weeks' gestation.^{19,20}

Trauma in Pregnancy

Table 1. Management of Blunt Trauma in Pregnancy

Evaluation

Primary maternal and fetal survey
Laboratory: blood type, Rh factor test, hematocrit measurement, Kleihauer-Betke test, coagulation studies
Consider obstetric ultrasonography
If greater than 20 weeks' gestation, monitor for contractions
If fewer than three contractions per hour, monitor for four hours, then discharge
If three to seven contractions per hour, monitor for 24 hours, then discharge

Discharge criteria

Resolution of contractions
Reassuring fetal heart tracing
Intact membranes
No vaginal bleeding
No uterine tenderness
All Rh-negative patients should receive Rh immune globulin therapy unless the injury is remote from the uterus (e.g., isolated distal extremity)¹²

Information from references 9 and 12.

Delivery increases venous return and cardiac output by 25% to 30%, may lead to a clear survival benefit for the mother, and has not been shown to be harmful.^{16,19-22} The American Heart Association recommends considering

perimortem cesarean delivery if ACLS has not returned spontaneous circulation within four to five minutes of cardiac arrest.¹⁸ *Table 2 and eTable B* describe considerations based on gestational age and issues related to perimortem cesarean delivery.

Prevention

MOTOR VEHICLE CRASHES

Overall, 2% of pregnant women are involved in a motor vehicle crash during their pregnancy,⁷ and an estimated 368 pregnant women per year die as a result.²³ Motor vehicle crashes account for more than 50% of all trauma during pregnancy, with 82% of fetal deaths occurring during these crashes.²⁴

Seat belt use actually decreases during pregnancy, because women fear that the seat belt will hurt the fetus. In blunt trauma, the most common cause of fetal death is maternal death.⁹ Incorrect seat belt use increases the risk of intrauterine injury and fetal death.²⁵ When women were properly restrained, adverse fetal outcomes occurred in 29% of motor vehicle crashes. In women who were improperly restrained, adverse fetal outcomes occurred in 50% of motor vehicle crashes.²⁶ Proper seat belt use should be a major prenatal counseling issue. The lap belt should be placed as low as possible under the protuberant portion of the abdomen and the shoulder belt positioned off to the side of the uterus, between the breasts and over the midportion of the clavicle (*Figure 4*). Placement of the lap belt over the dome of the uterus




Figure 2. Left lateral uterine displacement by tilting the pregnant woman 25 to 30 degrees.

Reprinted with permission from *Advanced Life Support in Obstetrics Provider Syllabus*. Leawood, Kan.: American Academy of Family Physicians; 2010.



Figure 3. Manual uterine displacement.

Reprinted with permission from *Advanced Life Support in Obstetrics Provider Syllabus*. Leawood, Kan.: American Academy of Family Physicians; 2010.

Table 2. Gestational Age in Emergency Hysterotomy

Gestational age less than 20 weeks: emergency hysterotomy is not indicated for a single fetus, but it may be indicated for more than one fetus.

Gestational age 20 to 23 weeks: emergency hysterotomy is indicated to enable successful resuscitation but not for survival of the delivered fetus.

Gestational age greater than 23 weeks: emergency hysterotomy is indicated to save the life of both the mother and fetus.

Emergency hysterotomy should not be delayed while attempting to listen for fetal heart tones or to perform an ultrasonography to document gestational age. A uterine fundus measurement of 3 to 4 cm above the umbilicus correlates with a 23- to 24-week singleton gestation, and is a quick, easy assessment of gestational age that can be made in the field. Omission of an emergency hysterotomy, or delay in performing the procedure, may lead to the unnecessary loss of two lives, although the decision is multidimensional.

Personnel with the appropriate skills and equipment should provide this service, in or out of the hospital.



Figure 4. Correct way to wear a seat belt in pregnancy.

ILLUSTRATION BY JOHN KARAFEOU

significantly increases pressure transmission to the uterus and has been associated with significant uterine and fetal injury. There should not be excessive slack in either belt, and both the lap and shoulder restraints should be applied as snugly as comfort will allow.

Air bag deployment reduces injury to pregnant women and does not increase the risk of adverse pregnancy outcomes.²⁷ The American College of Obstetricians and Gynecologists (ACOG)²⁸ and the National Highway Traffic Safety Administration²⁹ state that pregnant women who are occupants in motor vehicles should wear lap and shoulder seat belts and should not turn off air bags.

INTIMATE PARTNER VIOLENCE

Direct assault on the abdomen may occur in the setting of IPV. As a result, ACOG and the American Academy of Pediatrics recommend universal screening for IPV during pregnancy.^{30,31} Likewise, the American Academy of Family Physicians states that all family physicians should be alert for physical and behavioral signs and symptoms associated with abuse or neglect.³² The U.S. Preventive Services Task Force recommends that all women of childbearing age be screened for IPV.³³ A review of this topic was previously published in the May 15, 2011, issue of *American Family Physician* (<http://www.aafp.org/afp/2011/0515/p1165.html>).

A patient who presents with a vague or inconsistent history of trauma should raise suspicion for battering.¹²

Table 3. Possible Indicators of Intimate Partner Violence

- Diminished self-image, depression, or suicide attempts
- Frequent emergency department or physician's office visits
- Injuries inconsistent with the stated history
- Partner insists on being present for interview and examination, and monopolizes discussion
- Progressive isolation from support systems
- Reproductive coercion
- Self-abuse
- Self-blame for injuries
- Symptoms suggestive of substance abuse
- Unwanted touching or fondling

Information from references 12 and 30.

Battering may escalate in pregnancy. The abdomen is the most common target for blows, kicks, and other assaults. *Table 3* lists other historical elements and signs that may indicate IPV.^{12,30} Identified patients should be appropriately counseled and referred.³⁰

Data Sources: Drawing on the literature search completed in October 2013 for the American Academy of Family Physicians' Advanced Life Support in Obstetrics Chapter K: Maternal Resuscitation, we searched the Cochrane Database of Systematic Reviews, the National Guideline Clearinghouse, and PubMed utilizing the key words trauma, motor vehicle crashes, pregnancy, abruption, and perimortem cesarean delivery, both individually and in combination. Search date: August 14, 2014.

Trauma in Pregnancy

The opinions and assertions contained herein are the private views of the authors and are not to be construed as official or as reflecting the views of the U.S. Army, Navy, or Air Force Medical Departments or the U.S. Army, Navy, Air Force, or Public Health Service.

This article is one in a series on "Advanced Life Support in Obstetrics (ALSO)," initially established by Mark Deutchman, MD, Denver, Colo. The series is now coordinated by Larry Leeman, MD, MPH, ALSO Managing Editor, Albuquerque, N.M.

The Authors

NEIL J. MURPHY, MD, is a staff member in the Dept. of Obstetrics and Gynecology at Alaska Native Medical Center in Anchorage, and an associate professor in the Dept. of Family Medicine at the University of Washington in Seattle.

JEFFREY D. QUINLAN, MD, is an associate professor and vice chair of the Dept. of Family Medicine at the Uniformed Services University of the Health Sciences in Bethesda, Md.

Address correspondence to Neil J. Murphy, MD, Alaska Native Medical Center, 4320 Diplomacy Dr., PCC-OB/GYN, Anchorage, AK 99508 (e-mail: njmurphy@southcentralfoundation.com). Reprints are not available from the authors.

REFERENCES

1. Mirza FG, Devine PC, Gaddipati S. Trauma in pregnancy: a systematic approach. *Am J Perinatol*. 2010;27(7):579-586.
2. El Kady D. Perinatal outcomes of traumatic injuries during pregnancy. *Clin Obstet Gynecol*. 2007;50(3):582-591.
3. Mendez-Figueroa H, Dahlke JD, Vrees RA, Rouse DJ. Trauma in pregnancy: an updated systematic review. *Am J Obstet Gynecol*. 2013; 209(1):1-10.
4. El-Kady D, Gilbert WM, Anderson J, Danielsen B, Townner D, Smith LH. Trauma during pregnancy: an analysis of maternal and fetal outcomes in a large population. *Am J Obstet Gynecol*. 2004;190(6):1661-1668.
5. Smith R, Crane P; Perinatal Joint Practice Committee. Post-trauma care in pregnancy. University of Michigan. February 2003. Renewed January 24, 2011. http://obgyn.med.umich.edu/sites/obgyn.med.umich.edu/files/internal_resources_clinical/trauma.pdf. Accessed November 29, 2013.
6. Brown S, Mozurkewich E. Trauma during pregnancy. *Obstet Gynecol Clin North Am*. 2013;40(1):47-57.
7. Muench MV, Canterino JC. Trauma in pregnancy. *Obstet Gynecol Clin North Am*. 2007;34(3):555-583, xiii.
8. Pearlman MD, Tintinalli JE, Lorenz RP. A prospective controlled study of outcome after trauma during pregnancy. *Am J Obstet Gynecol*. 1990;162(6):1502-1507, discussion 1507-1510.
9. Shah AJ, Kilcline BA. Trauma in pregnancy. *Emerg Med Clin North Am*. 2003;21(3):615-629.
10. Pearlman MD, Tintinalli JE, Lorenz RP. Blunt trauma during pregnancy. *N Engl J Med*. 1990;323(23):1609-1613.
11. Cronholm PF, Fogarty CT, Ambuel B, Harrison SL. Intimate partner violence. *Am Fam Physician*. 2011;83(10):1165-1172.
12. American College of Surgeons. Trauma in pregnancy and intimate partner violence. In: *Advanced Trauma Life Support Student Course Manual*. 9th ed. Chicago, Ill.: American College of Surgeons; 2012.
13. Dahmus MA, Sibai BM. Blunt abdominal trauma: are there any predictive factors for abruptio placentae or maternal-fetal distress? *Am J Obstet Gynecol*. 1993;169(4):1054-1059.
14. Glantz C, Purnell L. Clinical utility of sonography in the diagnosis and treatment of placental abruption. *J Ultrasound Med*. 2002;21(8): 837-840.
15. Advanced Life Support in Obstetrics Provider Syllabus. Leawood, Kan.: American Academy of Family Physicians; 2010.
16. Jeejeebhoy FM, Zelop CM, Windrim R, Carvalho JC, Dorian P, Morrison LJ. Management of cardiac arrest in pregnancy: a systematic review. *Resuscitation*. 2011;82(7):801-809.
17. Sinz E, Navarro K, Soderberg ES, Callaway CW; American Heart Association. *Advanced Cardiovascular Life Support: Provider Manual*. Dallas, Tex.: American Heart Association; 2011.
18. Vanden Hoek TL, Morrison LJ, Shuster M, et al. Part 12: cardiac arrest in special situations: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care [published corrections appear in *Circulation*. 2011;123(6):e239, and *Circulation*. 2011;124(15):e405]. *Circulation*. 2010;122(18 suppl 3): S829-S861.
19. Katz VL, Dotters DJ, Droegemueller W. Perimortem cesarean delivery. *Obstet Gynecol*. 1986;68(4):571-576.
20. Katz V, Balderston K, DeFreest M. Perimortem cesarean delivery: were our assumptions correct? *Am J Obstet Gynecol*. 2005;192(6):1916-1920, discussion 1920-1921.
21. Ueland K, Akamatsu TJ, Eng M, Bonica JJ, Hansen JM. Maternal cardiovascular dynamics. VI. Cesarean section under epidural anesthesia without epinephrine. *Am J Obstet Gynecol*. 1972;114(6):775-780.
22. Einav S, Kaufman N, Sela HY. Maternal cardiac arrest and perimortem caesarean delivery: evidence or expert-based? *Resuscitation*. 2012;83 (10):1191-1200.
23. Weiss H. Causes of traumatic death during pregnancy [letter]. *JAMA*. 2001;285(22):2854-2855.
24. Mattox KL, Goetzl L. Trauma in pregnancy. *Crit Care Med*. 2005;33(10 suppl):S385-S389.
25. Weinberg L, Steele RG, Pugh R, Higgins S, Herbert M, Story D. The pregnant trauma patient. *Anaesth Intensive Care*. 2005;33(2):167-180.
26. Klinich KD, Flannagan CA, Rupp JD, Sochor M, Schneider LW, Pearlman MD. Fetal outcome in motor-vehicle crashes: effects of crash characteristics and maternal restraint. *Am J Obstet Gynecol*. 2008;198(4):450. e1-9.
27. Schiff MA, Mack CD, Kaufman RP, Holt VL, Grossman DC. The effect of air bags on pregnancy outcomes in Washington State: 2002-2005. *Obstet Gynecol*. 2010;115(1):85-92.
28. American College of Obstetricians and Gynecologists. Car safety for you and your baby. <http://www.acog.org/~media/For%20Patients/faq018.pdf?dmc=1&ts=20140703T2121569354>. Accessed July 3, 2014.
29. National Highway Traffic Safety Administration. Buckle up in pregnancy. http://www.nhtsa.dot.gov/people/injury/airbags/Internet_Services_Group/ISG-Restricted/Buckle-Up%20America/pregnancybrochure_BUA_PregnancyNHTSAchange.pdf. Accessed July 3, 2014.
30. American College of Obstetricians and Gynecologists. Intimate partner violence. ACOG Committee Opinion No. 518. *Obstet Gynecol*. 2012; 119(2 pt 1):412-417.
31. AAP Committee on Fetus and Newborn and ACOG Committee on Obstetric Practice Editors. *Guidelines for Perinatal Care*. 7th ed. Elk Grove, Ill.: American Academy of Pediatrics; 2012:131-132, 246-248.
32. American Academy of Family Physicians. AAFP policies. Violence (position paper). <http://www.aafp.org/about/policies/all/violence.html>. Accessed August 14, 2014.
33. U.S. Preventive Services Task Force. Screening for intimate partner violence and abuse of elderly and vulnerable adults. January 2013. <http://www.uspreventiveservicestaskforce.org/Page/Document/RecommendationStatementFinal/intimate-partner-violence-and-abuse-of-elderly-and-vulnerable-adults-screening>. Accessed July 3, 2014.

Table A. Modifications of Resuscitative Efforts in Pregnancy

Action	Rationale
Basic life support	
Perform manual uterine displacement, or 25- to 30-degree left lateral tilt	Decreased aortocaval compression
Increase chest wall compression force	Decreased chest wall compliance with breast hypertrophy and diaphragmatic elevation
Use cricoid pressure, if assistance is available	Decreased gastric aspiration
Perform compressions higher on the sternum (slightly above center of sternum)	Elevated diaphragm and contents
Remove fetal and uterine monitors before defibrillation	Loss of adequate cardiac shock dose; produces skin burns at monitor sites
Heimlich maneuver; use chest thrust if unable to encircle the gravid abdomen	Enlarged uterus displaces diaphragm
Advanced cardiac life support	
Start intravenous therapy above diaphragm	Aortocaval compression
Early tracheal intubation; use short laryngoscope handle and smaller endotracheal tube	Difficult ventilation with pharyngeal edema, breast hypertrophy, diaphragmatic elevation
Consider other etiologies (e.g., magnesium toxicity)	Magnesium used as tocolytic therapy
Consider left wide paddle, adhesive pad, or breast displacement	Dextrorotation of the heart; breast hypertrophy
Verify endotracheal tube with carbon dioxide detector	Esophageal detector more likely not to reinflate after compression
Alter ventilation volumes and rates	Tailor ventilator support to oxygenation and ventilation
Perform emergency hysterotomy after four minutes	Decreased aortocaval and venous compression
Continue all maternal resuscitative efforts (cardiopulmonary resuscitation, positioning, defibrillation, and drugs) during and after cesarean delivery	Decreased aortocaval and venous compression

Information from:

Berg RA, Hemphill R, Abella BS, et al. Part 5: adult basic life support: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care [published correction appears in *Circulation*. 2011;124(15):e402]. *Circulation*. 2010;122(18 suppl 3):S685-S705.

Jeejeebhoy FM, Zelop CM, Windrim R, Carvalho JC, Dorian P, Morrison LJ. Management of cardiac arrest in pregnancy: a systematic review. *Resuscitation*. 2011;82(7):801-809.

Sinz E, Navarro K; American Heart Association. Cardiac arrest associated with pregnancy. In: *ACLS for Experienced Providers: Manual and Resource Text*. Dallas, Tex.: American Heart Association; 2013.

Vanden Hoek TL, Morrison LJ, Shuster M, et al. Part 12: cardiac arrest in special situations: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care [published corrections appear in *Circulation*. 2011;123(6):e239, and *Circulation*. 2011;124(15):e405]. *Circulation*. 2010;122(18 suppl 3):S829-S861.

eTable B. Perimortem Hysterotomy Issues

<i>Factors to consider</i>	<i>Discussion</i>
Cardiac arrest	
<p>If the mother fails to respond to initial resuscitative efforts and the gestational age is greater than 20 weeks, personnel and equipment should be assembled for emergency hysterotomy. This will allow simultaneous continuation of resuscitative efforts and preparation for the cesarean delivery.</p> <p>Is the mother receiving appropriate basic life support and advanced cardiac life support care, including:</p> <ul style="list-style-type: none"> CPR with compressions performed with the mother angled to the left? Early intubation with verification of proper placement of the endotracheal tube? Administration of indicated intravenous medications to a venous site above the diaphragm? <p>Has the mother responded to arrest interventions?</p> <p>Are there any potentially reversible causes of arrest?</p>	<p>Survival probabilities for the mother and fetus decrease as the interval from maternal arrest increases.</p> <p>Physicians should aim for an interval of five minutes or less from maternal arrest to delivery of the fetus. This goal requires efficient assembly of personnel and equipment.</p> <p>Physicians should not wait until five minutes of unsuccessful resuscitation have passed before beginning to consider the need to deliver the fetus emergently. The need for hysterotomy should be considered within minutes to enable assembly of personnel and equipment.</p> <p>Physicians should ensure that the mother has received superior resuscitative efforts. She cannot be declared “refractory” to CPR and advanced cardiac life support unless all interventions have been implemented and implemented well.</p>
Mother-infant factors	
<p>Is the fetus old enough to survive?</p> <p>Has too much time passed for the mother to survive?</p> <p>Is the mother’s cardiac arrest caused by a chronic hypoxic state?</p> <p>What is the status of the fetus at the time of the mother’s cardiac arrest?</p>	<p>Recognition of gestational age is critically important. Survival is unlikely for the infant born at a gestational age less than approximately 23 to 24 weeks and a birth weight less than 500 g (1 lb, 2 oz).</p> <p>Do not lose sight of the goal of this dramatic event: a live, neurologically intact infant and mother.</p> <p>The potential for reasonable outcome should be carefully considered before pushing the margins of survivability.</p> <p>Even if the fetus is unlikely to survive (gestational age of 20 to 23 weeks), the mother may benefit from emergency hysterotomy.</p>
Setting and personnel	
<p>Are appropriate equipment and supplies available?</p> <p>Is hysterotomy within the rescuer’s skill “comfort zone”?</p> <p>Are skilled neonatal or pediatric support personnel available to care for the infant, especially if it is not full term?</p> <p>Are obstetric personnel immediately available to support the mother after delivery?</p> <p>In both in-hospital and out-of-hospital settings, is there adequate staff and equipment support? In out-of-hospital settings, is bystander support available?</p>	<p>Equipment should be regularly rechecked.</p> <p>Effective communication is key.</p> <p>The whole team should have periodic drills.</p>
Differential diagnosis	
<p>Consider whether persistent arrest is because of an immediately reversible problem (e.g., excess anesthesia, reaction to analgesia, severe bronchospasm). If it is, the problem should be corrected, and there may be no need for hysterotomy.</p> <p>Consider whether persistent arrest is because of a fatal, untreatable problem (e.g., massive amniotic fluid embolism). If it is, an immediate hysterotomy may save the fetus.</p>	<p>This also raises the problem of quickly reversible issues (improperly mixed medication [e.g., lidocaine, magnesium sulfate]).</p> <p>If the cause is reversible or subacute, then timely management of problems may obviate the need for emergency hysterotomy.</p>

CPR = cardiopulmonary resuscitation.

Adapted from Advanced Life Support in Obstetrics Provider Syllabus. Leawood, Kan.: American Academy of Family Physicians; 2010.