

Medicine by the Numbers

A Collaboration of *TheNNT.com* and AFP

The NNT Group rating system:

Green: Benefits greater than harms

Yellow: Unclear benefits

Red: No benefits

Black: Harms greater than benefits

▶ Tap Water vs. Sterile Saline for Wound Irrigation

CHRISTINA CHAO, MD, and DAN RUNDE, MD

TAP WATER VS. STERILE SALINE FOR WOUND IRRIGATION

No significant difference

Benefits

Harms

None were harmed by using tap water

Details for This Review

Study Population: Adults with superficial wounds

Efficacy End Points: Prevention of wound infection

Harm End Points: None identified

Narrative: Various solutions are available for use in wound irrigation. It has been suggested that normal saline may be preferable because of its isotonicity and sterility. This review examines tap water as an alternative to normal saline for wound cleansing and prevention of subsequent infection.

A Cochrane review, including three studies of adults and two studies of children, compared the rates of acute infection from the use of these two solutions for acute wound cleansing.¹ In adults (1,328 participants), there was a nonsignificant decrease in wound infection in the tap water group (relative risk = 0.66; 95% confidence interval [CI], 0.42 to 1.04). In children (535 participants), there was a nonsignificant increase in wound infections in the tap water group (relative risk = 1.07; 95% CI, 0.43 to 2.64).

A double-blind randomized controlled trial published since this Cochrane review found similar results. In the 663 participants studied, there was no difference between those irrigated with tap water vs. normal saline, with a trend toward decreased infections in the tap water group.²

Given the lack of adverse events and the affordability of tap water, tap water should be considered as preferable to normal saline for cleansing of acute wounds.

Caveats: In the Cochrane review studies, there was no standardization for wound infection diagnosis across the pooled data. In the case of chronic wounds, a small study found a similar, nonsignificant decrease in infection in the tap water group compared with the normal saline group (number needed to treat = 9; 95% CI, 0.01 to 2.96).³

Interestingly, two studies comparing the use of tap water for wound cleansing vs. no cleansing (one enrolling patients with acute head wounds and the other enrolling postoperative hernia patients caring for their wounds at home) found no difference in infection rates.^{4,5} However, these studies were small and of moderate quality, and further investigations are needed before these findings can be generalized to a broader group.

Copyright 2015 The NNT Group. Used with permission.

This series is coordinated by Dean A. Seehusen, MD, MPH, AFP Contributing Editor, and Daniel Runde, MD, from the NNT Group.

This review is available from the NNT Group at <http://www.thennt.com/nnt/tap-water-for-wound-irrigation/>.

Author disclosure: No relevant financial affiliations.

REFERENCES

1. Fernandez R, Griffiths R. Water for wound cleansing. *Cochrane Database Syst Rev*. 2012;(2):CD003861.
2. Weiss EA, Oldham G, Lin M, Foster T, Quinn JV. Water is a safe and effective alternative to sterile normal saline for wound irrigation prior to suturing: a prospective, double-blind, randomised, controlled clinical trial. *BMJ Open*. 2013;3(1):e001504.
3. Griffiths RD, Fernandez RS, Ussia CA. Is tap water a safe alternative to normal saline for wound irrigation in the community setting? *J Wound Care*. 2001;10(10):407-411.
4. Goldberg HM, Rosenthal SA, Nemetz JC. Effect of washing closed head and neck wounds on wound healing and infection. *Am J Surg*. 1981;141(3):358-359.
5. Riederer SR, Inderbitzi R. Does a shower put postoperative healing at risk? [in German]. *Chirurg*. 1997;68(7):715-717. ■