

Corticosteroid Injections for Common Musculoskeletal Conditions

ZOË J. FOSTER, MD; TYLER T. VOSS, DO; JACQUELYNN HATCH, DO; and ADAM FRIMODIG, DO
St. Joseph Mercy Livingston Family Medicine Residency, Brighton, Michigan

Family physicians considering corticosteroid injections as part of a comprehensive treatment plan for musculoskeletal diagnoses will find few high-quality studies to assist with evidence-based decision making. Most studies of corticosteroid injections for the treatment of osteoarthritis, tendinopathy, bursitis, or neuropathy include only small numbers of patients and have inconsistent long-term follow-up. Corticosteroid injections for the treatment of adhesive capsulitis result in short-term improvements in pain and range of motion. For subacromial impingement syndrome, corticosteroid injections provide short-term pain relief and improvement in function. In medial and lateral epicondylitis, corticosteroid injections offer only short-term improvement of symptoms and have a high rate of symptom recurrence. Corticosteroid injections for carpal tunnel syndrome may help patients avoid or delay surgery. Trigger finger and de Quervain tenosynovitis may be treated effectively with corticosteroid injections. Patients with hip or knee osteoarthritis may have short-term symptom relief with corticosteroid injections. (*Am Fam Physician*. 2015;92(8):694-699. Copyright © 2015 American Academy of Family Physicians.)

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz Questions on page 674.

Author disclosure: No relevant financial affiliations.

Family physicians often treat patients with musculoskeletal conditions.¹ In the United States, 8% of ambulatory visits are for musculoskeletal and soft tissue concerns, and more than 13% of these patients have arthritis as a comorbid chronic condition.² Although many common musculoskeletal diagnoses can be successfully treated with conservative treatments such as rest, ice or heat, splinting, oral nonsteroidal anti-inflammatory drugs (NSAIDs), and physical therapy, almost two-thirds of family physicians use corticosteroid injections as part of a treatment plan.³ Treatment of musculoskeletal conditions by a primary care physician can improve patient satisfaction⁴ and help avoid costly referrals. Discussion of the diagnostic benefits, techniques, and adverse effects of corticosteroid injections is beyond the scope of this article; however, given the reported rarity of adverse effects, physicians should consider them relatively safe. This article summarizes the research on corticosteroid injections to provide family physicians with evidence-based recommendations for their use (*Table 1*).

Adhesive Capsulitis

Adhesive capsulitis describes a continuum of pain and stiffness in the shoulder, followed

by contracture of the shoulder structures, then gradual recovery of range of motion.⁵⁻⁷ It is generally self-limiting, although some patients report persistent pain and limited range of motion after three years.^{5,7,8}

Most interventions for adhesive capsulitis result in improvement. In the short term (eight to 16 weeks), intra-articular or subacromial corticosteroid injections outperform oral corticosteroids, NSAIDs, and physical therapy.^{5,8,9} However, the difference in outcomes between groups is not statistically significant after one year.^{5,8,9} Surgery (manipulation with arthroscopic arthrolysis) is superior in terms of range of motion and function by six weeks, although patients who receive intra-articular corticosteroid injections have similar gains by 12 weeks.⁷

Physicians should recommend conservative treatment initially, adding oral or injected corticosteroids if there is not adequate improvement in pain, range of motion, or function. Before referring a patient for manipulation under anesthesia, physicians should consider a subacromial corticosteroid injection, because it poses less risk.

Subacromial Impingement Syndrome

Subacromial impingement syndrome includes rotator cuff syndrome, tendinitis,

Table 1. Corticosteroid Injections for Common Musculoskeletal Conditions: Relative Success and Duration of Pain Relief

Condition	Short-term relief	Long-term relief	Strength of evidence	Success of therapy
Adhesive capsulitis	++ to +++	++ to +++	++	Faster pain relief and improved range of motion and function in short term; equivalent to other options in long term
Carpal tunnel syndrome	+++	+	+	Short-term improvements in pain and function, with up to one-half of patients having a recurrence in the long term; reduces need for surgery in some patients
De Quervain tenosynovitis	++	NA	+	Short-term improvements in pain and function compared with placebo
Greater trochanteric bursitis	+++	+ to ++	++	Short-term pain relief, although no better than usual care in the long term
Hip osteoarthritis	++ to +++	+ to ++	++	Injections challenging to administer in office settings
Knee osteoarthritis	++ to +++	+	++	One to two weeks of pain relief after injection
Lateral epicondylitis	++ to +++	+	+	Short-term pain relief with significant risk of symptom rebound in long term
Medial epicondylitis	++	+	+	Short-term pain relief, although no long-term difference compared with placebo
Morton neuroma	NA	NA	+	Success of therapy is unknown because of poor-quality studies
Subacromial impingement syndrome	+	+	++ to +++	Short-term pain relief equal to systemic corticosteroids; similar long-term outcomes as other treatments
Trigger finger	++ to +++	++	+	Cure rates range from 54% to 86%
Wrist and hand osteoarthritis	NA	NA	+	No strong recommendation from the American College of Rheumatology

NA = not available; + = weak or conflicting evidence or poor success; ++ = fair evidence or success; +++ = good evidence or success.

and bursitis of the shoulder.¹⁰ It results from a combination of intrinsic and extrinsic factors that injure the bursa and rotator cuff tendons in the subacromial space.¹¹ Subacromial impingement syndrome is the most common diagnosis in patients with shoulder pain.^{12,13}

Corticosteroid injections provide a maximum of four weeks of pain relief.¹⁴ Intra-articular injections are no better than placebo, and subacromial injections provide only minimal pain relief.¹⁴ Ultrasound-guided injections have no benefit compared with landmark-guided injections.¹⁵ There is no difference in pain or disability at four weeks with systemic vs. subacromial injections.¹⁶

Corticosteroid injections for subacromial impingement syndrome have similar to worse outcomes compared with conservative treatments. A combination of physical therapy and corticosteroid injections provides better pain relief at six weeks compared with physical therapy alone, but by 12 and 24 weeks there is no difference between treatments.¹² Subacromial injections of triamcinolone are inferior to ketorolac for short-term improvement in forward flexion strength and patient

satisfaction.¹⁷ In an industry-sponsored study, application of a heated lidocaine/tetracaine patch (Synera) provided similar improvements in pain and range of motion as subacromial triamcinolone injections by six weeks.¹⁸ A Cochrane review found no difference in outcomes at four weeks between corticosteroid injections, ultrasonography, acupuncture, and oral NSAIDs.¹⁴

Physicians should recommend conservative therapies initially for treatment of subacromial impingement syndrome. The patient's preferences and duration of symptoms should also be considered, because corticosteroid injections are more effective for those with symptoms of less than 12 weeks' duration.¹⁹

Lateral Epicondylitis

Lateral epicondylitis (tennis elbow) is an overuse injury affecting the origin of the wrist extensor muscles on the lateral epicondyle.²⁰ Most cases resolve within six months to two years.²¹

In patients with symptoms of up to six weeks' duration, corticosteroid injection provides pain relief for

Corticosteroid Injections

four to six weeks.²⁰⁻²⁷ One trial showed corticosteroids to be superior to local anesthetic in terms of pain and disability at three weeks, but by 12 weeks the corticosteroid group had a 35% recurrence rate compared with no recurrences in the local anesthetic group.²⁷

Long-term outcomes after corticosteroid injection indicate no benefit or symptom worsening.^{21,26} The recurrence rate one year after a single corticosteroid injection is more than 50%, compared with a 12% recurrence rate in patients receiving a placebo injection.^{23,26} Physical therapy also has poor long-term results, with a one-year recurrence rate of 8% to 29%.^{23,26} Treatment with a muscle energy technique improved pain-free grip strength and pain at 52 weeks compared with a single injection of corticosteroid.²⁸

Corticosteroid injections can be part of a treatment plan for lateral epicondylitis, although patients should be warned that there is a high recurrence rate. Patients should be counseled that current evidence indicates that long-term outcomes are often the same, regardless of initial treatment.

Medial Epicondylitis

Medial epicondylitis (golfer's elbow) is pain over the medial epicondyle.²⁹ Corticosteroid injection can provide short-term improvements in pain, although there is no long-term difference in outcomes between corticosteroid injections and placebo at six weeks and one year.^{29,30} Platelet-rich plasma injection provides greater improvements in pain and function at 52 weeks compared with corticosteroids, although it is unclear whether these outcomes are clinically meaningful to patients.³¹ Given the lack of evidence, physicians should look to studies of lateral epicondylitis for guidance.

Carpal Tunnel Syndrome

Carpal tunnel syndrome is caused by entrapment of the median nerve under the transverse carpal ligament at the wrist.³² Genetic predisposition and obesity are risk factors.³³ Younger patients with shorter symptom duration are more likely to have spontaneous improvement compared with those who have bilateral symptoms and a positive Phalen test.³⁴

Corticosteroid injections are beneficial in improving short-term outcomes.³⁵⁻³⁸ Triamcinolone injections result in improved symptom scores and functional status at one week compared with placebo injection.³⁵ Methylprednisolone injection reduces symptom severity at 10 weeks compared with placebo.³⁶ At two weeks, systemic corticosteroids and local injections provide similar improvements in symptoms; however, patients

who receive injections have significantly better symptom scores by eight and 12 weeks.³⁹ Local corticosteroid injection and oral NSAIDs with a neutral wrist splint result in similar short-term improvements in symptom severity.⁴⁰ Two corticosteroid injections provide no additional improvement compared with only a single injection.⁴¹

Long-term outcomes are mixed in patients with carpal tunnel syndrome who are treated with corticosteroid injections; up to one-half have symptom recurrence.³⁵ Ultrasonography may improve the duration of effect and other parameters, but at significant cost.⁴² The need for surgery at one year is reduced with corticosteroid injection, but a significant number of patients (75% in one study) eventually undergo surgical treatment.³⁶

Physicians can tailor their treatment recommendations to the patient's preferences. Corticosteroid injection is an option as long as patients are aware that it may not prevent recurrence or the need for surgery.

De Quervain Tenosynovitis

De Quervain tenosynovitis is tendinosis of the abductor pollicis longus and extensor pollicis brevis in the first dorsal compartment of the wrist.⁴³ Research on this condition is limited by small patient numbers, differences in the type of injectable corticosteroid used, and poor long-term follow-up.⁴⁴

A 2009 Cochrane review supports the use of corticosteroid injections over splinting in patients with de Quervain tenosynovitis,⁴⁴ although the study on which the recommendation is based included only pregnant and lactating women.⁴⁵ A more recent study shows improvement in pain and function at one week after corticosteroid injection compared with placebo.⁴⁶ Corticosteroid injection alone results in improved outcomes compared with a combination of injection and splinting, whereas treatment with rest alone or NSAIDs alone does not improve symptoms.⁴³

Physicians should consider corticosteroid injections for patients with de Quervain tenosynovitis. Referral for surgery may be considered if symptoms do not resolve after injections,⁴⁶ because anatomic variants in the first extensor compartment may explain treatment failures.⁴³

Trigger Finger

Trigger finger is the triggering or locking of a finger caused by histologic changes in the A1 pulley that prevent smooth motion of the flexor digitorum profundus and superficialis tendons.⁴⁷ Rheumatoid arthritis, diabetes mellitus, carpal tunnel syndrome, and de Quervain tenosynovitis are associated with trigger finger.⁴⁸

Cure rates after corticosteroid injection are generally moderate, ranging from 54% to 86%.^{48,49} Compared with

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	References
Corticosteroid injections in the shoulder have only short-term benefits in adhesive capsulitis and subacromial impingement syndrome.	B	5, 7-9, 12, 14, 19
Corticosteroid injections for lateral and medial epicondylitis lead to short-term improvement but have a high rate of recurrence and are no better than other options in the long term.	B	21, 26-30
Corticosteroid injections can be considered for patients with carpal tunnel syndrome who wish to avoid or delay surgical treatment.	B	35, 36, 40
Corticosteroid injections for de Quervain tenosynovitis and trigger finger are effective early in therapy.	B	44, 46, 48, 50
Corticosteroid injections provide short-term relief from symptoms of knee and hip osteoarthritis in patients who wish to delay surgery.	B	57, 60-63

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.

placebo injections, corticosteroids result in less triggering and improved pain at one week, and these effects persist through one year.⁵⁰ In one study, corticosteroid injections resulted in less topical and joint pain at one week and one month compared with percutaneous and open releases; however, patients who underwent surgery had better outcomes at six months.⁴⁸ Symptom relapse occurred in 12% of patients who received one injection, 18% of those who received a second injection, and none who underwent surgery.⁴⁸ However, a Cochrane review notes that studies of corticosteroid injections for trigger finger are limited by sample size and short duration of follow-up.⁴⁷

Physicians should offer a corticosteroid injection for patients with trigger finger who are seeking a nonsurgical treatment option.

Wrist and Hand Osteoarthritis

The American College of Rheumatology makes “no strong recommendation” for intra-articular corticosteroid injection in patients with hand osteoarthritis.⁵¹ A Cochrane review of intra-articular corticosteroid injections for wrist or hand rheumatoid arthritis found only one study with moderate bias that recommended against splinting or resting the wrist after intra-articular corticosteroid injection.⁵² Physicians are best advised to refer patients who desire such an intervention to a subspecialist.

Greater Trochanteric Bursitis

Greater trochanteric bursitis is in the differential diagnosis for pain in the lateral hip. This condition has recently been referred to as greater trochanteric pain syndrome, reflecting the results of radiologic and histopathologic studies that indicate no inflammation of the subgluteus maximus bursa.⁵³

Compared with usual care (analgesics and physical therapy), corticosteroid injections provide greater pain relief at rest and with activity at six weeks and three months, but no differences in pain or function at six months and one

year.⁵⁴ Physicians may consider corticosteroid injections for patients with greater trochanteric pain who desire more prompt resolution of their symptoms.

Hip Osteoarthritis

Hip osteoarthritis affects more than one-third of patients older than 85 years.⁵⁵ Treatment with intra-articular corticosteroid injections in the outpatient setting is challenging given the need for radiographic guidance.⁵⁶ A small double-blind study comparing corticosteroid and placebo injections using an ultrasound-guided anterolateral approach showed improvement in pain on walking at two and four weeks for the corticosteroid group, although that improvement disappeared by three months.⁵⁷ Fluoroscopically guided corticosteroid injections provide improvement in pain, stiffness, and physical function at two months compared with placebo injections.⁵⁵ Most primary care physicians should consider orthopedic referral for patients who desire injection treatment for hip osteoarthritis.

Knee Osteoarthritis

Knee osteoarthritis is a common cause of knee pain in older adults. It is estimated that one in six persons older than 55 years presents with knee pain in a given year.⁵⁸ More than one-half of patients older than 85 years likely have symptomatic knee osteoarthritis.⁵⁹

Guidelines from the American College of Rheumatology conditionally recommend the use of intra-articular corticosteroid injections for treatment of knee osteoarthritis.⁵¹ The duration of pain relief is one to two weeks in most trials, with a few showing improvements lasting three to four weeks.⁶⁰⁻⁶³ Research uniformly supports the safety of intra-articular corticosteroid injections for treatment of knee osteoarthritis; however, these studies are limited by lack of histologic data and poor long-term follow-up.⁶⁴ A Cochrane review found weak evidence for the use of corticosteroid injections for the treatment of knee rheumatoid arthritis.⁵²

Corticosteroid Injections

Physicians may consider intra-articular corticosteroid injections before orthopedic referral for surgical evaluation in patients with knee osteoarthritis.

Morton Neuroma

Morton neuroma refers to pain between the metatarsal heads from the plantar digital nerve, most commonly affecting the third web space.⁶⁵

A Cochrane review notes that although corticosteroid injection is considered a standard treatment for Morton neuroma, few studies have investigated its safety and effectiveness, and those that are published are of poor quality.⁶⁵ Physicians may consider corticosteroid injection for these patients, but should be advised there is no good-quality evidence to support it.

Data Sources: PubMed, SPORTDiscus (EBSCO), Cochrane, and the AHRQ databases were searched for studies published within the past 10 years. Additional searches were performed using Essential Evidence Plus. Key words were corticosteroid injections and lateral epicondylitis/tennis elbow, carpal tunnel syndrome/median neuropathy, deQuervain's tenosynovitis/thumb tendinitis, rotator cuff tendinitis/shoulder bursitis/shoulder pain, greater trochanteric bursitis/lateral hip pain, and knee pain/knee osteoarthritis. Search dates: March 2014 through June 2015.

The authors thank Jillah Biza, Riecker Memorial Library at St. Joseph Mercy Hospital, Ann Arbor, Mich., for her assistance with literature searches and obtaining articles.

The Authors

ZOË J. FOSTER, MD, is a faculty physician at St. Joseph Mercy Livingston Family Medicine Residency, Brighton, Mich., and for the University of Michigan Primary Care Sports Medicine Fellowship Program, Ann Arbor. She is also a team physician for the University of Michigan; Eastern Michigan University, Ypsilanti; and Pinckney (Mich.) Community High School.

TYLER T. VOSS, DO, is a primary care sports medicine fellow at St. John Providence Health System, Brighton. At the time the article was written, he was a third-year resident at St. Joseph Mercy Livingston Family Medicine Residency.

JACQUELYNN HATCH, DO, is a third-year resident at St. Joseph Mercy Livingston Family Medicine Residency.

ADAM FRIMODIG, DO, is a third-year resident at St. Joseph Mercy Livingston Family Medicine Residency.

Address correspondence to Zoë J. Foster, MD, St. Joseph Mercy Hospital, 7575 Grand River, Ste. 210, Brighton, MI 48114 (e-mail: zoe_foster@ihacares.com). Reprints are not available from the authors.

REFERENCES

1. MacKay C, Canizares M, Davis AM, et al. Health care utilization for musculoskeletal disorders. *Arthritis Care Res (Hoboken)*. 2010;62(2):161-169.
2. Cherry DK, Hing E, Woodwell DA, et al. National ambulatory medical health care survey: 2006 summary. *Natl Health Stat Report*. 2008;(3):1-39.
3. American Academy of Family Physicians. Clinical procedures performed by physicians at their practice. <http://www.aafp.org/about/the-aafp/family-medicine-facts/table-12.html>. Accessed January 29, 2015.

4. Bhagra A, Syed H, Reed DA, et al. Efficacy of musculoskeletal injections by primary care providers in the office: a retrospective cohort study. *Int J Gen Med*. 2013;6:237-243.
5. Blanchard V, Barr S, Cerisola FL. The effectiveness of corticosteroid injections compared with physiotherapeutic interventions for adhesive capsulitis: a systematic review. *Physiotherapy*. 2010;96(2):95-107.
6. Griesser MJ, Harris JD, Campbell JE, et al. Adhesive capsulitis of the shoulder: a systematic review of the effectiveness of intra-articular corticosteroid injections. *J Bone Joint Surg Am*. 2011;93(18):1727-1733.
7. De Carli A, Vadalà A, Perugia D, et al. Shoulder adhesive capsulitis: manipulation and arthroscopic arthrolysis or intra-articular steroid injections? *Int Orthop*. 2012;36(1):101-106.
8. Lorbach O, Anagnostakos K, Scherf C, et al. Nonoperative management of adhesive capsulitis of the shoulder: oral cortisone application versus intra-articular cortisone injections. *J Shoulder Elbow Surg*. 2010;19(2):172-179.
9. Shin SJ, Lee SY. Efficacies of corticosteroid injection at different sites of the shoulder for the treatment of adhesive capsulitis. *J Shoulder Elbow Surg*. 2013;22(4):521-527.
10. Huisstede BM, Miedema HS, Verhagen AP, et al. Multidisciplinary consensus on the terminology and classification of complaints of the arm, neck and/or shoulder. *Occup Environ Med*. 2007;64(5):313-319.
11. Alqunae M, Galvin R, Fahey T. Diagnostic accuracy of clinical tests for subacromial impingement syndrome: a systematic review and meta-analysis. *Arch Phys Med Rehabil*. 2012;93(2):229-236.
12. Crawshaw DP, Helliwell PS, Hensor EM, et al. Exercise therapy after corticosteroid injection for moderate to severe shoulder pain: large pragmatic randomised trial. *BMJ*. 2010;340:c3037.
13. van der Windt DA, Koes BW, de Jong BA, et al. Shoulder disorders in general practice: incidence, patient characteristics, and management. *Ann Rheum Dis*. 1995;54(12):959-964.
14. Buchbinder R, Green S, Youd JM. Corticosteroid injections for shoulder pain. *Cochrane Database Syst Rev*. 2003;(1):CD004016.
15. Bloom JE, Rischin A, Johnston RV, et al. Image-guided versus blind glucocorticoid injection for shoulder pain. *Cochrane Database Syst Rev*. 2012;(8):CD009147.
16. Ekeberg OM, Bautz-Holter E, Tveitå EK, et al. Subacromial ultrasound guided or systemic steroid injection for rotator cuff disease: randomised double blind study. *BMJ*. 2009;338:a3112.
17. Min KS, St Pierre P, Ryan PM, et al. A double-blind randomized controlled trial comparing the effects of subacromial injection with corticosteroid versus NSAID in patients with shoulder impingement syndrome. *J Shoulder Elbow Surg*. 2013;22(5):595-601.
18. Radnovich R, Trudeau J, Gammaitoni AR. A randomized clinical study of the heated lidocaine/tetracaine patch versus subacromial corticosteroid injection for the treatment of pain associated with shoulder impingement syndrome. *J Pain Res*. 2014;7:727-735.
19. Gaujoux-Viala C, Dougados M, Gossec L. Efficacy and safety of steroid injections for shoulder and elbow tendonitis: a meta-analysis of randomized controlled trials. *Ann Rheum Dis*. 2009;68(12):1843-1849.
20. Smidt N, van der Windt DA, Assendelft WJ, et al. Corticosteroid injections, physiotherapy, or a wait-and-see policy for lateral epicondylitis: a randomised controlled trial. *Lancet*. 2002;359(9307):657-662.
21. Krogh TP, Bartels EM, Ellingsen T, et al. Comparative effectiveness of injection therapies in lateral epicondylitis: a systematic review and network meta-analysis of randomized controlled trials. *Am J Sports Med*. 2013;41(6):1435-1446.
22. Crowther MA, Bannister GC, Huma H, et al. A prospective, randomised study to compare extracorporeal shock-wave therapy and injection of steroid for the treatment of tennis elbow. *J Bone Joint Surg Br*. 2002;84(5):678-679.
23. Bisset L, Beller E, Jull G, et al. Mobilisation with movement and exercise, corticosteroid injection, or wait and see for tennis elbow: randomised trial. *BMJ*. 2006;333(7575):939.

24. Hay EM, Paterson SM, Lewis M, et al. Pragmatic randomised controlled trial of local corticosteroid injection and naproxen for treatment of lateral epicondylitis of elbow in primary care. *BMJ*. 1999;319(7215):964-968.
25. Assendelft WJ, Hay EM, Adsheed R, et al. Corticosteroid injections for lateral epicondylitis: a systematic overview. *Br J Gen Pract*. 1996;46(405):209-216.
26. Coombes BK, Bisset L, Brooks P, et al. Effect of corticosteroid injection, physiotherapy, or both on clinical outcomes in patients with unilateral lateral epicondylalgia: a randomized controlled trial. *JAMA*. 2013;309(5):461-469.
27. Mardani-Kivi M, Karimi-Mobarakeh M, Karimi A, et al. The effects of corticosteroid injection versus local anesthetic injection in the treatment of lateral epicondylitis: a randomized single-blinded clinical trial. *Arch Orthop Trauma Surg*. 2013;133(6):757-763.
28. Küçükşen S, Yilmaz H, Sallı A, et al. Muscle energy technique versus corticosteroid injection for management of chronic lateral epicondylitis: randomized controlled trial with 1-year follow-up. *Arch Phys Med Rehabil*. 2013;94(11):2068-2074.
29. Stahl S, Kaufman T. The efficacy of an injection of steroids for medial epicondylitis. A prospective study of sixty elbows. *J Bone Joint Surg Am*. 1997;79(11):1648-1652.
30. Coombes BK, Bisset L, Vicenzino B. Efficacy and safety of corticosteroid injections and other injections for management of tendinopathy: a systematic review of randomised controlled trials. *Lancet*. 2010;376(9754):1751-1767.
31. Peerbooms JC, Sluimer J, Bruijn DJ, et al. Positive effect of an autologous platelet concentrate in lateral epicondylitis in a double-blind randomized controlled trial: platelet-rich plasma versus corticosteroid injection with a 1-year follow-up. *Am J Sports Med*. 2010;38(2):255-262.
32. Katz JN, Simmons BP. Clinical practice. Carpal tunnel syndrome. *N Engl J Med*. 2002;346(23):1807-1812.
33. Bland JD. Carpal tunnel syndrome. *BMJ*. 2007;335(7615):343-346.
34. Padua L, Padua R, Aprile I, et al. Carpal tunnel syndrome. Multiperspective follow-up of untreated carpal tunnel syndrome: a multicenter study. *Neurology*. 2001;56(11):1459-1466.
35. Peters-Veluthamaningal C, Winters JC, Groenier KH, et al. Randomised controlled trial of local corticosteroid injections for carpal tunnel syndrome in general practice. *BMC Fam Pract*. 2010;11:54.
36. Atroshi I, Flondell M, Hofer M, et al. Methylprednisolone injections for the carpal tunnel syndrome: a randomized, placebo-controlled trial. *Ann Intern Med*. 2013;159(5):309-317.
37. Ly-Pen D, Andréu JL, de Blas G, et al. Surgical decompression versus local steroid injection in carpal tunnel syndrome: a one-year, prospective, randomized, open, controlled clinical trial. *Arthritis Rheum*. 2005;52(2):612-619.
38. Marshall S, Tardif G, Ashworth N. Local corticosteroid injection for carpal tunnel syndrome. *Cochrane Database Syst Rev*. 2007;(2):CD001554.
39. Wong SM, Hui AC, Tang A, et al. Local vs systemic corticosteroids in the treatment of carpal tunnel syndrome. *Neurology*. 2001;56(11):1565-1567.
40. Celiker R, Arslan S, Inanici F. Corticosteroid injection vs. nonsteroidal antiinflammatory drug and splinting in carpal tunnel syndrome. *Am J Phys Med Rehabil*. 2002;81(3):182-186.
41. Wong SM, Hui AC, Lo SK, et al. Single vs. two steroid injections for carpal tunnel syndrome: a randomised clinical trial. *Int J Clin Pract*. 2005;59(12):1417-1421.
42. Makhlof T, Emil NS, Sibbitt WL Jr, et al. Outcomes and cost-effectiveness of carpal tunnel injections using sonographic needle guidance. *Clin Rheumatol*. 2014;33(6):849-858.
43. Richie CA III, Briner WW Jr. Corticosteroid injection for treatment of de Quervain's tenosynovitis: a pooled quantitative literature evaluation. *J Am Board Fam Pract*. 2003;16(2):102-106.
44. Peters-Veluthamaningal C, van der Windt DA, Winters JC, et al. Corticosteroid injection for de Quervain's tenosynovitis. *Cochrane Database Syst Rev*. 2009;(3):CD005616.
45. Avci S, Yilmaz C, Sayli U. Comparison of nonsurgical treatment measures for de Quervain's disease of pregnancy and lactation. *J Hand Surg Am*. 2002;27(2):322-324.
46. Peters-Veluthamaningal C, Winters JC, Groenier KH, et al. Randomised controlled trial of local corticosteroid injections for de Quervain's tenosynovitis in general practice. *BMC Musculoskelet Disord*. 2009;10:131.
47. Peters-Veluthamaningal C, van der Windt DA, Winters JC, et al. Corticosteroid injection for trigger finger in adults. *Cochrane Database Syst Rev*. 2009;(1):CD005617.
48. Sato ES, Gomes Dos Santos JB, et al. Treatment of trigger finger: randomized clinical trial comparing the methods of corticosteroid injection, percutaneous release and open surgery. *Rheumatology (Oxford)*. 2012;51(1):93-99.
49. van Middelkoop M, Huisstede BM, Glerum S, et al. Effectiveness of interventions of specific complaints of the arm, neck, or shoulder (CANS): musculoskeletal disorders of the hand. *Clin J Pain*. 2009;25(6):537-552.
50. Peters-Veluthamaningal C, Winters JC, Groenier KH, et al. Corticosteroid injections effective for trigger finger in adults in general practice: a double-blinded randomised placebo controlled trial. *Ann Rheum Dis*. 2008;67(9):1262-1266.
51. Hochberg MC, Altman RD, April KT, et al. American College of Rheumatology 2012 recommendations for the use of nonpharmacologic and pharmacologic therapies in osteoarthritis of the hand, hip, and knee. *Arthritis Care Res (Hoboken)*. 2012;64(4):465-474.
52. Wallen M, Gillies D. Intra-articular steroids and splints/rest for children with juvenile idiopathic arthritis and adults with rheumatoid arthritis. *Cochrane Database Syst Rev*. 2006;(1):CD002824.
53. Silva F, Adams T, Feinstein J, et al. Trochanteric bursitis: refuting the myth of inflammation. *J Clin Rheumatol*. 2008;14(2):82-86.
54. Brinks A, van Rijn RM, Willemsen SP, et al. Corticosteroid injections for greater trochanteric pain syndrome: a randomized controlled trial in primary care. *Ann Fam Med*. 2011;9(3):226-234.
55. Lambert RG, Hutchings EJ, Grace MG, et al. Steroid injection for osteoarthritis of the hip: a randomized, double-blind, placebo-controlled trial. *Arthritis Rheum*. 2007;56(7):2278-2287.
56. Dorleijn DM, Luijsterburg PA, Reijman M, et al. Effectiveness of intramuscular corticosteroid injection versus placebo injection in patients with hip osteoarthritis: design of a randomized double-blinded controlled trial. *BMC Musculoskelet Disord*. 2011;12:280.
57. Qvistgaard E, Christensen R, Torp-Pedersen S, et al. Intra-articular treatment of hip osteoarthritis: a randomized trial of hyaluronic acid, corticosteroid, and isotonic saline. *Osteoarthritis Cartilage*. 2006;14(2):163-170.
58. Peat G, McCarney R, Croft P. Knee pain and osteoarthritis in older adults: a review of community burden and current use of primary health care. *Ann Rheum Dis*. 2001;60(2):91-97.
59. Murphy L, Schwartz TA, Helmick CG, et al. Lifetime risk of symptomatic knee osteoarthritis. *Arthritis Rheum*. 2008;59(9):1207-1213.
60. Cheng OT, Souzdalnitski D, Vrooman B, et al. Evidence-based knee injections for the management of arthritis. *Pain Med*. 2012;13(6):740-753.
61. Arroll B, Goodyear-Smith F. Corticosteroid injections for osteoarthritis of the knee: meta-analysis. *BMJ*. 2004;328(7444):869.
62. Hepper CT, Halvorson JJ, Duncan ST, et al. The efficacy and duration of intra-articular corticosteroid injection for knee osteoarthritis: a systematic review of level I studies. *J Am Acad Orthop Surg*. 2009;17(10):638-646.
63. Bellamy N, Campbell J, Robinson V, et al. Intraarticular corticosteroid for treatment of osteoarthritis of the knee. *Cochrane Database Syst Rev*. 2006;(2):CD005328.
64. Raynauld JP, Buckland-Wright C, Ward R, et al. Safety and efficacy of long-term intraarticular steroid injections in osteoarthritis of the knee: a randomized, double-blind, placebo-controlled trial. [published correction appears in *Arthritis Rheum*. 2003;48(11):3300]. *Arthritis Rheum*. 2003;48(2):370-377.
65. Thomson CE, Gibson JN, Martin D. Interventions for the treatment of Morton's neuroma. *Cochrane Database Syst Rev*. 2004;(3):CD003118.