

Common Questions About Pressure Ulcers

JAQUELINE G.M. RAETZ, MD, and KEREN H. WICK, PhD, *University of Washington, Seattle, Washington*

Patients with limited mobility due to physical or cognitive impairment are at risk of pressure ulcers. Primary care physicians should examine at-risk patients because pressure ulcers are often missed in inpatient, outpatient, and long-term care settings. High-risk patients should use advanced static support surfaces to prevent pressure ulcers and air-fluidized beds to treat pressure ulcers. Physicians should document the size and clinical features of ulcers. Cleansing should be done with saline or tap water, while avoiding caustic agents, such as hydrogen peroxide. Dressings should promote a moist, but not wet, wound healing environment. The presence of infection is determined through clinical judgment; if uncertain, a tissue biopsy should be performed. New or worsening pain may indicate infection of a pressure ulcer. When treating patients with pressure ulcers, it is important to keep in mind the patient's psychological, behavioral, and cognitive status. The patient's social, financial, and caregiver resources, as well as goals and long-term prognosis, should also be considered in the treatment plan. (*Am Fam Physician*. 2015;92(10):888-894. Copyright © 2015 American Academy of Family Physicians.)

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz Questions on page 869.

Author disclosure: No relevant financial affiliations.

Pressure ulcers are injuries of the skin or underlying tissue that occur as a result of pressure alone or in combination with shearing forces. The ulcers typically occur over bony prominences.¹ During pressure ulcer development, external pressure exceeds capillary blood flow pressure, leading to ischemia and tissue injury.² Pressure ulcers are common, affecting up to 3 million Americans³ and costing about \$11 billion annually in the United States.⁴ The most important risks for pressure ulcer development are impaired mobility, decreased perfusion, and edema or a stage I pressure ulcer.⁵ Additional risk factors include poor general health status, low nutritional status, and skin moisture from conditions such as incontinence.⁵ Advanced age and cognitive impairment are also risk factors.⁵ Multifactorial interventions to reduce pressure ulcers are among the top-10 strategies strongly encouraged by the Agency for Healthcare Research and Quality for adoption in all health care systems.⁶ When treating patients with pressure ulcers, it is important to keep in mind the patient's psychological, behavioral, and cognitive status. The patient's social, financial, and caregiver resources, as well as goals and long-term prognosis, should also be considered when developing a treatment plan.

What Are the Most Effective Ways to Prevent Pressure Ulcers in Hospitalized or Immobile Patients?

For patients at risk of pressure ulcers, it is important to examine the skin and initiate a multicomponent intervention.⁷ Advanced static support surfaces should be used for high-risk patients rather than standard hospital mattresses.^{3,8,9}

EVIDENCE SUMMARY

Pressure ulcers may develop in as little as four to six hours.¹⁰ Primary care physicians should examine at-risk patients periodically because ulcers are often missed in inpatient, outpatient, and long-term care settings. The most common sites of pressure ulcers are indicated in *Figure 1*. It is important to identify patients at high risk, initiate preventive measures, and monitor for ulcer development. Moderate-quality evidence from a systematic review of 26 studies supports a multicomponent intervention to prevent pressure ulcers. This usually includes an assessment, use of a specialized support surface, repositioning, mobilization, avoiding friction, and management of nutrition and moisture. Other features include an on-site clinician to lead the program, a multidisciplinary team, audit and feedback, ongoing education, simplification, and standardization.⁷ Multidisciplinary care is crucial, including primary care physi-

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	References
Multicomponent interventions should be used to prevent pressure ulcers in high-risk patients.	B	7
Advanced static support surfaces, rather than standardized hospital mattresses, should be used to prevent pressure ulcers in high-risk patients.	A	3, 8, 9
Hydrocolloid or foam dressings should be used for the treatment of pressure ulcers.	B	4, 22
Cleansing pressure ulcers with caustic agents, such as povidone-iodine (Betadine) or Dakin's solutions, should be avoided.	C	23, 24
Air-fluidized beds should be used for the treatment of pressure ulcers.	A	4
New or worsening pain may indicate infection of a pressure ulcer.	C	32

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.

cians, nurses (including those specializing in wound care), social workers, podiatrists, surgeons, dietitians, physical and occupational therapists, and infectious disease consultants as needed.

The Braden, Waterlow, and Norton scales are commonly used for pressure ulcer risk assessment. These are five- to six-item scales that assess risk factors such as physical and mental condition, mobility, moisture, continence, and nutrition. However, they are weak predictors of the risk of ulcers, and may be no better at reducing the risk of formation than a nurse's or physician's clinical judgment.^{3,9,11} The 2015 clinical practice guideline from the American College of Physicians suggests that risk scales may be most useful to clinicians with limited experience in caring for patients who develop pressure ulcers.⁹

Adequate evidence supports the use of advanced static support surfaces rather than standard hospital mattresses to prevent pressure ulcers in high-risk patients.^{3,8,9} These support surfaces include foam, water, gel, and air mattresses or mattress overlays. Other commonly used measures include repositioning, nutritional supplementation, creams, dressings, and pads placed over bony prominences. There is inadequate evidence on the effectiveness of these measures, although all have a low risk of harm.^{3,9,12,13} There is also no evidence to suggest an optimal interval at which to reposition patients,^{14,15} although every two hours is recommended based on expert opinion.¹⁶ It is reasonable to consider consultation with a dietitian and use of skin moisturizers to prevent pressure ulcers.¹⁷ Even with optimal care, pressure ulcers may still develop in certain patients.

How Should Pressure Ulcers Be Assessed?

Clinicians should take a medical and pressure-ulcer history, including assessment of pain. They should examine and measure the length, width, and depth of ulcers. The presence of exudate, necrotic tissue, eschar, slough, undermining, and tunneling or sinus tracts should be documented. The presence of healing tissue (pink granulation tissue or epithelialization) should be noted. The wound should be staged using the National Pressure Ulcer Advisory Panel's 1 to 4 staging system.¹⁸

EVIDENCE SUMMARY

The most commonly used staging system for pressure ulcers is the National Pressure Ulcer Advisory Panel staging system (Table 1).^{18,19} This system was recently








Figure 1. Common sites where pressure ulcers develop.

Table 1. Pressure Ulcer Staging System

Stage	Description	Typical presentation
Stage 1: nonblanchable erythema of intact skin	Intact skin with nonblanchable redness of a localized area usually over a bony prominence; darkly pigmented skin may not have visible blanching, its color may differ from the surrounding area; area may be painful, firm, soft, warmer, or cooler compared with adjacent tissue; stage I may be difficult to detect in patients with dark skin tones; may indicate "at-risk" persons	
Stage 2: partial-thickness skin loss with exposed dermis	Partial-thickness loss of skin or tissue presenting as a shallow open ulcer with a red-pink wound bed, without slough; may present as an intact or open/ruptured serum-filled or serosanguineous blister; presents as a shiny or dry, shallow ulcer without slough or bruising*; this category should not be used to describe skin tears, tape burns, incontinence-associated dermatitis, maceration, or excoriation	
Stage 3: full-thickness skin loss	Full-thickness skin loss; subcutaneous fat may be visible but bone, tendon, or muscle is not exposed; slough may be present but does not obscure the depth of tissue loss; may include undermining and tunneling; depth of ulcer varies by anatomic location; bridge of the nose, ear, occiput, and malleolus do not have (adipose) subcutaneous tissue and ulcers can be shallow; in contrast, areas of significant adiposity can develop extremely deep ulcers; bone/tendon is not visible or directly palpable	
Stage 4: full-thickness skin and tissue loss	Full-thickness tissue loss with exposed bone, tendon, or muscle; slough or eschar may be present; often includes undermining and tunneling; depth of ulcer varies by anatomic location; bridge of the nose, ear, occiput, and malleolus do not have (adipose) subcutaneous tissue and ulcers can be shallow; ulcers can extend into muscle or supporting structures (e.g., fascia, tendon, joint capsule), making osteomyelitis or osteitis likely to occur; exposed bone/muscle is visible or directly palpable	
Additional categories/stages for the United States		
Unstageable/pressure injury: obscured full-thickness skin and tissue loss	Full-thickness tissue loss in which actual depth of ulcer is completely obscured by slough (yellow, tan, gray, green, or brown) or eschar (tan, brown, or black) in the wound bed; until enough slough or eschar is removed to expose the base of the wound, the true depth cannot be determined, but it will be stage 3 or 4; stable (dry, adherent, intact without erythema or fluctuance) eschar on the heels serves as a natural (biological) cover and should not be removed	
Deep-tissue pressure injury: persistent non-blanchable deep red, maroon or purple discoloration	Purple or maroon localized area of discolored intact skin or blood-filled blister caused by damage to underlying soft tissue from pressure or shear; area may be preceded by tissue that is painful, firm, mushy, boggy, warmer, or cooler compared with adjacent tissue; deep-tissue injury may be difficult to detect in patients with dark skin tones; evolution may include a thin blister over a dark wound bed; wound may further evolve and become covered by thin eschar; evolution may be rapid, exposing additional layers of tissue even with optimal treatment	

*—Bruising indicates deep-tissue injury.

Adapted with permission from National Pressure Ulcer Advisory Panel. NPUAP pressure injury stages. <http://www.npuap.org/resources/educational-and-clinical-resources/npuap-pressure-injury-stages>, and Pressure injury staging illustrations. <http://www.npuap.org/resources/educational-and-clinical-resources/pressure-injury-staging-illustrations/>. Accessed December 15, 2016.

updated, and now uses Arabic numerals and refers to "pressure injury" rather than "pressure ulcer," since not all injuries involve ulceration of the tissue. There are four stages of pressure injuries. In a stage 1 injury, the skin is intact with nonblanchable redness, or differing color from surrounding skin in darkly pigmented skin. The area may be warm, cool, soft, firm, or painful compared

with other areas. Patients with stage 1 injuries are at risk of developing additional pressure injuries. Stage 2 pressure injuries involve partial-thickness skin loss. They are shallow and have a red-pink wound bed. An intact blister is also considered a stage 2 injury. There should be no slough (dead tissue that is often a yellow-gray color and tightly adhered) or bruising in a stage 2 injury.

A stage 3 injury involves full-thickness skin loss. It may be shallow or deep, and may have undermining or tunneling. A stage 4 injury involves full-thickness tissue loss of the dermis with exposed bone, tendon, or muscle (Figure 2). Additionally, injuries may be classified as “unstageable/unclassified: full-thickness skin or tissue loss—depth unknown” if slough or eschar is present, because true depth and staging cannot be determined until it is removed. Injuries may also be classified as “suspected deep-tissue injury—depth unknown.”¹⁸

What Is the Best Way to Cleanse and Dress a Pressure Ulcer?

Pressure ulcers should be cleansed with saline or tap water.^{1,20,21} They should be debrided of slough and necrotic tissue. A hydrocolloid, foam, or other nonadherent dressing that promotes a moist wound environment should be used.^{4,22} Wet-to-dry dressings, povidone-iodine (Betadine) solution, and Dakin’s solution should be avoided.^{23,24} Pain associated with cleansing or dressing pressure ulcers should be treated appropriately.

EVIDENCE SUMMARY

There is insufficient evidence to advise using one cleansing solution or technique over another.²⁰ In general, saline is favored to cleanse pressure ulcers, but tap water may be used if saline is not available.^{1,21} Solutions that may impede granulation tissue formation, such as hydrogen peroxide, chlorine-based solutions (Dakin’s), or povidone-iodine solution, should be avoided.^{23,24} However, a time-limited course of properly diluted antiseptics may be used in an infected wound to control bacterial burden or in palliative situations with wounds

that are not expected to heal and have a foul odor caused by a heavy bacterial burden.¹

Debridement of necrotic tissue and slough is important to promote healing and decrease the risk of infection. Debridement can be autolytic (provided by the wound’s own drainage), enzymatic (collagenase is commonly available and well tolerated), mechanical (whirlpool), or surgical (sharp debridement).¹

A moist wound environment is believed to assist in healing and aid in autolytic debridement.²⁵ Low-quality evidence supports the use of a hydrocolloid or foam dressing over standard therapy.^{4,22} Hydrocolloid dressings are most useful with light to moderate wound drainage, and foam dressings are available in a range of thicknesses for drier wounds to those with heavier exudate.²⁶ If these are not available, it is important that any dressing promote moist (but not wet) wound healing and be nonadhesive. Wet-to-dry dressings should be avoided because the debridement from removal of a fully dry dressing may impede healing and can cause significant pain.¹

Other dressing options include saline and gauze (wet to moist), transparent films (which cover and create a moist environment), hydrogels (which provide moisture), and alginates (which absorb excess moisture).²⁶ Table 2 lists options for dressings based on the type of ulcer.^{1,22,26,27}

A dry, intact eschar over the heels without erythema or fluctuance should not be removed because it serves as a natural biologic cover.¹ It is important to assess and treat pain associated with pressure ulcers, particularly during debridement and dressing.¹

Does Nutritional Supplementation Improve Prognosis?

Protein supplementation may improve outcomes in the treatment of pressure ulcers.^{4,22} There is no evidence to support the use of vitamin C or zinc to treat pressure ulcers in patients who are not deficient of these nutrients.^{4,22}

EVIDENCE SUMMARY

A 2013 systematic review and a 2015 guideline by the American College of Physicians cite moderate-quality evidence supporting protein supplementation to treat pressure ulcers.^{4,22} However, a 2014 Cochrane review concluded that there is no evidence that nutritional interventions, including protein, provide benefit in the prevention or treatment of pressure ulcers.²⁸ Although commonly prescribed, only one good-quality study has evaluated the effect of vitamin C supplementation, and it revealed no change in the rate of wound healing. Insufficient evidence was found on the effect of zinc supplementation.^{4,22}



Figure 2. Stage 4 sacral pressure ulcer. Note yellow slough as well as exposed muscle.

Pressure Ulcers

Table 2. Treatment Options for Pressure Ulcer Dressings

Ulcer type*†	Best evidence-based recommendation	Other options or wound filler	Wraps/covers
Shallow and dry	Thin hydrocolloid Thin polyurethane foam	Transparent films Hydrogel‡	Nonadherent gauze
Shallow and wet	Hydrocolloid with or without absorbent paste or powder Foam	Alginate	Nonadherent gauze or absorbent contact layer
Deep and dry	Hydrocolloid (over filler dressing) Foam	Fill with damp gauze, copolymer starch, or hydrogel	Transparent thin film, polyurethane foam, or nonadherent gauze
Deep and wet	Foam	Fill with alginate, single gauze strip/roll, foam, or other absorbent dressing	Transparent thin film or polyurethane foam
Fragile or friable periwound skin	Consider use of silicone dressing	—	—

NOTE: The objective is to maintain a moist environment, absorb excess exudate, facilitate autolysis, fill cavities, and protect the wound from infection. Avoid using wet-to-dry gauze dressings.

*—Dressings are not usually needed for stage I pressure ulcers.

†—In general, shallow ulcers are defined as stage I or II and deep ulcers as stage III or IV.

‡—Not shown to be better than other standard dressings, per Cochrane review.²⁷

Information from references 1, 22, 26, and 27.

Which Other Interventions May Help When Treating Pressure Ulcers?

Relieving the source of pressure, repositioning the patient, and using air-fluidized beds are recommended to treat pressure ulcers.^{4,22} Moderate-strength evidence supports the use of radiant heat dressings and electrical stimulation.^{4,22} Alternating-pressure surfaces, platelet-derived growth factor, and light therapy may also help.^{4,22} Surgical consultation should be considered for advanced ulcers not responding to standard therapy.

EVIDENCE SUMMARY

Treatment of pressure ulcers should alleviate the causes and risks contributing to ulcer formation. Clinicians should attempt to remove all pressure over the ulcer. This may include simple techniques such as “floating heels” (Figure 3) and frequent repositioning (every two hours). However, repositioning has not been studied in randomized controlled trials.²⁹

Three systematic reviews found moderate-strength evidence supporting the use of air-fluidized beds, radiant heat dressings (a noncontact dressing attached to a heating element that provides warmth to increase capillary blood flow and resistance to infection), and

electrical stimulation (direct electric current delivered through the wound bed using surface electrodes) to improve pressure ulcer healing.^{4,22,30} Electrical stimulation is specifically recommended in the American College of Physicians guidelines.²² With electrical stimulation, high-voltage pulse current should be used because it has been better studied, penetrates deeper, and has a lower risk of burns. Daily one-hour sessions are commonly used. Most studies used stimulus pulse



Figure 3. A simple measure to prevent or treat pressure ulcers to the heels is to offload pressure by placing a pillow under the legs.

settings of a 50-microsecond duration, approximately 100 Hz in frequency, and 100 to 150 V in intensity.³⁰ Electrical stimulation should not be used in patients with cancer because it could stimulate cancerous cells, or in patients with osteomyelitis because it may cause premature closure.³⁰

Low-strength evidence supports the use of alternating-pressure surfaces, platelet-derived growth factor, and light therapy (infrared, visible, and ultraviolet spectrum light).^{4,22} However, a Cochrane review concluded that because of the low-quality evidence, the effects of phototherapy are uncertain.³¹

If consistent with a patient's goals of care, surgical consultation is recommended based on expert opinion and usual practice for stage 3 or 4 ulcers not responding to standard therapy. There is insufficient evidence to support one surgical technique over another, and dehiscence is common.²²

When Should Osteomyelitis or Other Infectious Complications Be Considered?

The decision to treat a pressure ulcer for infection should be a clinical one. Ideally, evidence of infection is confirmed by tissue or bone biopsy. Superficial cultures usually are not useful other than to rule out methicillin-resistant Staphylococcus aureus (MRSA) colonization. Systemic rather than topical antibiotics should be used to treat an infected pressure ulcer.^{1,32}

EVIDENCE SUMMARY

The diagnosis of wound infection is usually a clinical determination. Although tissue or bone biopsy has long been considered the preferred method to identify infection or osteomyelitis, this is often not practical in the clinical setting.^{1,32} All open pressure ulcers contain bacteria, which, unless they reach high colony counts, should not impair wound healing.²³ Therefore, culture is not usually helpful to determine infection, other than to rule out the presence of MRSA. If a superficial culture is taken, evidence suggests that the Levine technique should be used, which involves a swab rotated over a 1-cm square patch of wound with enough pressure to extract fluid from the wound for five seconds.³² When bone is exposed in a pressure ulcer, osteomyelitis can be clinically presumed.

Of all clinical signs of infection, a systematic review suggests that worsening pain (in patients with intact sensation) may be one of the best predictors of chronic wound infection.³² Other signs of an acute spreading infection may include erythema around the ulcer's edges, induration, warmth, purulent drainage, or worsening of

chronic drainage. Additional signs of infection include no evidence of healing for two weeks, friable granulation tissue, foul odor, new necrotic tissue, or lack of uniform spread of granulation tissue across the base of the wound (referred to as pocketing or bridging). Systemic signs may include fever, lymph node enlargement, or possible delirium or confusion.¹

Pressure ulcers that are large, have necrotic tissue or a foreign body, have been present a long time, or are repeatedly contaminated by stool are at greater risk of becoming infected. Patients at higher risk of infection are also at higher risk of developing ulcers initially, including patients with diabetes mellitus, undernutrition, poor tissue perfusion, or immune suppression.

To identify osteomyelitis on imaging, radiography is often used first, although magnetic resonance imaging and nuclear medicine tests are more sensitive and specific than conventional plain radiography.³³ Systemic antibiotics should be used to treat infected ulcers. When treating osteomyelitis, infectious disease consultation, if available, should be considered to help tailor antibiotic therapy. There is little role for the use of topical antibiotics.¹

Data Sources: A search was performed using Essential Evidence Plus. The term pressure ulcer was searched in the Cochrane Database of Systematic Reviews and the evidence-based guidelines from the National Guideline Clearinghouse. Additionally, a PubMed search of systematic reviews was done using the search term pressure ulcer. Search dates: January 2014 through July 2015.

The authors thank Elliot Twiggs, MD, for his assistance with Figure 3.

The Authors

JAQUELINE G.M. RAETZ, MD, is an assistant professor in the Department of Family Medicine at the University of Washington in Seattle.

KEREN H. WICK, PhD, is an associate professor and director of research and graduate programs at the MEDEX Northwest Physician Assistant Program at the University of Washington.

Address correspondence to Jaqueline G.M. Raetz, MD, University of Washington, 331 NE Thornton Place, Seattle, WA 98125 (e-mail: jraetz@uw.edu). Reprints are not available from the authors.

REFERENCES

- Haesler E, ed. National Pressure Ulcer Advisory Panel, European Pressure Ulcer Advisory Panel and Pan Pacific Pressure Injury Alliance. Prevention and treatment of pressure ulcers: quick reference guide. Perth, Australia: Cambridge Media; 2014. <http://www.npuap.org/wp-content/uploads/2014/08/Quick-Reference-Guide-DIGITAL-NPUAP-EPUAP-PPPIA.pdf>. Accessed August 4, 2015.
- Grey JE, Harding KG, Enoch S. Pressure ulcers. *BMJ*. 2006;332(7539):472-475.
- Chou R, Dana T, Bougatsos C, et al. Pressure ulcer risk assessment and prevention: a systematic comparative effectiveness review. *Ann Intern Med*. 2013;159(1):28-38.

Pressure Ulcers

- Smith ME, Totten A, Hickam DH, et al. Pressure ulcer treatment strategies: a systematic comparative effectiveness review. *Ann Intern Med.* 2013;159(1):39-50.
- Coleman S, Gorecki C, Nelson EA, et al. Patient risk factors for pressure ulcer development: systematic review. *Int J Nurs Stud.* 2013;50(7):974-1003.
- Shekelle PG, Pronovost PJ, Wachter RM, et al. The top patient safety strategies that can be encouraged for adoption now. *Ann Intern Med.* 2013;158(5 pt 2):365-368.
- Sullivan N, Schoelles KM. Preventing in-facility pressure ulcers as a patient safety strategy: a systematic review. *Ann Intern Med.* 2013;158(5 pt 2):410-416.
- McInnes E, Jammali-Blasi A, Bell-Syer SE, Dumville JC, Cullum N. Support surfaces for pressure ulcer prevention. *Cochrane Database Syst Rev.* 2011;(4):CD001735.
- Qaseem A, Mir TP, Starkey M, Denberg TD. Risk assessment and prevention of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2015;162(5):359-369.
- Gefen A. How much time does it take to get a pressure ulcer? Integrated evidence from human, animal, and in vitro studies. *Ostomy Wound Manage.* 2008;54(10):26-28,30-35.
- Moore ZE, Cowman S. Risk assessment tools for the prevention of pressure ulcers. *Cochrane Database Syst Rev.* 2014;(2):CD006471.
- Moore ZE, Webster J. Dressings and topical agents for preventing pressure ulcers. *Cochrane Database Syst Rev.* 2013;(8):CD009362.
- McGinnis E, Stubbs N. Pressure-relieving devices for treating heel pressure ulcers. *Cochrane Database Syst Rev.* 2014;(2):CD005485.
- Gillespie BM, Chaboyer WP, McInnes E, Kent B, Whitty JA, Thalib L. Repositioning for pressure ulcer prevention in adults. *Cochrane Database Syst Rev.* 2014;(4):CD009958.
- Whitney J, Phillips L, Aslam R, et al. Guidelines for the treatment of pressure ulcers. *Wound Repair Regen.* 2006;14(6):663-679.
- Institute for Clinical Systems Improvement (ICSI). Pressure ulcer prevention and treatment protocol. Health care protocol. Bloomington, Minn.: Institute for Clinical Systems Improvement (ICSI); January 2012. <http://www.guideline.gov/content.aspx?id=36059#Section420>. Accessed August 4, 2015.
- Reddy M, Gill SS, Rochon PA. Preventing pressure ulcers: a systematic review. *JAMA.* 2006;296(8):974-984.
- National Pressure Ulcer Advisory Panel. NPUAP pressure ulcer stages/categories. <http://www.npuap.org/resources/educational-and-clinical-resources/npuap-pressure-ulcer-stagescategories/>. Accessed August 4, 2015.
- National Pressure Ulcer Advisory Panel. Pressure ulcer category/staging illustrations. <http://www.npuap.org/resources/educational-and-clinical-resources/pressure-ulcer-categorystaging-illustrations/>. Accessed August 4, 2015.
- Moore ZE, Cowman S. Wound cleansing for pressure ulcers. *Cochrane Database Syst Rev.* 2013;(3):CD004983.
- Fernandez R, Griffiths R. Water for wound cleansing. *Cochrane Database Syst Rev.* 2012;(2):CD003861.
- Qaseem A, Humphrey LL, Forciea MA, Starkey M, Denberg TD. Treatment of pressure ulcers: a clinical practice guideline from the American College of Physicians. *Ann Intern Med.* 2015;162(5):370-379.
- Rodeheaver GT. Pressure ulcer debridement and cleansing: a review of current literature. *Ostomy Wound Manage.* 1999;45(1A suppl):80S-85S.
- Lyder CH, Ayello EA. Pressure ulcers: a patient safety issue. In: Hughes RG, ed. *Patient Safety and Quality: An Evidence-Based Handbook for Nurses*. Rockville, Md.: Agency for Healthcare Research and Quality; 2008. <http://www.ncbi.nlm.nih.gov/books/NBK2650/>. Accessed August 4, 2015.
- Kerstein MD, Gemmen E, van Rijswijk L, et al. Cost and cost effectiveness of venous and pressure ulcer protocols of care. *Dis Manag Health Outcomes.* 2001;9(11):651-663.
- Kroshinsky D, Strazzula L. Pressure ulcers. Merck Manuals Professional Edition. March 2013. <http://www.merckmanuals.com/professional/dermatologic-disorders/pressure-ulcers/pressure-ulcers/>. Accessed August 4, 2015.
- Dumville JC, Stubbs N, Keogh SJ, Walker RM, Liu Z. Hydrogel dressings for treating pressure ulcers. *Cochrane Database Syst Rev.* 2015;(2):CD011226.
- Langer G, Fink A. Nutritional interventions for preventing and treating pressure ulcers. *Cochrane Database Syst Rev.* 2014;(6):CD003216.
- Moore ZE, Cowman S. Repositioning for treating pressure ulcers. *Cochrane Database Syst Rev.* 2015;(1):CD006898.
- Kawasaki L, Mushahwar VK, Ho C, Dukelow SP, Chan LL, Chan KM. The mechanisms and evidence of efficacy of electrical stimulation for healing of pressure ulcer: a systematic review. *Wound Repair Regen.* 2014;22(2):161-173.
- Chen C, Hou WH, Chan ES, Yeh ML, Lo HL. Phototherapy for treating pressure ulcers. *Cochrane Database Syst Rev.* 2014;(7):CD009224.
- Reddy M, Gill SS, Wu W, Kalkar SR, Rochon PA. Does this patient have an infection of a chronic wound? *JAMA.* 2012;307(6):605-611.
- Pineda C, Vargas A, Rodriguez AV. Imaging of osteomyelitis: current concepts. *Infect Dis Clin North Am.* 2006;20(4):789-825.