

Implementing AHRQ Effective Health Care Reviews

Helping Clinicians Make Better Treatment Choices

Management of Binge-Eating Disorder in Adults

Practice Pointers by JULIANA LLANO, MD, *Georgetown University/Providence Hospital Family Medicine Residency Program, Washington, District of Columbia*, and KENNETH W. LIN, MD, *Georgetown University Medical Center, Washington, District of Columbia*



More online
at <http://www.aafp.org/afp>.

The Agency for Healthcare Research and Quality (AHRQ) conducts the Effective Health Care Program as part of its mission to produce evidence to improve health care and to make sure the evidence is understood and used. A key clinical question based on the AHRQ Effective Health Care Program systematic review of the literature is presented, followed by an evidence-based answer based upon the review. AHRQ's summary is accompanied by an interpretation by an AFP author that will help guide clinicians in making treatment decisions. For the full review, clinician summary, and consumer summary, go to <http://effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=2212>.

This series is coordinated by Kenny Lin, MD, MPH, Associate Deputy Editor for AFP Online.

A collection of Implementing AHRQ Effective Health Care Reviews published in AFP is available at <http://www.aafp.org/afp/ahrq>.

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz on page 288.

Author disclosure: No relevant financial affiliations.

Key Clinical Issue

What are the benefits and harms of treatments for adults with binge-eating disorder?

Evidence-Based Answer

Therapist-led cognitive behavior therapy (CBT) reduces binge-eating frequency and increases binge-eating abstinence. (Strength of Recommendation [SOR]: A, based on consistent, good-quality patient-oriented evidence.) In short-term studies (six to 16 weeks), lisdexamfetamine, second-generation antidepressants, and topiramate increased binge-eating abstinence and reduced binge-eating frequency and eating-related obsessions and compulsions. (SOR: B, based on inconsistent or limited-quality patient-oriented evidence.) Adverse effects of pharmacologic interventions were rarely severe (*eTable A*).

Practice Pointers

Binge-eating disorder is characterized by recurrent episodes of eating unusually large portions in a discrete period (two hours or less), at least once a week for three months. Core features include lack of control over eating, eating rapidly, feeling uncomfortably full, eating alone, and feeling guilty or depressed. Binge-eating disorder is not associated with inappropriate compensatory behaviors (e.g., purging).¹ The lifetime prevalence in the United States is 2.8%. It is more common in females, adolescents, and obese individuals. Severity is characterized by the number of episodes per week.²

Binge-eating disorder can be treated with psychological and behavior therapies (*eTable B*) and pharmacotherapy. Psychotherapeutic options include behavioral weight loss therapy, CBT, interpersonal

psychotherapy, and dialectical behavior therapy. CBT focuses on changing negative thoughts and undesirable behaviors.³ It can be therapist-led (individual or group sessions), guided self-help, or structured self-help. The U.S. Food and Drug Administration approved lisdexamfetamine for treatment of moderate to severe binge-eating disorder. Other treatments include topiramate and second-generation antidepressants.⁴

This Agency for Healthcare Research and Quality (AHRQ) review included 57 studies and one systematic review on the effectiveness and adverse effects of treatments for adults with binge-eating disorder. Meta-analyses provided strong evidence that lisdexamfetamine and second-generation antidepressants increase binge-eating abstinence. Second-generation antidepressants also decrease binge-eating frequency, and eating obsessions and compulsions. The U.S. Food and Drug Administration warns about the potential for abuse and dependence with lisdexamfetamine use, as well as an increased risk of sudden death, stroke, and myocardial infarction.²

Meta-analyses provided strong evidence that therapist-led CBT reduces binge-eating frequency and increases binge-eating abstinence. There is moderate evidence that behavioral weight loss therapy, which incorporates strategies such as caloric restriction and increased physical activity to promote weight loss, decreases body mass index more than CBT, but it is not clearly associated with reduced binge-eating behaviors. There was insufficient evidence to determine the effectiveness of other psychotherapies or combinations of pharmacologic and psychological treatments.

The AHRQ review is consistent with the American Psychiatric Association practice

Clinical Bottom Line: Summary of Key Findings for the Efficacy and Comparative Effectiveness of Interventions to Treat Binge-Eating Disorder

Psychological and behavior therapy interventions

Therapist-led CBT vs. wait-list*

CBT increased binge-eating abstinence (RR = 4.95; 95% CI, 3.06 to 8.00). ●●●

CBT decreased the frequency of binge-eating episodes per week (MD = -2.32; 95% CI, -4.56 to -0.09). ●●●

CBT decreased eating-related psychopathology. ●●●

No differences were found in body mass index or symptoms of depression. ●●○

Therapist-led CBT vs. behavioral weight loss therapy

Behavioral weight loss therapy decreased body mass index more than CBT at the end of treatment. ●●○

CBT decreased binge-eating frequency more than behavioral weight loss therapy at the end of treatment and for up to 12 months of follow-up. ●○○

No differences were found in binge-eating abstinence or symptoms of depression. ●○○

Pharmacologic interventions

Lisdexamfetamine† (a central nervous system stimulant) vs. placebo

Lisdexamfetamine increased binge-eating abstinence (RR = 2.61; 95% CI, 2.04 to 3.33). ●●●

Lisdexamfetamine decreased binge-eating days per week, weight, and eating-related obsessions and compulsions. ●●●

Second-generation antidepressants (as a class) vs. placebo

Antidepressants increased binge-eating abstinence (RR = 1.67; 95% CI, 1.24 to 2.26) and decreased the frequency of binge-eating episodes per week (MD = -0.67; 95% CI, -1.26 to -0.09). ●●●

Antidepressants decreased the frequency of binge-eating days per week (MD = -0.90; 95% CI, -1.48 to -0.32) and eating-related obsessions and compulsions. ●●○

Antidepressants decreased symptoms of depression (MD = -1.98; 95% CI, -3.67 to -0.28) ●○○

No differences were found in weight or body mass index. ●○○

Topiramate (an anticonvulsant) vs. placebo

Topiramate increased binge-eating abstinence and decreased binge-eating frequency, weight, and eating-related obsessions and compulsions. ●●○

Topiramate improved general and eating-related psychological functioning, and decreased impulsivity and disability in family and other social domains. ●○○

Strength of evidence scale

High: ●●● High confidence that the evidence reflects the true effect. Further research is very unlikely to change our confidence in the estimate of effect.

Moderate: ●●○ Moderate confidence that the evidence reflects the true effect. Further research may change our confidence in the estimate of effect and may change the estimate.

Low: ●○○ Low confidence that the evidence reflects the true effect. Further research is likely to change our confidence in the estimate of effect and is likely to change the estimate.

Insufficient: ○○○ Evidence either is unavailable or does not permit a conclusion.

CBT = cognitive behavior therapy; CI = confidence interval; FDA = U.S. Food and Drug Administration; MD = mean difference; RR = relative risk.

*—Wait-list refers to patients who received no treatment at all.

†—Lisdexamfetamine is not indicated by the FDA for weight loss. The FDA notes that the use of other sympathomimetic drugs for weight loss has been associated with serious cardiovascular adverse events, and the safety and effectiveness of lisdexamfetamine for the treatment of obesity have not been established.

Adapted from the Agency for Healthcare Research and Quality, Effective Health Care Program. Management and outcomes of binge-eating disorder in adults: current state of the evidence. Executive summary. Rockville, Md.: Agency for Healthcare Research and Quality; May 2016. <https://effectivehealthcare.ahrq.gov/ehc/products/563/2212/binge-eating-clinician-160517.pdf>. Accessed November 21, 2016.

guideline for the treatment of patients with eating disorders, which strongly recommends individual and group CBT for binge-eating disorders, as well as guided self-help programs.⁵ There is little evidence regarding the optimal duration of treatment. One study suggested that 16 sessions of CBT are more effective than eight sessions.⁶ Another controlled study found that a 10-month sequence of CBT followed by behavioral

weight loss therapy is not significantly superior to six-month courses of CBT or behavioral weight loss therapy alone.⁷

In 2011, the World Federation of Societies of Biological Psychiatry Task Force on Eating Disorders identified 26 randomized controlled trials of pharmacologic treatments for binge-eating disorder and concluded that evidence supports the use of imipramine, sertraline,

citalopram or escitalopram, and topiramate.⁸ The optimal treatment duration is unclear.

Based on the AHRQ review, a reasonable approach is to refer patients with binge-eating disorder to a therapist-led CBT program as first-line therapy. Patients who are unable to access a program or who do not respond to CBT may be prescribed lisdexamfetamine or a second-generation antidepressant. To find therapists who specialize in eating disorders, go to the <http://locator.apa.org/>.

EDITOR'S NOTE: American Family Physician *SOR ratings are different from the AHRQ Strength of Evidence (SOE) ratings.*

Address correspondence to Juliana Llano, MD, at jal320@georgetown.edu. Reprints are not available from the authors.

REFERENCES

1. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders*. 5th ed. Washington, DC: American Psychiatric Association; 2013.
2. Agency for Healthcare Research and Quality, Effective Health Care Program. Clinician summary. Management and outcomes of binge-eating disorder in adults: current state of the evidence. Rockville, Md.: Agency for Healthcare Research and Quality; May 2016. <http://effectivehealthcare.ahrq.gov/index.cfm/search-for-guides-reviews-and-reports/?pageaction=displayproduct&productid=2212>. Accessed November 21, 2016.
3. Coffey SF, Banducci AN, Vinci C. Common questions about cognitive behavior therapy for psychiatric disorders. *Am Fam Physician*. 2015; 92(9):807-812.
4. Devlin MJ. Binge-eating disorder comes of age. *Ann Intern Med*. 2016; 165(6):445-446.
5. Yager J, Devlin MJ, Halmi KA, et al.; American Psychiatric Association. Guideline watch (August 2012): practice guideline for the treatment of patients with eating disorders. 3rd ed. http://psychiatryonline.org/pb/assets/raw/sitewide/practice_guidelines/guidelines/eatingdisorders-watch.pdf. Accessed November 29, 2016.
6. Schlup B, Meyer AH, Munsch S. A non-randomized direct comparison of cognitive-behavioral short- and long-term treatment for binge eating disorder. *Obesity Facts*. 2010;3(4):261-266.
7. Grilo CM, Masheb RM, Wilson GT, Gueorguieva R, White MA. Cognitive-behavioral therapy, behavioral weight loss, and sequential treatment for obese patients with binge-eating disorder: a randomized controlled trial. *J Consult Clin Psychol*. 2011;79(5):675-685.
8. Aigner M, Treasure J, Kaye W, Kasper S; WFSBP Task Force on Eating Disorders. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for the pharmacological treatment of eating disorders. *World J Biol Psychiatry*. 2011;12(6):400-443. ■

NEW AAFP BENEFIT
Primary+
 Your CME and Professional Portfolio

Use this new AAFP member benefit to:

- Report CME
- Log procedures
- Track professional requirements
- And more

aafp.org/primary-plus

AMERICAN ACADEMY OF
 FAMILY PHYSICIANS

eTable A. Summary of Key Findings for the Adverse Effects of Pharmacologic Interventions to Treat Binge-Eating Disorder

Fluvoxamine vs. placebo

Fluvoxamine was associated with a higher number of events related to GI upset, sympathetic nervous system arousal, and sleep disturbance. ●○○

Lisdexamfetamine vs. placebo

Lisdexamfetamine was associated with greater insomnia (RR = 2.66; 95% CI, 1.63 to 4.31). ●●●

Lisdexamfetamine was associated with a greater risk of headache (RR = 1.63; 95% CI, 1.13 to 2.36). ●●●

Lisdexamfetamine was associated with a higher number of events related to GI upset, sympathetic nervous system arousal, and decreased appetite. ●●○

Topiramate vs. placebo

Topiramate was associated with a higher number of events related to sympathetic nervous system arousal. ●●○

Topiramate was associated with a higher number of other adverse events, including upper respiratory tract infection, taste perversion, difficulty with attention and memory, dizziness, confusion, and back pain. ●●○

No difference was found in the number of headaches. ●●○

No difference was found in the number of events related to GI upset or sleep disturbance. ●○○

Strength of evidence scale

High: ●●● High confidence that the evidence reflects the true effect. Further research is very unlikely to change our confidence in the estimate of effect.

Moderate: ●●○ Moderate confidence that the evidence reflects the true effect. Further research may change our confidence in the estimate of effect and may change the estimate.

Low: ●○○ Low confidence that the evidence reflects the true effect. Further research is likely to change our confidence in the estimate of effect and is likely to change the estimate.

Insufficient: ○○○ Evidence either is unavailable or does not permit a conclusion.

CI = confidence interval; GI = gastrointestinal; RR = relative risk.

Adapted from the Agency for Healthcare Research and Quality, Effective Health Care Program. Management and outcomes of binge-eating disorder in adults: current state of the evidence. Executive summary. Rockville, Md.: Agency for Healthcare Research and Quality; May 2016. <https://effectivehealthcare.ahrq.gov/ehc/products/563/2212/binge-eating-clinician-160517.pdf>. Accessed November 21, 2016.

eTable B. Psychological and Behavior Therapy Interventions for Binge-Eating Disorder

Behavioral weight loss therapy

Treatment that incorporates various behavior strategies, such as caloric restriction and increased physical activity, to promote weight loss.

CBT

Psychotherapy that focuses on identifying relationships among thoughts, feelings, and behaviors and aims to change participants' negative thoughts about themselves and the world and, by doing so, reduce negative emotions and undesirable behavior patterns. CBT is delivered in various ways (e.g., therapist-led individual and group sessions, structured self-help, and guided self-help).

In therapist-led CBT, a therapist is present for the duration of each group or individual session to provide psychoeducation, teach new skills, and support participants.

Dialectical behavior therapy

Psychotherapy that helps participants understand how negative feelings can lead to binge eating as a coping mechanism. It focuses on mindfulness, emotion regulation, and distress tolerance. It is delivered in individual sessions or as group therapy.

Interpersonal psychotherapy

Psychotherapy that helps participants understand how problems with social interactions can lead to binge eating as a coping mechanism. It helps participants learn to cope with negative emotions stemming from problems with social interactions and develop healthy interpersonal skills. It is delivered in individual sessions or as group therapy.

CBT = cognitive behavior therapy.

Adapted from the Agency for Healthcare Research and Quality, Effective Health Care Program. Management and outcomes of binge-eating disorder in adults: current state of the evidence. Executive summary. Rockville, Md.: Agency for Healthcare Research and Quality; May 2016. <https://effectivehealthcare.ahrq.gov/ehc/products/563/2212/binge-eating-clinician-160517.pdf>. Accessed November 21, 2016.