

Laceration Repair: A Practical Approach

RANDALL T. FORSCH, MD, MPH; SAHOKO H. LITTLE, MD, PhD; and CHRISTA WILLIAMS, MD
University of Michigan Medical School, Ann Arbor, Michigan

The goals of laceration repair are to achieve hemostasis and optimal cosmetic results without increasing the risk of infection. Many aspects of laceration repair have not changed over the years, but there is evidence to support some updates to standard management. Studies have been unable to define a “golden period” for which a wound can safely be repaired without increasing risk of infection. Depending on the type of wound, it may be reasonable to close even 18 or more hours after injury. The use of nonsterile gloves during laceration repair does not increase the risk of wound infection compared with sterile gloves. Irrigation with potable tap water rather than sterile saline also does not increase the risk of wound infection. Good evidence suggests that local anesthetic with epinephrine in a concentration of up to 1:100,000 is safe for use on digits. Local anesthetic with epinephrine in a concentration of 1:200,000 is safe for use on the nose and ears. Tissue adhesives and wound adhesive strips can be used effectively in low-tension skin areas. Wounds heal faster in a moist environment and therefore occlusive and semiocclusive dressings should be considered when available. Tetanus prophylaxis should be provided if indicated. Timing of suture removal depends on location and is based on expert opinion and experience. (*Am Fam Physician*. 2017;95(10):628-636. Copyright © 2017 American Academy of Family Physicians.)



More online
at <http://www.aafp.org/afp>.

CME This clinical content conforms to AAFP criteria for continuing medical education (CME). See CME Quiz Questions on page 622.

Author disclosure: No relevant financial affiliation.

► **Patient information:**
A handout on this topic is available at <http://www.aafp.org/afp/2008/1015/p952.html>.

Approximately 6 million patients present to emergency departments for laceration treatment every year.¹ Although many patients seek care at emergency departments or urgent care centers, primary care physicians are an important resource for urgent laceration treatment. Many aspects of laceration repair have not changed, but there is evidence to support some updates to standard management.

Approach to the Wound

The goals of laceration repair are to achieve hemostasis and optimal cosmetic results without increasing the risk of infection. Important considerations include timing of the repair, wound irrigation techniques, providing a clean field for repair to minimize contamination, and appropriate use of anesthesia. An article on wound care was previously published in *American Family Physician*.²

EVALUATING THE WOUND

When a patient presents with a laceration, the physician should obtain a history, including tetanus vaccination status, allergies, and time and mechanism of injury, and then assess wound size, shape, and location.³ If active bleeding persists after application of direct pressure, hemostasis should be obtained using hemostat, ligation, or sutures

before further evaluation. Hemostasis controls bleeding, prevents hematoma formation, and allows for deeper inspection of the wound.³ The next step is to determine whether vessels, tendons, nerves, joints, muscles, or bones are damaged. Anesthesia may be necessary to achieve hemostasis and to explore the wound. Devitalized and necrotic tissue in a traumatic wound should be identified and removed to reduce risk of infection.^{4,5}

If a foreign body (e.g., dirt particles, wood, glass) is suspected but cannot be identified visually, then radiography, ultrasonography, or computed tomography may be needed. About one-third of foreign bodies may be missed on initial inspection.⁶

Injuries that require subspecialist consultation include open fractures, tendon or muscle lacerations of the hand, nerve injuries that impair function, lacerations of the salivary duct or canaliculus, lacerations of the eyes or eyelids that are deeper than the subcutaneous layer, injuries requiring sedation for repair, or other injuries requiring treatment beyond the knowledge or skill of the physician.

TIMING OF WOUND CLOSURE

No randomized controlled trials (RCTs) have compared primary and delayed closure

SORT: KEY RECOMMENDATIONS FOR PRACTICE

<i>Clinical recommendation</i>	<i>Evidence rating</i>	<i>References</i>
Noninfected wounds caused by clean objects may undergo primary closure up to 18 hours after injury. Head wounds may be repaired up to 24 hours after injury.	B	2, 7-9
Using potable tap water instead of sterile saline for wound irrigation does not increase the risk of infection.	A	2, 10-12
Use of clean nonsterile examination gloves rather than sterile gloves during wound repair does not significantly increase risk of infection.	A	11, 18-20
If there is no concern for vascular compromise to an appendage, local anesthetic containing epinephrine in a concentration of up to 1:100,000 is safe for use in laceration repair of the digits, including for digital blockade.	B	29, 30

A = consistent, good-quality patient-oriented evidence; B = inconsistent or limited-quality patient-oriented evidence; C = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <http://www.aafp.org/afpsort>.

of nonbite traumatic wounds.⁷ One systematic review and a prospective cohort study of 2,343 patients found that lacerations repaired after 12 hours have no significant increase in infection risk compared with those repaired earlier.¹ A case series of 204 patients found no increased risk of infection in wounds repaired at less than 19 hours.⁸ Noninfected wounds caused by clean objects may undergo primary closure up to 18 hours after injury. Head wounds may be repaired up to 24 hours after injury.⁸ Factors that may increase the likelihood of infection include wound contamination, laceration length greater than 5 cm, laceration located on the lower extremities, and diabetes mellitus.⁹

WOUND IRRIGATION

Irrigation cleanses the wound of debris and dilutes bacterial load before closure. However, there is no strong evidence that cleansing a wound increases healing or reduces infection.¹⁰ A Cochrane review and several RCTs support the use of potable tap water, as opposed to sterile saline, for wound irrigation.^{2,10-13} To dilute the wound's bacterial load below the recommended 10⁵ organisms per mL,¹⁴ 50 to 100 mL of irrigation solution per 1 cm of wound length is needed.¹⁵ Optimal pressure for irrigation is around 5 to 8 psi.¹⁶ This can be achieved by using a 19-gauge needle with a 35-mL syringe or by placing the wound under a running faucet.^{16,17}

Physicians should wear protective gear, such as a mask with shield, during irrigation.

CLEAN VS. STERILE GLOVES

Use of clean nonsterile examination gloves, rather than sterile gloves, during wound repair has little to no impact on rate of subsequent wound infection. An RCT of 493 patients undergoing skin excision with primary closure revealed that clean gloves were not inferior to sterile gloves regarding infection risk.¹⁸ A larger RCT with 816 patients and good follow-up revealed no statistically significant difference in the incidence of infection between clean and sterile glove use.¹⁹ Smaller observational studies support these findings.^{11,20}

Lacerations are considered contaminated at presentation, and physicians should make every effort to avoid introducing additional bacteria to the wound. However, strict sterile techniques appear to be unnecessary. Sutures, needles, and other instruments that touch the wound should be sterile, but

everything else only needs to be clean.

ANESTHETIZING THE WOUND

Topical and injectable local anesthetics reduce pain during treatment of lacerations and may be used alone or in combination.²¹⁻²³ Topical anesthetics (*eTable A*) are particularly useful when treating children. Topical agents commonly used in the United States include lidocaine/epinephrine/tetracaine and lidocaine/prilocaine. Lidocaine/prilocaine is not approved by the U.S. Food and Drug Administration for use on nonintact skin, although it has been used this way in numerous studies.

When using an injectable local anesthetic, the pain associated with injection can be reduced by using a high-gauge needle, buffering the anesthetic, warming the anesthetic to body temperature, and injecting the anesthetic slowly.²⁴⁻²⁸ Lidocaine may be buffered by adding 1 mL of sodium bicarbonate to 9 mL of lidocaine 1% (with or without epinephrine).²⁷

If there is no concern for vascular compromise to an appendage, then local anesthetic containing epinephrine in a concentration of up to 1:100,000 is safe for use in laceration repair of the digits, including for digital blockade.^{29,30} Local anesthetic containing epinephrine in a concentration of 1:200,000 is safe for laceration repair of the nose and ears.³¹ A systematic review documents the safe use of lidocaine with epinephrine (in a

Laceration Repair

concentration up to 1:80,000) in more than 10,000 procedures involving digits without any reported incidence of necrosis.³⁰ Only two studies examined the safety of epinephrine-containing anesthetics in patients with peripheral vascular disease. Although no patients had ischemic complications, the studies were small. Concern for peripheral vascular compromise should be considered a contraindication to the use of an epinephrine-containing anesthetic.

Table 1. Laceration Closure Techniques

Technique	Comments
Simple interrupted sutures	General tissue approximation Can be used for most wounds
Simple running sutures	Fast and effective for long lacerations All sutures are lost if one suture is cut by mistake or removed for drainage
Horizontal mattress sutures (Figure 1)	Effective for everting wound edges Can cause skin necrosis and excessive scars
Vertical mattress sutures (Figure 2)	Most effective for everting wound edges Can cause skin necrosis and excessive scars
Half-buried mattress sutures (Figure 3)	Most effective in everting triangular wound edges in flap repair
Running subcuticular sutures	Fast and effective in accurate skin edge apposition Does not allow for drainage Suited for closing clean wounds, such as surgical wounds in the operating room
Interrupted dermal sutures	Effective in accurate skin edge apposition and wound eversion Allows for minimal drainage Suited for closing clean wounds
Staples	Fast, creates loose closure Allows for drainage Suited for unclean wounds Should be avoided if cosmetic outcome is important
Wounds adhesive strips	Fast, no anesthesia required Used to approximate clean, simple, small lacerations with little tension and without bleeding
Tissue adhesive	Fast, no anesthesia required Used to approximate clean, simple, small lacerations with little tension and without bleeding

NOTE: For a video of suture techniques, see <https://www.youtube.com/watch?v=-ZWUgKiBxfk>.

Wound Repair

Laceration closure techniques are summarized in *Table 1*. For a video of suturing techniques, see <https://www.youtube.com/watch?v=-ZWUgKiBxfk>. There are no significant studies to guide technique choice. Compared with multilayer repair, single layer repair has similar cosmetic results for facial lacerations³² and is faster and more cost-effective for scalp lacerations.³³ Running sutures reportedly have less dehiscence than interrupted sutures in surgical wounds.³⁴ Mattress sutures (*Figures 1³⁵ and 2³⁵*) are effective for everting wound edges.^{36,37} Half-buried mattress sutures are useful for everting triangular edges in flap repair (*Figure 3*). Cosmetic outcomes of facial wounds repaired without deep dermal sutures are similar to layered closure.³⁷ The approach to repair varies by wound location. Nonbite and bite wounds are treated differently because of differences in infection risk. *Figure 4* is an algorithm for the management of lacerations.

FACIAL LACERATIONS

Debridement of facial wounds should be conservative because of increased blood supply to the face. Removing subcutaneous fat may lead to depression of the scar.³⁸ Single layer 5-0 or 6-0 nylon sutures are sufficient.³²

LIP LACERATION THROUGH VERMILION BORDER

An optimal cosmetic result depends on reapproximation of the vermilion border. Therefore, the first skin suture should be placed at this border. The border should be marked before anesthetic injection because the anesthetic may blur the border. The muscle layer and oral mucosa should be repaired with 3-0 or 4-0 absorbable sutures, and skin should be repaired with 6-0 or 7-0 nylon sutures.

EYELID

The patient should be referred to ophthalmology if the laceration involves the eye itself, the tarsal plate, or the eyelid margin, or penetrates deeper than the subcutaneous layer. Laceration through the portion of the upper or lower lid medial to the punctum often damages the lacrimal duct or the




ILLUSTRATION BY RENEE CANNON

Figure 1. Horizontal mattress sutures.

Reprinted with permission from Forsch RT. *Essentials of skin laceration repair*. Am Fam Physician. 2008;78(8):948.




ILLUSTRATION BY RENEE CANNON

Figure 2. Vertical mattress sutures.

Reprinted with permission from Forsch RT. *Essentials of skin laceration repair*. Am Fam Physician. 2008;78(8):948.




ILLUSTRATION BY RENEE CANNON

Figure 3. Half-buried mattress sutures.

medial canthal ligament and requires referral to an ophthalmologist or plastic surgeon. Laceration of upper or lower eyelid skin can be repaired with 6-0 nylon sutures.

EYEBROW

The edges of the eyebrow serve as landmarks, so the eyebrow should not be shaved. Placing a single suture at each margin first ensures good alignment.³⁷

EAR

Cartilage has poor circulation and is prone to infection and necrosis. It needs to be covered with skin to heal. A single bite with reverse cutting needle or tapered needle (6-0 polypropylene sutures) should be used to approximate skin and perichondrium simultaneously. Ear trauma often causes a hematoma, and applying a pressure dressing can be difficult. Fluffed gauze under a circumferential head wrap can achieve adequate pressure to prevent a hematoma.

SCALP

A rich blood supply to the scalp causes lacerations to bleed significantly. After ruling out intracranial injury, bleeding should be controlled with direct pressure for adequate exploration of the wound. Shaving the area is rarely necessary. If the galea is lacerated more than 0.5 cm it should be repaired with 2-0 or 3-0 absorbable sutures.³⁹ Skin can be repaired using staples; interrupted, mattress, or running sutures, such as 3-0 or 4-0 nylon sutures; or the hair apposition technique (Figure 5³⁵). Staples are faster and more cost-effective than sutures with no difference in complications.⁴⁰ The hair apposition technique using tissue adhesive has the lowest cost and highest patient satisfaction for scalp repair.⁴¹ A video of the hair apposition technique is available at <https://lacerationrepair.com/alternative-wound-closure/hair-apposition-technique/>.

HANDS AND FOREARM

Lacerations of the fingers, hands, and forearms can be repaired by a family physician if deep tissue injury is not suspected. These lacerations are repaired with 4-0 or 5-0 nylon sutures. Any suspicion of injury involving tendon, nerve, muscle, vessels, bone, or the nail bed warrants immediate referral to a hand surgeon. Traditionally, a large subungual hematoma involving more than 25% of the visible nail indicated nail removal for nail bed inspection and repair, but a recent review concluded that a subungual hematoma without significant fingertip injury can be treated with trephining (drainage through a hole) alone.⁴²

Management of Acute Lacerations




Figure 4. Algorithm for the management of acute lacerations.

BITE WOUNDS

Up to 19% of bite wounds become infected. Cat bites are much more likely to become infected compared with dog or human bites (47% to 58% of cat bites, 8% to 14% of dog bites, and 7% to 9% of human bites).⁴³ The risk of infection increases as time from injury to repair increases, regardless of suture material.⁴ Evidence on optimal timing of primary closure and antibiotic treatment is lacking.^{4,44}

Cosmesis was improved with suturing compared with no suturing in RCTs of patients with dog bites, although the infection rate was the same.^{44,45} Therefore, dog bite wounds should be repaired, especially facial wounds because they are less prone to infection.^{4,46} Cat bites, with higher infection rates, have better outcomes without primary closure, especially when not located on the face or scalp. Bite wounds with a high risk of infection, such as cat bites, deep puncture wounds, or wounds longer than 3 cm,⁴³ should be treated with prophylactic amoxicillin/clavulanate (Augmentin).^{47,48} Clindamycin may be used in patients with a penicillin allergy.⁴⁹




Figure 5. Hair apposition technique for laceration closure. Opposing strands of hair are brought together with a simple twist and are secured with a drop of tissue adhesive.

Reprinted with permission from Forsch RT. Essentials of skin laceration repair. *Am Fam Physician*. 2008;78(8):949.

Table 2. Commonly Used Suture Materials

Material	Common needle type*	Time to lose 50% strength	Configuration	Typical use
Absorbable				
Chromic	Reverse cutting	10 to 14 days	Monofilament	Mucosa, eye wounds
Glycolide/lactide polymer (polyglactin 910 [Vicryl])	Conventional or reverse cutting	2 to 3 weeks	Braided	Deep dermal, muscle, fascia, oral mucosa, genitalia wounds
Polyglactone (Monocryl)	Conventional and reverse cutting	7 to 10 days	Monofilament	Dermal, subcuticular wounds
Polydioxanone (PDS II)	Reverse cutting	4 weeks	Monofilament	Muscle, fascia, dermal wounds
Nonabsorbable				
Nylon (Ethilon)	Cutting edge	> 10 years	Monofilament	Skin
Polypropylene (Prolene)	Tapered point, blunt tip	Indefinite	Monofilament	Mostly used in vascular surgeries; can be used for skin, tendon, and ligaments, depending on the needles
Silk	Does not come with needle	1 year	Braided	Used for hemostasis in ligation of vessels or for tying over bolsters

*—A variety of needles are available to order, but the most typical needles likely to be stocked are listed.

Information from references 50 and 51.

CHOOSING THE APPROPRIATE SUTURE MATERIAL

Physicians should use the smallest suture that will give sufficient strength to reapproximate and support the healing wound.^{50,51} Commonly used sutures are included in *Table 2*^{50,51}; however, good evidence is lacking regarding the appropriate suture size for laceration repair. The 5-0 or 6-0 sutures should be used for the face, and 4-0 sutures should be used for most other areas. The 3-0 sutures work well for the thicker skin on the back, scalp, palms, and soles.^{50,51}

A meta-analysis of 19 studies of skin closure for surgical wounds and traumatic lacerations found no significant difference in cosmetic outcome, wound infection, or wound dehiscence between absorbable and nonabsorbable sutures.^{52,53} A systematic review did not show any advantage of monofilament sutures over braided sutures with regard to cosmetic outcome, wound infection, or wound dehiscence.⁵⁴

USE OF TISSUE ADHESIVE OR WOUND ADHESIVE STRIPS

The two types of tissue adhesive available in the United States are *n*-butyl-2-cyanoacrylate (Histoacryl Blue, PeriAcryl) and 2-octyl cyanoacrylate (Dermabond, Surgiseal). *Table 3* shows the criteria for tissue adhesive use. A Cochrane review found these adhesives to be comparable in cosmesis, procedure time, discomfort, and complications.⁵⁵ They work well in clean, linear wounds that are not under tension. They are not generally used in hair-bearing areas (except in the hair apposition technique). There is a slightly higher likelihood of wound dehiscence with tissue adhesives than with sutures, with a number needed to harm of 25 for tissue adhesives.^{52,53}

Tissue adhesive should not be applied to misaligned wound edges. Care should be taken to avoid getting tissue adhesive into the wound or accidentally adhering gauze or instruments to the wound. If tissue adhesive is misapplied, it should be wiped off quickly with dry gauze. To remove dry adhesive, petroleum-based ointment should be applied and wiped away after 30 minutes.

Wound adhesive strips can also be used. One analysis suggests that wound adhesive strips are the most cost-effective method of closure for appropriate low-tension wounds.⁵⁶ The strips are applied perpendicular to the vector of the wound to approximate and secure the edges. One study found the same cosmetic outcomes

Table 3. Criteria for Use of Tissue Adhesives

Wound less than 12 hours old
Linear (not stellate)
Hemostatic
Not crossing a joint
Not crossing a mucocutaneous junction
Not in a hair-bearing area (unless hair apposition technique is being used)
Not under significant tension (or tension relieved with deep absorbable sutures)
Not grossly contaminated
Not infected
Not devitalized
Not a result of mammalian bite
No chronic condition that might impair wound healing

Laceration Repair

with adhesive strips vs. tissue adhesive when used to repair facial lacerations.⁵⁷

Laceration Aftercare

Once a wound has been adequately repaired, consideration should be given to the elements of aftercare. Although patients have traditionally been instructed to keep wounds covered and dry for 24 hours, one study found that uncovering wounds for routine bathing within the first 12 hours after closure did not increase the risk of infection.⁵⁸

A small prospective study showed that traumatic lacerations repaired with sutures had lower rates of infection when antibiotic ointment was applied rather than petroleum jelly. The lowest rate of infection occurred with the use of an ointment containing bacitracin and neomycin.⁵⁹ Therefore, topical antibiotic ointment should be applied to traumatic lacerations repaired with sutures unless the patient has a specific antibiotic allergy. A meta-analysis did not show benefit with the use of prophylactic systemic antibiotics for reducing wound infections in simple, nonbite wounds.⁶⁰

Wounds heal most quickly in a moist environment.⁶¹ Occlusive and semioclusive dressings lead to faster wound healing, decreased wound contamination, decreased infection rates, and increased comfort compared with dry gauze dressings.⁶² Choice of moisture retentive dressing should be based on the amount of exudate expected. Transparent film (e.g., Tegaderm) and hydrocolloid dressings are readily available and suited for repaired wounds without drainage. Film dressings allow for visualization of the wound to monitor for signs of infection. Gauze dressings with petroleum gel with or without an antibiotic are commonly used for wounds with some drainage. Foam dressings are more absorptive but mostly used for chronically draining wounds. When using interactive dressings such as film dressings, hydrocolloid dressings, or foam dressings, they should be changed according to package recommendations, which is anywhere from three to seven days or when fluid accumulation separates the dressing from the surrounding skin.⁶²

Patients with contaminated or high-risk (e.g., deep puncture) wounds who have not had a tetanus booster for more than five years should receive a tetanus vaccine. Patients who have not had at least three doses of a tetanus vaccine or who have an unknown tetanus

Table 4. Tetanus Wound Management

Tetanus vaccination history	Clean, minor wounds		Contaminated or high-risk wounds*	
	Tdap or Td†	Tetanus immune globulin	Tdap or Td†	Tetanus immune globulin
Unknown or fewer than 3 doses	Yes	No	Yes	Yes
3 or more doses	No‡	No	No§	No

Td = tetanus and diphtheria toxoids; Tdap = tetanus toxoid, reduced diphtheria toxoid, and acellular pertussis.

*—Examples are wounds contaminated with dirt, feces, soil, or saliva; deep puncture wounds; avulsions; and wounds resulting from missiles, crushing injury, burns, or frostbite.

†—Tdap is preferred over Td for adults who have never received Tdap. Single-antigen tetanus toxoid is no longer available in the United States.

‡—Yes, if it has been more than 10 years since the last dose of a tetanus toxoid-containing vaccine.

§—Yes, if it has been more than 5 years since the last dose of a tetanus toxoid-containing vaccine.

Adapted from Tetanus. In: Hamborsky J, Kroger A, Wolfe C, eds. Epidemiology and Prevention of Vaccine-Preventable Diseases. Atlanta, Ga.: Centers for Disease Control and Prevention; 2015:344.

vaccine history should also receive a tetanus immune globulin. Patients with a clean and minor wound should receive the tetanus vaccine only if they have not had a tetanus vaccine for more than 10 years. Tetanus immune globulin is not indicated for clean, minor wounds (Table 4).⁶³

Sutures should be removed after an appropriate interval depending on location (Table 5³⁵). This is based on expert opinion and experience.

Table 5. Timing of Suture or Staple Removal

Wound location	Timing of removal (days)
Face	3 to 5
Scalp	7 to 10
Arms	7 to 10
Trunk	10 to 14
Legs	10 to 14
Hands or feet	10 to 14
Palms or soles	14 to 21

Adapted with permission from Forsch RT. Essentials of skin laceration repair. Am Fam Physician. 2008;78(8):950.

This article updates previous articles on this topic by Forsch³⁵ and by Zuber.⁶⁴

Data Sources: The authors used an Essential Evidence summary based on the key words facial laceration, laceration, and tissue adhesives. The search included relevant POEMs, Cochrane reviews, diagnostic test data, and a custom PubMed search. Key words were skin laceration, skin repair, local anesthesia, sterile technique, sterile gloves, and wound irrigation. Search dates: April 2015 and January 5, 2017.

The Authors

RANDALL T. FORSCH, MD, MPH, is an assistant professor in the Department of Family Medicine at the University of Michigan Medical School in Ann Arbor.

SAHOKO H. LITTLE, MD, PhD, is an assistant professor in the Department of Family Medicine at the University of Michigan Medical School. She is also an attending physician at the Comprehensive Wound Care Clinic, University of Michigan.

CHRISTA WILLIAMS, MD, is a clinical lecturer in the Department of Family Medicine at the University of Michigan Medical School.

Address correspondence to Randall T. Forsch, MD, MPH, University of Michigan Medical School, 1301 Catherine, Ann Arbor, MI 48109-5624 (e-mail: rforsch@umich.edu). Reprints are not available from the authors.

REFERENCES

- Zehlabchi S, Tan A, Yadav K, Badawy A, Lucchesi M. The impact of wound age on the infection rate of simple lacerations repaired in the emergency department. *Injury*. 2012;43(11):1793-1798.
- Worster B, Zawora MQ, Hsieh C. Common questions about wound care. *Am Fam Physician*. 2015;91(2):86-92.
- American College of Emergency Physicians. Clinical policy for the initial approach to patients presenting with penetrating extremity trauma. *Ann Emerg Med*. 1994;23(5):1147-1156.
- Edlich RF, Thacker JG, Buchanan L, Rodeheaver GT. Modern concepts of treatment of traumatic wounds. *Adv Surg*. 1979;13:169-197.
- Haury B, Rodeheaver G, Vensko J, Edgerton MT, Edlich RF. Debridement: an essential component of traumatic wound care. *Am J Surg*. 1978;135(2):238-242.
- Anderson MA, Newmeyer WL III, Kilgore ES Jr. Diagnosis and treatment of retained foreign bodies in the hand. *Am J Surg*. 1982;144(1):63-67.
- Eliya-Masamba MC, Banda GW. Primary closure versus delayed closure for non bite traumatic wounds within 24 hours post injury. *Cochrane Database Syst Rev*. 2013;(10):CD008574.
- Berk WA, Osbourne DD, Taylor DD. Evaluation of the 'golden period' for wound repair: 204 cases from a third world emergency department. *Ann Emerg Med*. 1988;17(5):496-500.
- Quinn JV, Polevoi SK, Kohn MA. Traumatic lacerations: what are the risks for infection and has the 'golden period' of laceration care disappeared? *Emerg Med J*. 2014;31(2):96-100.
- Fernandez R, Griffiths R. Water for wound cleansing. *Cochrane Database Syst Rev*. 2012;(2):CD003861.
- Xia Y, Cho S, Greenway HT, Zelac DE, Kelley B. Infection rates of wound repairs during Mohs micrographic surgery using sterile versus nonsterile gloves: a prospective randomized pilot study. *Dermatol Surg*. 2011;37(5):651-656.
- Moscato RM, Mayrose J, Reardon RF, Janicke DM, Jehle DV. A multicenter comparison of tap water versus sterile saline for wound irrigation. *Acad Emerg Med*. 2007;14(5):404-409.
- Weiss EA, Oldham G, Lin M, Foster T, Quinn JV. Water is a safe and effective alternative to sterile normal saline for wound irrigation prior to suturing: a prospective, double-blind, randomised, controlled clinical trial. *BMJ Open*. 2013;3(1).
- Marshall KA, Edgerton MT, Rodeheaver GT, Magee CM, Edlich RF. Quantitative microbiology: its application to hand injuries. *Am J Surg*. 1976;131(6):730-733.
- Wheeler CB, Rodeheaver GT, Thacker JG, Edgerton MT, Edlich RF. Side-effects of high pressure irrigation. *Surg Gynecol Obstet*. 1976;143(5):775-778.
- Moscato RM, Reardon RF, Lerner EB, Mayrose J. Wound irrigation with tap water. *Acad Emerg Med*. 1998;5(11):1076-1080.
- Singer AJ, Hollander JE, Subramanian S, Malhotra AK, Villez PA. Pressure dynamics of various irrigation techniques commonly used in the emergency department. *Ann Emerg Med*. 1994;24(1):36-40.
- Heal C, Sriharan S, Buttner PG, Kimber D. Comparing non-sterile to sterile gloves for minor surgery: a prospective randomised controlled non-inferiority trial. *Med J Aust*. 2015;202(1):27-31.
- Perelman VS, Francis GJ, Rutledge T, Foote J, Martino F, Dranitsaris G. Sterile versus nonsterile gloves for repair of uncomplicated lacerations in the emergency department: a randomized controlled trial. *Ann Emerg Med*. 2004;43(3):362-370.
- Creamer J, Davis K, Rice W. Sterile gloves: do they make a difference? *Am J Surg*. 2012;204(6):976-979.
- Adler AJ, Dubinsky I, Eisen J. Does the use of topical lidocaine, epinephrine, and tetracaine solution provide sufficient anesthesia for laceration repair? *Acad Emerg Med*. 1998;5(2):108-112.
- Ernst AA, Marvez-Valls E, Nick TG, Weiss SJ. LAT (lidocaine-adrenaline-tetracaine) versus TAC (tetracaine-adrenaline-cocaine) for topical anesthesia in face and scalp lacerations. *Am J Emerg Med*. 1995;13(2):151-154.
- Eidelman A, Weiss JM, Enu IK, Lau J, Carr DB. Comparative efficacy and costs of various topical anesthetics for repair of dermal lacerations: a systematic review of randomized, controlled trials. *J Clin Anesth*. 2005;17(2):106-116.
- Young KD. What's new in topical anesthesia. *Clin Pediatr Emerg Med*. 2007;8(4):232-239.
- Scarfone RJ, Jasani M, Gracely EJ. Pain of local anesthetics: rate of administration and buffering. *Ann Emerg Med*. 1998;31(1):36-40.
- Bartfield JM, Gennis P, Barbera J, Breuer B, Gallagher EJ. Buffered versus plain lidocaine as a local anesthetic for simple laceration repair. *Ann Emerg Med*. 1990;19(12):1387-1389.
- Fatovich DM, Jacobs IG. A randomized controlled trial of buffered lidocaine for local anesthetic infiltration in children and adults with simple lacerations. *J Emerg Med*. 1999;17(2):223-228.
- Hogan ME, vanderVaart S, Perampaladas K, Machado M, Einarson TR, Taddio A. Systematic review and meta-analysis of the effect of warming local anesthetics on injection pain. *Ann Emerg Med*. 2011;58(1):86-98.e1.
- Chowdhry S, Seidenstricker L, Cooney DS, Hazani R, Wilhelmi BJ. Do not use epinephrine in digital blocks: myth or truth? Part II. A retrospective review of 1111 cases. *Plast Reconstr Surg*. 2010;126(6):2031-2034.
- Shridharani SM, Manson PN, Magarakis M, Broyles JM, Whitaker IS, Rodriguez ED. The safety and efficacy of epinephrine in hand surgery: a systematic review of the literature and international survey. *Eur J Plast Surg*. 2014;37(4):183-188.
- Häfner HM, Röcken M, Breuninger H. Epinephrine-supplemented local anesthetics for ear and nose surgery: clinical use without complications in more than 10,000 surgical procedures. *J Dtsch Dermatol Ges*. 2005;3(3):195-199.
- Singer AJ, Gulla J, Hein M, Marchini S, Chale S, Arora BP. Single-layer versus double-layer closure of facial lacerations: a randomized controlled trial. *Plast Reconstr Surg*. 2005;116(2):363-368.
- Adeolu AA, Olabanji JK, Komolafe EO, Ademuyiwa AO, Awe AO, Oladele AO. A prospective study of two methods of closing surgical scalp wounds. *Br J Neurosurg*. 2012;26(1):75-77.

Laceration Repair

34. Iwase K, Higaki J, Tanaka Y, Kondoh H, Yoshikawa M, Kamiike W. Running closure of clean and contaminated abdominal wounds using a synthetic monofilament absorbable looped suture. *Surg Today*. 1999; 29(9):874-879.
35. Forsch RT. Essentials of skin laceration repair. *Am Fam Physician*. 2008; 78(8):945-951.
36. Jones JS, Gartner M, Drew G, Pack S. The shorthand vertical mattress stitch: evaluation of a new suture technique. *Am J Emerg Med*. 1993; 11(5):483-485.
37. Lammers RL, Smith ZE. Methods of wound closure. In: Roberts JR, Custalow CB, Thomsen TW, Hedges JR, eds. *Roberts and Hedges' Clinical Procedures in Emergency Medicine*. Philadelphia, Pa.: Elsevier/Saunders; 2014.
38. Grabb WC, Klainert HE, eds. *Techniques in Surgery: Facial and Hand Injuries*. Somerville, NJ: Ethicon Inc.; 1980.
39. Hollander JE, Singer AJ, Valentine SM, Shofer FS. Risk factors for infection in patients with traumatic lacerations. *Acad Emerg Med*. 2001;8(7): 716-720.
40. Kanegaye JT, Vance CW, Chan L, Schonfeld N. Comparison of skin stapling devices and standard sutures for pediatric scalp lacerations: a randomized study of cost and time benefits. *J Pediatr*. 1997;130(5): 808-813.
41. Kavalci C, Cevik Y, Durukan P, Sayhan MB. Comparison of different suture techniques. *J Clin Anal Med*. 2015;6(1):15-17.
42. Batrick N, Hashemi K, Freij R. Treatment of uncomplicated subungual haematoma. *Emerg Med J*. 2003;20(1):65.
43. Jaindl M, Oberleitner G, Endler G, Thallinger C, Kovar FM. Management of bite wounds in children and adults—an analysis of over 5000 cases at a level I trauma centre. *Wien Klin Wochenschr*. 2016;128(9-10): 367-375.
44. Medeiros I, Saconato H. Antibiotic prophylaxis for mammalian bites. *Cochrane Database Syst Rev*. 2001;(2):CD001738.
45. Paschos NK, Makris EA, Gantsos A, Georgoulis AD. Primary closure versus non-closure of dog bite wounds. a randomised controlled trial. *Injury*. 2014;45(1):237-240.
46. Rui-feng C, Li-song H, Ji-bo Z, Li-qiu W. Emergency treatment on facial laceration of dog bite wounds with immediate primary closure: a prospective randomized trial study. *BMC Emerg Med*. 2013;13(suppl 1):S2.
47. Henton J, Jain A. Cochrane corner: antibiotic prophylaxis for mammalian bites (intervention review). *J Hand Surg Eur Vol*. 2012;37(8): 804-806.
48. Evgeniou E, Markeson D, Iyer S, Armstrong A. The management of animal bites in the United Kingdom. *Eplasty*. 2013;13:e27.
49. Ellis R, Ellis C. Dog and cat bites. *Am Fam Physician*. 2014;90(4):239-243.
50. *Wound Closure Manual*. Somerville, NJ: Ethicon Inc.; 2007.
51. Bullocks JM. *Plastic Surgery Emergencies: Principles and Techniques*. New York, NY: Thieme; 2008.
52. Mouzas GL, Yeadon A. Does the choice of suture material affect the incidence of wound infection? A comparison of dexon (polyglycolic acid) sutures with other commonly used sutures in an accident and emergency department. *Br J Surg*. 1975;62(12):952-955.
53. Xu B, Xu B, Wang L, et al. Absorbable versus nonabsorbable sutures for skin closure: a meta-analysis of randomized controlled trials. *Ann Plast Surg*. 2016;76(5):598-606.
54. Slieker JC, Daams F, Mulder IM, Jeekel J, Lange JF. Systematic review of the technique of colorectal anastomosis. *JAMA Surg*. 2013;148(2): 190-201.
55. Farion K, Osmond MH, Hartling L, et al. Tissue adhesives for traumatic lacerations in children and adults. *Cochrane Database Syst Rev*. 2002; (3):CD003326.
56. Zempysky WT, Zehrer CL, Lyle CT, Hedblom EC. Economic comparison of methods of wound closure: wound closure strips vs. sutures and wound adhesives. *Int Wound J*. 2005;2(3):272-281.
57. Zempysky WT, Parrotti D, Grem C, Nichols J. Randomized controlled comparison of cosmetic outcomes of simple facial lacerations closed with Steri Strip Skin Closures or Dermabond tissue adhesive. *Pediatr Emerg Care*. 2004;20(8):519-524.
58. Heal C, Buettner P, Raasch B, et al. Can sutures get wet? Prospective randomised controlled trial of wound management in general practice. *BMJ*. 2006;332(7549):1053-1056.
59. Hood R, Shermock KM, Emerman C. A prospective, randomized pilot evaluation of topical triple antibiotic versus mupirocin for the prevention of uncomplicated soft tissue wound infection. *Am J Emerg Med*. 2004;22(1):1-3.
60. Cummings P, Del Beccaro MA. Antibiotics to prevent infection of simple wounds: a meta-analysis of randomized studies. *Am J Emerg Med*. 1995;13(4):396-400.
61. Winter GD. Formation of the scab and the rate of epithelization of superficial wounds in the skin of the young domestic pig. *Nature*. 1962; 193:293-294.
62. Korting HC, Schöllmann C, White RJ. Management of minor acute cutaneous wounds: importance of wound healing in a moist environment. *J Eur Acad Dermatol Venereol*. 2011;25(2):130-137.
63. Tetanus. In: Hamborsky J, Kroger A, Wolfe C, eds. *Epidemiology and Prevention of Vaccine-Preventable Diseases*. Atlanta, Ga.: Centers for Disease Control and Prevention; 2015.
64. Zuber TJ. The mattress sutures: vertical, horizontal, and corner stitch. *Am Fam Physician*. 2002;66(12):2231-2236.

eTable A. Topical Anesthetics for Laceration Repair

<i>Agent</i>	<i>Forms</i>	<i>Recommended age</i>	<i>Dosage</i>	<i>Application</i>	<i>Onset of action</i>	<i>Duration of action</i>
Lidocaine/ epinephrine/ tetracaine	Solution, gel	Older than 1 month	Up to 3 mL	Apply with a cotton-tipped applicator or soaked cotton ball	20 to 30 minutes	1 hour
Lidocaine/ prilocaine*	Cream	Older than 3 months for nonintact skin; any age for intact skin	Term neonate \geq 37 weeks to 2 months of age: maximum of 1 g on 10 cm ² for 1 hour 3 to 11 months of age: maximum of 2 g on 20 cm ² for 1 hour 1 to 5 years of age: maximum of 10 g on 100 cm ² for 4 hours \geq 5 years of age: maximum of 20 g on 200 cm ² for 4 hours	Apply to intact skin with an occlusive cover	Peaks around 60 minutes	1 to 4 hours

NOTE: Topical anesthetics are used for lacerations less than 5 cm long and are most effective on the scalp and face. They should be avoided on the digits, nose, and ear lobes.

*—Lidocaine/prilocaine is not approved by the U.S. Food and Drug Administration for nonintact skin, although it has been used this way in numerous studies.

Information from Young KD. What's new in topical anesthesia. Clin Pediatr Emerg Med. 2007;8(4):232-239.