

## Evidence-Based Approach to Irritable Bowel Syndrome

Joel Heidelbaugh, MD, FAAFP, FACG

FMX

## ACTIVITY DISCLAIMER

The material presented here is being made available by the American Academy of Family Physicians for educational purposes only. Please note that medical information is constantly changing; the information contained in this activity was accurate at the time of publication. This material is not intended to represent the only, nor necessarily best, methods or procedures appropriate for the medical situations discussed. Rather, it is intended to present an approach, view, statement, or opinion of the faculty, which may be helpful to others who face similar situations.

The AAFP disclaims any and all liability for injury or other damages resulting to any individual using this material and for all claims that might arise out of the use of the techniques demonstrated therein by such individuals, whether these claims shall be asserted by a physician or any other person. Physicians may care to check specific details such as drug doses and contraindications, etc., in standard sources prior to clinical application. This material might contain recommendations/guidelines developed by other organizations. Please note that although these guidelines might be included, this does not necessarily imply the endorsement by the AAFP.

FMX

## DISCLOSURE

It is the policy of the AAFP that all individuals in a position to control content disclose any relationships with commercial interests upon nomination/invitation of participation. Disclosure documents are reviewed for potential conflict of interest (COI), and if identified, conflicts are resolved prior to confirmation of participation. Only those participants who had no conflict of interest or who agreed to an identified resolution process prior to their participation were involved in this CME activity.

All individuals in a position to control content for this session have indicated they have no relevant financial relationships to disclose.

The content of my material/presentation in this CME activity will include discussion of unapproved or investigational uses of products or devices as indicated:

Faculty will briefly discuss the following commonly-used treatments for irritable bowel syndrome (IBS), which have not been approved by the U.S. FDA for treatment of various associated symptoms:

- Anti-spasmodics and anti-depressants (TCAs, and SSRIs) for the treatment of abdominal cramping and bloating
- Probiotics for the treatment of abdominal bloating
- Fiber, milk of magnesia (MOM) and polyethylene glycol (PEG) for the treatment of constipation
- Loperamide, diphenoxylate/atropine, and cholestyramine for the treatment of diarrhea

FMX

## Joel Heidelbaugh, MD, FAAFP, FACG

Clinical Professor, Departments of Family Medicine and Urology/Director of Medical Student Education and Clerkship Director, Department of Family Medicine/Director, Patients and Populations Branch, University of Michigan Medical School, Ann Arbor

Dr. Heidelbaugh is a family physician who has 17 years of teaching experience. His specialty topics include gastrointestinal disorders, men's health, and primary care urology. He is a member of the American Gastroenterological Association guideline panels for irritable bowel syndrome, inflammatory bowel disease, and Lynch syndrome. He is the co editor and co author of the textbook ROME IV: Functional Gastrointestinal Disorders for Primary Care and Non GI Clinicians, published through the Rome Foundation. In addition, he is the consulting editor of Primary Care: Clinics in Office Practice and the president elect of the American Society for Men's Health. Dr. Heidelbaugh believes that increasing awareness and education about gastrointestinal and men's health issues is an important trend in medical education, clinical practice, and research.

FMX

## Learning Objectives

1. Apply evidence-based diagnostic criteria to evaluate patients presenting with recurrent or episodic abdominal pain for IBS.
2. Establish referral and follow-up protocol with a gastroenterologist for patients exhibiting red flags for other disease, for which endoscopic evaluation should be considered.
3. Develop treatment plans that involve positive patient-physician communication, shared decision making, and follow-up strategies that result in symptom relief and improved quality of life.

FMX

## Audience Engagement System

The image displays three sequential screenshots of a mobile application interface for CME events.   
Step 1: A dashboard with a grid of icons representing different CME categories.   
Step 2: A list of CME events with columns for Date, Title, and Status. One event is highlighted with a red arrow.   
Step 3: A detailed view of a selected CME event, including its title, location, date, and duration. A red arrow points to a 'CME Report / Evaluation' button.

FMX

## Epidemiology, Burden of Illness and Misconceptions

FMX

## High Prevalence and Incidence

- Common in western Europe and North America; less common in Asia
- Estimated at 11% worldwide, 12% in US
- Patients often present between age 30 – 50 years of age
- Decrease in reporting in older patients
  - (Are abnormal stools really abnormal?)
- 1.5 times more common in women than in men, commonly overlooked in men
- Average 14 hours lost work productivity per 40-hour week
- Graded decrease in prevalence with increasing income
  - 8 - 16% in those making less than \$20,000/year
  - 3 - 5% in those making more than \$75,000/year
- 2nd only to GERD in burden of GI illness
  - \$1.6 billion – direct costs
  - \$19.2 billion – indirect costs

Canavan C, et al. *Clin Epidemiol.* 2014;6:71-80; Hungin AP, et al. *Aliment Pharmacol Ther* 2005;21:1365-1375.

FMX

## IBS Myths and Misconceptions

- Misconceptions
  - Purely a psychological-based diagnosis
  - Unknown mechanism(s)
  - Difficult to diagnose/"diagnosis of exclusion"
- Reality
  - IBS is a REAL disorder and multifactorial
  - Majority of patients do not have psychological comorbidities!
  - Serotonin implicated in pathogenesis
  - Diagnosis can be made accurately in primary care!

FMX

## AES Question #1

Which of the following are common patient perceptions regarding IBS?

1. IBS causes colon cancer
2. If you change your diet, you will cure IBS
3. IBS and IBD are related
4. IBS does not lead to malnutrition
5. IBS will stay the same regardless of age

FMX

## Patient Perceptions of IBS

| Common Knowledge  | Common Misconceptions  | Most Desired Information  |
|---|--|---|
| Combination of abdominal pain and constipation<br>• And/or diarrhea<br>• And/or bloating  | Can develop into:<br>• Colitis<br>• Surgical problem<br>• Malnutrition<br>• Cancer | • What foods to avoid?<br>• What causes IBS?<br>• Coping strategies to reduce symptoms? |
| Triggers include stress at work, relationships, combinations of other (emotional) factors | Will worsen with age   |   |

Halpert A, et al. *Am J Gastroenterol.* 2007;102:1972-1982.

FMX

## Primary Care Perceptions

- PCPs were less likely than gastroenterologists to believe that IBS was related to prior physical or sexual abuse, previous infection, or learned behavior, but were more likely to state that diet caused IBS, or due to a linkable genetic etiology<sup>1</sup>
- A study found that PCPs in the Netherlands considered smoking, caffeine, diet, "hasty lifestyle," and lack of exercise as potential triggers for IBS symptoms, while PCPs in the UK considered food, infection, and travel as other possible triggers<sup>2</sup>

1. Lacy BE, et al. *Scand J Gastroenterology.* 2006;41:892-902.  
2. Casiday RE, et al. *Fam Pract.* 2009;26:34-39.

FMX

## Health-Related Quality of Life

- Patients with IBS have same physical HRQOL scores as patients with diabetes, and lower physical HRQOL scores than patients with depression and GERD<sup>1,2</sup>
- Psychological HRQOL scores are lower than patients with chronic renal failure, and can be so severe as to raise risk of suicidal behavior<sup>3</sup>

1. El-Serag HB, et al. *Aliment Pharmacol Ther.* 2002;16:1171-1185.  
 2. Gralnek IM, et al. *Gastroenterology.* 2000;119:654-660.  
 3. Spiegel BMR, et al. *Aliment Pharmacol Ther.* 2007;26:183-193.


FMX

## Health-Related Quality of Life


- IBS has a significant negative impact on HRQOL in most patients
- Failure to recognize this impact could undermine the physician-patient relationship and lead to dissatisfaction with care provided
- Imperative that PCPs:
  - Identify the predominant IBS symptom (C or D or M?)
  - Gauge symptom severity
  - Understand the negative impact on HRQOL, especially relative to the psychological impact

FMX

## Pathophysiology of IBS



## Impact on Quality of Life



*Functional Bowel Disorder Severity Index. IBS Patients: Their Illness, Experience, and Unmet Needs. www.aboutibs.org/pdfs/IBSpatients.pdf. Accessed March 29, 2010.*

## AES Question #2

Which of the following conditions is associated with IBS?

1. Coronary artery disease
2. Seasonal allergies
3. Interstitial cystitis
4. Rheumatoid arthritis
5. Hidradenitis suppurativa


FMX

## Other Associated Conditions

- GI motility disorders
  - Dyspepsia, GERD, cyclic vomiting, gastroparesis, etc.
- Psychiatric disorders
  - Anxiety, depression, somatoform disorders, PTSD
- Chronic back pain
- Fibromyalgia, chronic fatigue syndrome
- Chronic headaches, "migraines"
- Chronic pelvic pain – men and women
- Functional urinary symptoms (e.g., interstitial cystitis)
- Dysmenorrhea
- Sexual dysfunction

FMX

## IBS and Symptom Overlap



Frissora CL, Koch KL. *Curr Gastroenterol Rep.* 2005;7:264-271.

## Our Practices

- Most cases of IBS are diagnosed by gastroenterology
- Most research is done by gastroenterology
- Most patients with all forms of IBS are managed by primary care!
- Imperative that primary care has optimal understanding of perceptions and evidence behind diagnosis and treatment!

FMX

## Diagnosis of Functional GI Disorders

FMX

## Making the Diagnosis of IBS

- Take a careful history
- Look for warning signs
- Perform a thorough exam
- Use the Rome IV criteria
- Classify into the appropriate subtype
- Perform limited diagnostic tests


FMX

## Key Points in the History

- Onset of symptoms (chronicity)
- Is abdominal pain present?
- Is constipation or diarrhea present?
- Other GI symptoms (think overlap)
- Presence of common non-GI symptoms
- Prior tests?
- Prior treatments?

FMX

## Rome IV Criteria for IBS



Mearin F, Lacy BE, et al. *Gastroenterology.* 2016;150:1393-1407.

## Rome IV: Limited Diagnostic Tests

- In the appropriate patient, consider
  - CBC, ESR or CRP, fecal calprotectin
  - Celiac serologies
- All patients do not require testing
- No role for colonoscopy in all patients

Mearin F, Lacy BE, et al. *Gastroenterology*. 2016;150:1393–1407.


FMX

## IBS: The Value of the Physical Exam

- Identify other causes of symptoms
- Confirmation that complaints are being taken seriously
- Reassurance of absence of concerning physical exam findings


FMX

## IBS Subtypes Are Based on Stool Consistency




Adapted from: Longstreth GF, et al. *Gastroenterology*. 2006;130:1480-1491.

## Stool Consistency but not Frequency Predicts Colon Transit Time Bristol Stool Form Scale



## IBS-C vs. Chronic Idiopathic Constipation



FMX

## AES Question #3

Which of the following are considered "alarm features" for chronic constipation?

1. Alternating diarrhea and constipation
2. Celiac disease
3. Worsening GERD
4. New onset constipation in the elderly
5. Macrocytic anemia


FMX

## Alarm Features for Chronic Constipation

- Age > 50 years; > 45 years if African-American
- New onset constipation in elderly
- Severe symptoms not investigated
- Rectal bleeding
- Weight loss
- Family history of organic GI disease
- Palpable abdominal/rectal mass

**Investigate and treat appropriately; colonoscopy may be indicated**

Paré P, et al. *Can J Gastroenterol.* 2007;21(suppl B):3B-22B.



## Pretest Probability of Organic Disease<sup>1</sup>

| Organic Disease                         | IBS Patients (%) | Control/Population (%) |
|---|------------------|------------------------|
| Colitis/IBD                             | 0.51-0.98        | 0.3-1.2                |
| Colorectal cancer                       | 0-0.51           | 0-6 (varies with age)  |
| Lactose malabsorption                   | 38               | 26                     |
| Thyroid dysfunction                     | 4.2              | 5-9                    |
| Celiac disease                          | 3.6              | 0.7                    |
| Celiac disease: antibodies <sup>2</sup> | 7.3              | 4.8                    |
| Celiac disease: confirmed <sup>2</sup>  | 0.41             | 0.44                   |

1. American College of Gastroenterology Task Force on Irritable Bowel Syndrome, et al. *Am J Gastroenterol.* 2009;104(suppl 1):S1-S35.  
 2. Cash BD, et al. *Gastroenterology.* 2011;141:1187-1193.

## When to Refer to a Specialist?

- Warning Signs or Alarm Features
  - New onset of symptoms over age 50
  - Rectal bleeding/bloody stools
  - Unexplained weight loss
  - Family history of colon cancer or inflammatory bowel disease
  - Abnormal physical examination
- Severe or progressive symptoms
- Failure to improve with fiber or laxative therapies

FMX

## Other Tests for Chronic Constipation

For patients with severe symptoms or poor response to laxatives consider:

- **Balloon expulsion:** Suspected outlet problems/dyssynergia
- **Anorectal manometry:** Suspected dyssynergia; Hirschsprung disease
- **Defecography:** Suspected outlet problems/pelvic floor dysfunction
- **Colonic transit testing:** Sitz markers or wireless pH-motility capsule

|                              | Balloon Expulsion Test |                        |
|------------------------------|------------------------|------------------------|
|                              | Normal                 | Abnormal               |
| Normal Anorectal Manometry   | No outlet dysfunction  | Outlet dysfunction     |
| Abnormal Anorectal Manometry | No outlet dysfunction  | Dyssynergic Defecation |

*Cash BD, et al. Rev Gastroenterol Disord. 2007;7:116-133.*

## Treatment of Constipation- and Diarrhea-Related Functional GI Disorders

FMX

## Improving Patient-Physician Interactions

- Listen (it's hard - don't interrupt!!!)
- Understand fears
- Review goals
- Set expectations
  - Short-term
  - Long-term
- Develop a logical treatment plan together
- Arrange for routine follow-up

FMX

## Managing IBS: What Do Your Patients Want?

- They want you to listen
  - Understand their history (symptoms, work, home)
- Education about their condition
  - Address questions or concerns
  - Address uncertainty of IBS
- Reassurance
- A positive diagnosis
  - Review diagnostic criteria and results with patients
- Symptom improvement

FMX

## Graded Treatment of Chronic Constipation




## IBS & Low FODMAP Diet

### Fermentable Oligo-, Di-, Monosaccharides And Polyols


|  |                        |   |
|--|------------------------|---|
|  | <b>Excess Fructose</b> | Honey, apples, pears, peaches, mangos, fruit juice, dried fruit       |
|  | <b>Fructans</b>        | Wheat (large amounts), rye (large amounts), onions, leeks, zucchini   |
|  | <b>Sorbitol</b>        | Apricots, peaches, artificial sweeteners, artificially sweetened gums |
|  | <b>Raffinose</b>       | Lentils, cabbage, Brussels sprouts, asparagus, green beans, legumes   |

## Improvements in IBS Symptom Scores: Low FODMAP vs Control Diet



FMX



## IBS & Low-Gluten Diet



Quality of evidence: very low.  
VAS, visual analog scale.  
Biesiekierski JR, et al. *Am J Gastroenterol*. 2011;106:508-514.

● Gluten  
● Placebo  
VAS Score 0 = None 100 = Worst

FMX



### Practical Advice on Fiber and Antispasmodics for IBS-C

**Fiber supplementation**

- Mild constipation symptoms
- May or may not help with pain
- Soluble probably better than insoluble
  - "Start low and titrate slow"
  - Gas and bloating are the main side effects

**Anti-spasmodics**

- For postprandial abdominal pain
- Not effective for chronic abdominal pain
- Avoid in the elderly

Eswaran S, et al. *Am J Gastroenterol.* 2013;108:718-727.

### Fiber and Stool Softeners for IBS-C


- Fiber and stool softeners (docusate) are most useful in patients with mild, infrequent constipation
- Best evidence for psyllium up to 25 grams/day
- Their role in patients with significantly delayed colon transit or dyssynergia is limited
- Fiber may worsen symptoms in patients with significantly delayed colon transit or dyssynergic defecation

### AES Question #4

Which of the following novel mechanisms has shown promise in the pharmacologic treatment of IBS?


1. Chloride channel activation
2. Sodium channel activation
3. Potassium channel activation
4. Magnesium channel activation

FMX






## Lubiprostone for IBS-C: Data From 2 Phase III Trials




Drossman DA, et al. *Aliment Pharmacol Ther.* 2009;29(3):329-341.

## Effects of Guanylate Cyclase-C Receptor Activation



Brierley SM. *Curr Opin Pharmacol.* 2012;12:632-640.  
Eutamene H, et al. *Neurogastroenterol Motil.* 2010;22:312-e84.  
Comiskey S, et al. *Am J Gastroenterol.* 2012;107 (suppl 1):S700 (abs 1725).

## Linaclootide Phase 3 IBS-C Trial



ITT Population, Observed Cases, LS-Means presented, P-values based on ANCOVA at each week. Bars represent 95% confidence intervals.

Chey WD, et al. *Am J Gastroenterol.* 2012;107:1702-1712.

## Linaclootide Phase 3 IBS-C Trial

TEAEs Occurring in  $\geq 3\%$  in Lin 266 µg Group and Lin 266 µg > Placebo

|                       | Weeks 1-12       |                     | Weeks 1-26       |                     |
|-----------------------|------------------|---------------------|------------------|---------------------|
|                       | Placebo<br>N=403 | Lin 266 µg<br>N=402 | Placebo<br>N=403 | Lin 266 µg<br>N=402 |
| Any TEAE              | 187 (46%)        | 212 (53%)           | 228 (57%)        | 263 (65%)           |
| Diarrhea              | 7 (2%)           | 71 (18%)            | 10 (2%)          | 79 (20%)            |
| Nausea                | 17 (4%)          | 17 (4%)             | 24 (6%)          | 23 (6%)             |
| URI                   | 13 (3%)          | 14 (3%)             | 22 (5%)          | 22 (5%)             |
| Abdominal pain        | 14 (3%)          | 15 (4%)             | 16 (4%)          | 18 (4%)             |
| Flatulence            | 7 (2%)           | 13 (3%)             | 9 (2%)           | 15 (4%)             |
| Gastroenteritis viral | 4 (1%)           | 8 (2%)              | 9 (2%)           | 15 (4%)             |
| Headache              | 8 (2%)           | 13 (3%)             | 11 (3%)          | 13 (3%)             |

TEAE: treatment emergent adverse events  
Chey WD, et al. *Am J Gastroenterol.* 2012;107:1702-1712.

## Loperamide\* for IBS-D

- Low doses (2 mg once or twice daily) may be effective to decrease stool frequency and improve stool consistency<sup>1</sup>
- 2 randomized controlled trials in IBS (N=42) show efficacy for diarrhea<sup>2,3</sup>
- No impact on symptoms of abdominal discomfort, bloating, or global IBS<sup>2,3</sup>
- Adverse effects: dizziness, abdominal pain/bloating, constipation, dry mouth, fatigue<sup>1</sup>

\*FDA approved for diarrhea, but not for IBS-D.  
1. US FDA, CDER. Loperamide NDA 017694. Label 10/21/2005. 2. Lavé B et al. *Scand J Gastroenterol Suppl.* 1987;130:77-80. 3. Hovdenak N. *Scand J Gastroenterol Suppl.* 1987;130:81-84.

FMX

## Diphenoxylate-Atropine\* for IBS-D

- A synthetic opiate agonist with a structure similar to meperidine
- Atropine is added to discourage deliberate abuse or overdose of diphenoxylate
- Schedule V controlled substance
- First approved by FDA in 1960
- No prospective studies

\*FDA approved for diarrhea, but not IBS-D.  
Lacy BE, et al. *Ther Adv Gastroenterol.* 2009;2:221-238.

FMX

## Antispasmodics for IBS-D

- 22 randomized controlled trials comparing 12 different antispasmodics vs placebo (N=1778 patients)
- Significant heterogeneity among studies
- Many agents not available in US
- Appear most useful for abdominal pain
- In meta-analysis, symptoms persist in 39% of patients receiving antispasmodics vs 56% of placebo-treated patients (RR: 0.68; 95% CI: 0.57-0.81)

Ford AC, et al. *BMJ*. 2008;337:a2313.

FMX

## Bile Acid Sequestrants\* for IBS-D

- 20%-30% of patients with IBS-D or functional diarrhea have bile acid malabsorption (BAM)<sup>1</sup>
- Up to 10% of IBS-D patients have evidence of severe BAM<sup>1</sup>
- Bile acids accelerate colonic transit, increase stool frequency, and reduce consistency<sup>1</sup>
- Best test: <sup>75</sup>SeHCAT<sup>1</sup>
- Cholestyramine, colessevelam hydrochloride may improve sx<sup>1,2</sup>

\*Off-label use, not FDA approved for IBS-D.

1. Wedlake L, et al. *Aliment Pharmacol Ther*. 2009;30:707-717.

2. Wong BS, et al. *Dig Dis Sci*. 2012;57:1222-1226.

FMX

## Alosetron for IBS-D

- A 5-HT<sub>2</sub> antagonist
- Reduces stool frequency and abdominal pain; improves urgency
- Treatment population: women with chronic, severe IBS-D who have failed other treatments
  - Dose: 0.5-1.0 mg QD to BID
- 8 large, R, DB, PC trials have shown benefits for global and individual symptoms of IBS (NNT = 8)
- Patient education regarding possible serious adverse effects
  - 0.95 cases of ischemic colitis/1000 patient-years
  - 0.36 cases of severe constipation/1000 patient-years
- If ischemic colitis occurs, it is usually within the first month
- Prescribing program mandated by FDA


Ford AC, et al. *Am J Gastroenterol*. 2014;109(Suppl 1):S2-S26.

FDA, Alosetron REMS.

<http://www.fda.gov/downloads/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/UCM227960.pdf>

## Treatment With Rifaximin - IBS-D

IBS-Related Abdominal Pain and Stool Consistency (Worst-Case Analysis)



Responder: Patient responding to IBS-related abdominal pain (>30% improvement) and stool consistency (>50% decrease in number of BMs with type 6 or 7) from baseline for ≥2 of the 4 weeks.

BM, bowel movement.

Lacy BE, et al. *Gastroenterology*. 2015;148(suppl 1):S657.


## Eluxadoline for IBS-D

- Mu (μ) opioid receptor agonist
- Delta (δ) opioid receptor antagonist
- Low systemic absorption and bioavailability
  - Low potential for drug-drug interactions
- Common adverse events include nausea and constipation
- No signs of dependence or withdrawal during clinical trials

FMX

## Eluxadoline for IBS-D

Primary Endpoint: Composite Responders – Pooled Data



Lembo A, et al. *N Engl J Med*. 2016;374:242-253.


FMX

## Complementary and Alternative Treatments for Functional GI Disorders

FMX

## Probiotics: Putative Mechanisms of Action


- Competitive inhibition
- Barrier protection
- Immune effects
- Anti-inflammatory effects
- Production of various substances (enzymes, SCFA, bacteriocidal agents)
- Ability to alter local pH and physiology
- Provides nutrition to colonocytes



SCFA: short chain fatty acids  
Camilleri M. *J Clin Gastroenterol.* 2006;40:264-269.

FMX

## *Bifidobacteria Infantis* 35624 for IBS Global Assessment of Relief



SQA: (Subjects' Global Assessment) a yes/no response to the following question:  
"Please consider how you felt in the past week in regard to your IBS, in particular your general well being, and symptoms of abdominal discomfort or pain, bloating, or distension, and altered bowel habit. Compared with the way you felt before beginning the medication, have you had adequate relief of your IBS symptoms?"

Whorwell PJ, et al. *Am J Gastroenterol.* 2006;101:1581-1590.

FMX

## Randomized Trial of Dried Plums for Chronic Constipation

- Dried plums contain large amounts of fiber and sorbitol
- 8-week RCT single-blinded cross-over study examined treatment with dried plums (prunes, 50 g/day or 6 prunes) compared with psyllium (11 g/day fiber) taken BID
  - 40 patients with chronic constipation
  - Dried plums resulted in a greater improvement in constipation symptoms compared with psyllium
    - More CSBMs (3.5 ± 0.2 vs 2.8 ± 0.2, P = 0.006)
    - Softer stools (3.2 vs 2.8 on the BSFS, P = 0.02)
    - Overall constipation symptoms (both groups improved, NS)



Attaluri A, et al. *Aliment Pharmacol Ther.* 2011;33(7):822-828.

FMX

## Current Therapeutic Guidelines: American Gastroenterological Association

FMX

## AGA Guideline for Pharmacologic Treatment of IBS

- Multidisciplinary guideline development
  - Gastroenterologists
  - Epidemiologists
  - Primary Care
  - Patient Representative
  - Rigorous COI exclusions (*NB: No one is paid!*)
- Clinical Practice and Quality Measures Committee
  - Guideline
  - Technical Review

FMX

## AGA Guideline for Pharmacologic Treatment of IBS

- **GRADE Methodology**
  - Grading of **R**ecommendations **A**ssessment, **D**evelopment, and **E**valuation System
  - High/Moderate/Low **Q**uality Evidence
  - Strong/Conditional (Weak) **S**trength of Evidence
- **PICO**
  - Patient **P**roblem or **P**opulation
  - **I**ntervention
  - **C**omparison
  - **O**utcome(s)

FMX

## AGA Guideline for Pharmacologic Treatment of IBS

1. Use **linaclotide** (over no drug tx) in IBS-C<sup>1</sup>
  - 1. *Strong recommendation; high quality evidence*
2. Use **lubiprostone** (over no drug tx) in IBS-C<sup>1</sup>
  - 1. *Conditional recommendation; moderate quality evidence*
3. Use **laxatives** [PEG] (over no drug tx) in IBS-C
  - 1. *Conditional recommendation; low quality evidence*
4. Use **rifaximin** (over no drug tx) in IBS-D<sup>2</sup>
  - 1. *Conditional recommendation; moderate quality evidence*
  - 1. <sup>1</sup> PIs who value avoiding higher out-of-pocket costs may prefer alternate treatments
  - 2. <sup>2</sup> Not approved by FDA, cost may be high, no evidence to support repetitive treatment

Weinberg DS, Smalley W, Heidelbaugh JJ, Sultan S. American Gastroenterological Association Institute Guideline on the Pharmacological Management of Irritable Bowel Syndrome. *Gastroenterology* 2014;147(5):1146-1148.

## AGA Guideline for Pharmacologic Treatment of IBS

5. Use **alosetron** (over no drug tx) in IBS-D
  - *Conditional recommendation; moderate evidence*
6. Use **loperamide** (over no drug tx) in IBS-D
  - *Conditional recommendation; very low quality evidence*
7. Use **TCAs** (over no drug tx) in IBS
  - *Conditional recommendation; low quality evidence*
8. Do not use **SSRIs** for IBS patients
  - *Conditional recommendation; low quality evidence*
9. Use **antispasmodics** (over no drug tx) in IBS
  - *Conditional recommendation; low quality evidence*

Weinberg DS, Smalley W, Heidelbaugh JJ, Sultan S. American Gastroenterological Association Institute Guideline on the Pharmacological Management of Irritable Bowel Syndrome. *Gastroenterology* 2014;147(5):1146-1148.

## Practice Recommendations

- Constipation is a multi-symptom condition that must be appropriately defined
- The main causes of constipation are slow colon transit and/or disordered defecation - more often functional than organic in etiology
- **Diet and lifestyle changes** help with most mild, moderate, or intermittent constipation symptoms, and should always be tried first
- Laxatives including osmotics, stimulants, and prosecretory agents improve symptoms in many patients
- Biofeedback and physical therapy are the preferred treatments for dyssynergic defecation and have modest results
- When patients fail to respond to laxatives, diagnostic testing can help to determine the etiology of constipation symptoms
- Gain skill in using newer agents, know when to refer to gastroenterology
- A multi-disciplinary approach is optimal for severely affected patients

FMX

## Questions



FMX

## Contact Information

Joel J. Heidelbaugh, MD, FAAFP, FACP  
[jheidel@umich.edu](mailto:jheidel@umich.edu)

FMX