



Body System: Geriatrics		
Session Topic: Polypharmacy in the Elderly		
Educational Format		Faculty Expertise Required
REQUIRED	Interactive Lecture	Expertise in the field of study. Experience teaching in the field of study is desired. Preferred experience with audience response systems (ARS). Utilizing polling questions and engaging the learners in Q&A during the final 15 minutes of the session are required.
OPTIONAL	Problem-Based Learning (PBL)	Expertise teaching highly interactive, small group learning environments. Case-based, with experience developing and teaching case scenarios for simulation labs preferred. Other workshop-oriented designs may be accommodated. A typical PBL room is set for 50-100 participants, with 7-8 each per round table. <u>Please describe your interest and plan for teaching a PBL on your proposal form.</u>
Professional Practice Gap	Learning Objective(s) that will close the gap and meet the need	Outcome Being Measured
<ul style="list-style-type: none"> Family physicians have gaps in knowledge and performance in evaluating for potentially adverse drug events, among elderly patients receiving multiple medications. Family physicians have gaps in knowledge and performance in developing systematic approach to managing elderly patients with multiple chronic conditions that focuses on the quality-of-life outcomes most valued by the patient, and includes applicable REMS. Family physicians have knowledge and performance gaps in developing collaborative care plans to address the needs of patients who have poor health literacy or reduced cognitive function, or those patients who have language barriers, in order to foster appropriate self- 	<ol style="list-style-type: none"> Use evidence-based criteria (e.g. BEERS, STOPP, START) to evaluate for potentially adverse drug events, among elderly patients receiving multiple medications. Develop a systematic approach, including applicable REMS, to managing elderly patients with multiple chronic conditions that focus on the quality-of-life outcomes most valued by the patient. Develop collaborative care plans to address the needs of patients who have poor health literacy or reduced cognitive function, or those patients who have language barriers, in order to foster appropriate self-administration of medications. Counsel elderly patients and caregivers about tools, resources, and strategies to aid in the self-administration of medications. 	Learners will submit written commitment to change statements on the session evaluation, indicating how they plan to implement presented practice recommendations.



<p>administration of medications.</p> <ul style="list-style-type: none"> Family physicians have knowledge and performance gaps in educating elderly patients and caregivers about tools, resources, and strategies to aid in the self-administration of medications. Family physicians have gaps in knowledge in balancing achieving care guidelines and reducing polypharmacy. 		
<p>ACGME Core Competencies Addressed (select all that apply)</p>		
X	Medical Knowledge	Patient Care
X	Interpersonal and Communication Skills	Practice-Based Learning and Improvement
	Professionalism	Systems-Based Practice
<p>Faculty Instructional Goals</p>		
<p>Faculty play a vital role in assisting the AAFP to achieve its mission by providing high-quality, innovative education for physicians, residents and medical students that will encompass the art, science, evidence and socio-economics of family medicine and to support the pursuit of lifelong learning. By achieving the instructional goals provided, faculty will facilitate the application of new knowledge and skills gained by learners to practice, so that they may optimize care provided to their patients.</p> <ul style="list-style-type: none"> Provide up to 3 evidence-based recommended practice changes that can be immediately implemented, at the conclusion of the session; including SORT taxonomy & reference citations Facilitate learner engagement during the session Address related practice barriers to foster optimal patient management Provide recommended journal resources and tools, during the session, from the American Family Physician (AFP), Family Practice Management (FPM), and Familydoctor.org patient resources; those listed in the <u>References</u> section below are a good place to start <ul style="list-style-type: none"> Visit http://www.aafp.org/journals for additional resources Visit http://familydoctor.org for patient education and resources Provide updates on new treatment therapies, changes to therapies, or warnings associated with existing therapies. Provide recommendations regarding new FDA approved medications; including safety, efficacy, tolerance, and cost considerations relative to currently available options. Include relevant FDA REMS education for any applicable medications. Provide specific examples utilizing BEERS/STOPP/START criteria to evaluate for potentially adverse drug events, including examples of when physicians should not rely on such criteria alone 		



- Provide specific examples and strategies for systematically managing elderly patients with chronic conditions, including applicable REMS, with an emphasis on reducing adverse drug events
- Provide specific strategies, tools, and resources to address the needs of patients who have poor health literacy or reduced cognitive function, or those patients who have language barriers, in order to foster appropriate self-administration of medications. These strategies should provide an approach to comply with the most meaningful guidelines for the elderly with appropriate medications and reduction of polypharmacy.
- Provide specific strategies and resources to assist physicians in providing consultation to elderly patients and caregivers about tools, resources, and strategies to aid in the self-administration of medications
- Provide recommendations regarding guidelines for Medicare reimbursement.
- Provide recommendations to maximize office efficiency and guideline adherence to the management of medications for elderly patients.
- Provide an overview of newly available treatments, including efficacy, safety, contraindications, and cost/benefit relative to existing treatments.
- Provide instructions regarding the incorporation and use of the PCMH/ACO/Primary Care Core Measure Set into practice.

Needs Assessment:

Among older Americans, aged 60 and over, more than 76% used two or more prescription drugs and 37% used five or more.¹ The risk of adverse drug events is considerable in older patients, leading to one in six hospital admissions because of an adverse drug event.² In fact, polypharmacy has been shown to increase a patient's risk for falls and postoperative hip fracture, precipitate confusion, and incontinence.³⁻⁵ Polypharmacy associated with multi-morbidity is burdensome for patients, likely leads to a reduction of overall drug benefit, and is an additive effect of harms and side effects.⁶ Older adults are often negatively impacted by polypharmacy and are at greater risk for adverse drug events due to metabolic changes, drug-drug interactions, prescribing cascades, and are at greater risk for hip fractures.^{2,7} Unintentional weight loss in older adults may also be due to medication use as polypharmacy can interfere with taste or cause nausea.⁸

The American Academy of Family Physicians (AAFP) CME Needs Assessment Survey indicates that family physicians have statistically significant and meaningful knowledge gaps to provide optimal manage of polypharmacy and medication interactions.^{9,10} More specifically, CME outcomes data from 2014-2017 AAFP FMX (formerly Assembly): *Polypharmacy* sessions suggest that physicians have knowledge and practice gaps with regard to using medication reconciliation tools; BEERS, START and STOP medication lists; and reviewing medication lists with patients; establishing protocols to minimize adverse drug events for hospitalized patients, and patients being discharged from the hospital; and provide counseling regarding side effects and stopping unnecessary medications.¹¹⁻¹⁴

Physicians are often challenged with the need to manage multiple chronic conditions in their older patients because clinical practice guidelines, applied without consideration of other chronic



conditions and medications, can lead to adverse drug events.¹⁵ Additionally, physicians are often reluctant to stop medications, especially if they did not initiate the treatment and the patient seems to be tolerating the therapy. The decision to discontinue medication is determined in part by the goals of care for that patient and the risks of adverse effects for that patient; and targets for treatment, based on outcomes evidence from studies in younger patients, may not be appropriate for older adults.¹⁵ Instead, physicians must be able to develop a systematic approach that identifies the quality-of-life outcomes most valued by the patient that should take precedence over the routine implementation of practice guidelines that recommend medications for generic clinical scenarios.^{2,16} A recent Cochrane review indicated that Warfarin (Coumadin), insulin, and digoxin accounted for one in three of ER visits, by older patients, because of an adverse drug event, whereas drugs on the Beers list accounted for less than 9 percent. Interventions to improve appropriate polypharmacy have been shown to reduce inappropriate prescribing and medication-related problems, although it is unclear if these intervention improve appropriate polypharmacy, such as pharmaceutical care.¹⁷ The rate of adverse drug events may be reduced by using validated risk calculators for bleeding in patients taking warfarin, setting less stringent goals for A1C levels in older patients with comorbidities, and avoiding high doses of digoxin or use of the drug without proper indications.¹⁸ Recent studies show that 42% of older adults received potentially inappropriate medications (PIM) but that a portion of them were not used for particularly long or were used for individuals who lacked specific medical conditions targeted in the Beers criteria as problematic or who had diagnoses for which use was justified. Even with these qualifications in the definition, 30.9% of community-dwelling older adults were prescribed PIMs, some of which are known to be associated with falls, delirium, declines in cognitive and physical functioning, and other potentially serious health outcomes.¹⁹

Family physicians should be knowledgeable about factors that influence pharmacokinetics in older adults; be aware of sources of adverse drug events; medications to avoid when possible in older adults; have a systematic approach for detecting adverse drug events; be aware of practical considerations to reduce the risk of adverse drug events; and know how to effectively use BEERS, STOPP, and START criteria.^{2,7,19-21} Family physicians should also be familiar with applicable REMS for medications they are prescribing to their elderly patients.²²

Elderly patients also have difficulty with self-administration of medication. Some studies suggest that as many as half of the elderly population lacks the adequate literacy skills to self-administer medications, particularly for multiple chronic conditions.^{23,24} Physicians should consider the following evidence based recommendations to address concerns of health literacy among elderly patients:²⁵

- Use universal health literacy precautions with all patients, regardless of their literacy or education levels.
- Prioritize and limit information to three key points for each visit.
- Use the teach-back method to assess patient comprehension of information.
- Simplify forms and offer assistance with form completion.

However, non-adherence is not simply a knowledge discrepancy, but it can also involve feelings, reactions to the physician, cost, availability, and competing medical belief systems.⁷ Physicians should have strategies in place, including group counseling and collaborative care plans, to address the needs of patients who have poor health literacy, or those patients who have language



barriers, in order to foster appropriate self-administration of medications.²⁶⁻³¹ Elderly patients who do not live in some level of managed care facility will often receive assistance from a family caregiver. In fact, in 2009, nearly 66 million Americans provided some level of care for an elderly family member.³² However, a majority of caregivers (81%) feel inadequately trained, having never received any formal education in caregiving.³² Family physicians should be prepared to counsel both elderly patients and caregivers about tools that can help manage self-administration of medications, such as home telemedicine, telehealth, disease-state monitoring systems, automatic medication dispenser, and vibrating alarm watches.³²

Physicians should consider the National Action Plan for ADE Prevention from the Office of Disease Prevention and Health Promotion (ODPHP), on behalf of the Federal Interagency Steering Committee for Adverse Drug Events.³³ Additionally, physicians may want to consider the following evidence-based recommendations:²

- The use of antipsychotics, antidepressants, benzodiazepines, and sedative/hypnotics in older adults should be limited to reduce the risk of falls.
- The number of medications in older adults should be limited, because each new medication adds more than one adverse drug event each year and taking six or more medications increases this risk fourfold.
- Ideally, the primary care physician should prescribe all medications, because each additional prescriber increases the risk of adverse drug events in older adults by 30 percent.
- Patients with several chronic health conditions should be asked often about adverse drug events, because the odds of an event double for four or five conditions and triple for six or more.

Best Practice Recommendations from the Choosing Wisely® campaign:³⁴

- Do not prescribe a medication without conducting a drug regimen review. (American Geriatrics Society)
- Avoid using prescription appetite stimulants or high-calorie supplements for the treatment of anorexia or cachexia in older adults; instead, optimize social supports, discontinue medications that may interfere with eating, provide appealing food and feeding assistance, and clarify patient goals and expectations. (American Geriatrics Society)

The ADE Action Plan addresses a defined group of ADEs that are considered to be common, clinically significant, preventable, and measurable; resulting from high-priority drug classes; and occurring largely in high-risk populations.

Three key drug classes identified as initial targets for the ADE Action Plan include:

1. **Anticoagulants** (primary ADE of concern: bleeding)
2. **Diabetes agents** (primary ADE of concern: hypoglycemia)
3. **Opioids** (primary ADE of concern: accidental overdoses, over-sedation, respiratory depression)

To align the efforts of federal health agencies to reduce patient harms from these specific ADEs nationally, the ADE Action Plan identifies a four-pronged approach:

- **Surveillance** — Coordinate existing federal surveillance resources and data to assess the health burden and rates of ADEs.



- **Prevention** — Share existing evidence-based prevention tools across federal agencies and with non-federal health care providers and patients.
- **Incentives and Oversight** — Explore opportunities, including financial incentives and oversight authorities, to promote ADE prevention.
- **Research** — Identify current knowledge gaps and future research needs (unanswered questions) for ADE prevention.

Unfortunately, physicians are rarely taught formally during residency how to take people off medications, or deprescribing.³⁵

The American Academy of Family Physicians Academy has participated in the Core Measures Collaborative (the Collaborative) convened by America's Health Insurance Plans (AHIP) since August 2014. The Collaborative is a multi-stakeholder effort working to define core measure sets of various specialties promoting alignment and harmonization of measure use and collection across both public and private payers.

Participants in the Collaborative included Centers for Medicare and Medicaid Services (CMS), the National Quality Forum (NQF), private payers, provider organizations, employers, and patient and consumer groups. This effort exists to decrease physician burden by reducing variability in measure selection, specifications and implementation— making quality measurement more useful and meaningful for consumers, employers, as well as public and private clinicians.

With significant AAFP input, a PCMH/ACO/Primary Care Core Measure Set has been developed for primary care. The goal of this set is to decrease burden and allow for more congruence between payer reporting programs.³⁶

Resources: Evidence-Based Practice Recommendations/Guidelines/Performance Measures

- Reducing the risk of adverse drug events in older adults²
- Using medications appropriately in older adults⁵
- American Geriatric Society: Care of Older Adults with Multi-morbidity⁶
- American Geriatric Society: updated BEERS criteria²⁰
- Minimizing adverse drug events in older patients⁷
- Appropriate use of polypharmacy for older patients¹⁸
- National Action Plan for Adverse Drug Event Prevention³³
- Unintentional Weight Loss in Older Adults⁸
- Health literacy: the gap between physicians and patients²⁸
- Health Literacy in Primary Care Practice²⁵
- The role of health literacy in patient-physician communication²⁹
- Help your patient "get" what you just said: a health literacy guide³⁰
- Engaging Patients in Collaborative Care Plans³¹
- Medication adherence: we didn't ask and they didn't tell³⁷
- What Needs to Change to Make Deprescribing Doable³⁵
- Adding health education specialists to your practice³⁸



- Envisioning new roles for medical assistants: strategies from patient-centered medical homes³⁹
- The benefits of using care coordinators in primary care: a case study⁴⁰
- Health Coaching: Teaching Patients to Fish⁴¹
- FamilyDoctor.org - Seniors: Managing Your Medications (patient resource)⁴²
- Caregiver Resource – FCA: Caregivers' Guide to Medications and Aging⁴³
- FamilyDoctor.org. Caregiving: Caring for an Elderly Relative - Managing Medicines (patient resource)⁴⁴
- FamilyDoctor.org. Falls: How to Lower Your Risk (patient resource)⁴⁵

References

1. Centers for Disease Control and Prevention. NCHS Data Brief: Prescription Drug Use Continues to Increase: U.S. Prescription Drug Data for 2007-2008. 2010; <http://www.cdc.gov/nchs/data/databriefs/db42.htm>. Accessed July, 2013.
2. Pretorius RW, Gataric G, Swedlund SK, Miller JR. Reducing the risk of adverse drug events in older adults. *American family physician*. 2013;87(5):331-336.
3. Ziere G, Dieleman JP, Hofman A, Pols HA, van der Cammen TJ, Stricker BH. Polypharmacy and falls in the middle age and elderly population. *British journal of clinical pharmacology*. 2006;61(2):218-223.
4. Lai SW, Liao KF, Liao CC, Muo CH, Liu CS, Sung FC. Polypharmacy correlates with increased risk for hip fracture in the elderly: a population-based study. *Medicine*. 2010;89(5):295-299.
5. Willams CM. Using medications appropriately in older adults. *American family physician*. 2002;66(10):1917-1924.
6. American Geriatrics Society, Expert Panel on the Care of Older Adults with Multimorbidity. Guiding principles for the care of older adults with multimorbidity: an approach for clinicians. 2012; <http://www.guideline.gov/content.aspx?id=39322&search=polypharmacy>. Accessed 7/17/2013.
7. Pham CB, Dickman RL. Minimizing adverse drug events in older patients. *American family physician*. 2007;76(12):1837-1844.
8. Gaddey HL, Holder K. Unintentional weight loss in older adults. *American family physician*. 2014;89(9):718-722.
9. CME Training Session Needs Survey. In. Leawood KS: AAFP; 2016.
10. AAFP. 2012 CME Needs Assessment: Clinical Topics. In: American Academy of Family Physicians; 2012.
11. American Academy of Family Physicians (AAFP). AAFP FMX CME Outcomes Report. In. Leawood KS: AAFP; 2015.
12. American Academy of Family Physicians (AAFP). AAFP Assembly CME Outcomes Report. In. Leawood KS: AAFP; 2014.
13. American Academy of Family Physicians (AAFP). 2017 AAFP FMX CME Outcomes Report. In. Leawood KS: AAFP; 2017.
14. American Academy of Family Physicians (AAFP). AAFP FMX CME Outcomes Report. In. Leawood KS: AAFP; 2016.



15. Boyd CM, Darer J, Boulton C, Fried LP, Boulton L, Wu AW. Clinical practice guidelines and quality of care for older patients with multiple comorbid diseases: implications for pay for performance. *JAMA : the journal of the American Medical Association*. 2005;294(6):716-724.
16. Steinman MA, Hanlon JT. Managing medications in clinically complex elders: "There's got to be a happy medium". *JAMA : the journal of the American Medical Association*. 2010;304(14):1592-1601.
17. Patterson SM, Hughes C, Kerse N, Cardwell CR, Bradley MC. Interventions to improve the appropriate use of polypharmacy for older people. *Cochrane Database Syst Rev*. 2012;5:CD008165.
18. Hitzeman N, Belsky K. Appropriate use of polypharmacy for older patients. *American family physician*. 2013;87(7):483-484.
19. Davidoff AJ, Miller GE, Sarpong EM, Yang E, Brandt N, Fick DM. Prevalence of potentially inappropriate medication use in older adults using the 2012 Beers criteria. *J Am Geriatr Soc*. 2015;63(3):486-500.
20. By the American Geriatrics Society Beers Criteria Update Expert P. American Geriatrics Society 2015 Updated Beers Criteria for Potentially Inappropriate Medication Use in Older Adults. *J Am Geriatr Soc*. 2015;63(11):2227-2246.
21. Brown JD, Hutchison LC, Li C, Painter JT, Martin BC. Predictive Validity of the Beers and Screening Tool of Older Persons' Potentially Inappropriate Prescriptions (STOPP) Criteria to Detect Adverse Drug Events, Hospitalizations, and Emergency Department Visits in the United States. *J Am Geriatr Soc*. 2016;64(1):22-30.
22. U.S. Food and Drug Administration (FDA). Approved Risk Evaluation and Mitigation Strategies (REMS). 2013; <http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm111350.htm>. Accessed May, 2013.
23. Marvanova M, Roumie CL, Eden SK, Cawthon C, Schnipper JL, Kripalani S. Health literacy and medication understanding among hospitalized adults. *Journal of hospital medicine : an official publication of the Society of Hospital Medicine*. 2011;6(9):488-493.
24. Kripalani S, Henderson LE, Chiu EY, Robertson R, Kolm P, Jacobson TA. Predictors of medication self-management skill in a low-literacy population. *Journal of general internal medicine*. 2006;21(8):852-856.
25. Hersh L, Salzman B, Snyderman D. Health Literacy in Primary Care Practice. *American family physician*. 2015;92(2):118-124.
26. Anderson KM, Siems LV, Holloway SC, Sultana N, Braund WE, Harris LM. Group counselling improves quality for patients with limited health literacy. *Quality in primary care*. 2012;20(1):5-13.
27. Edison K, Jeanetta S, Staiculescu I. Practice gaps--providing appropriate patient education materials for non-English-speaking patients. *Archives of dermatology*. 2011;147(2):244.
28. Safer RS, Keenan J. Health literacy: the gap between physicians and patients. *American family physician*. 2005;72(3):463-468.
29. Williams MV, Davis T, Parker RM, Weiss BD. The role of health literacy in patient-physician communication. *Family medicine*. 2002;34(5):383-389.
30. Roett MA, Wessel L. Help your patient "get" what you just said: a health literacy guide. *The Journal of family practice*. 2012;61(4):190-196.



31. Mauksch L, Safford B. Engaging Patients in Collaborative Care Plans. *Family practice management*. 2013;20(3):35-39.
32. Collins LG, Swartz K. Caregiver care. *American family physician*. 2011;83(11):1309-1317.
33. The Office of Disease Prevention and Health Promotion (ODPHP). National Action Plan for ADE Prevention. 2014; <http://health.gov/hcq/ade-action-plan.asp>. Accessed June, 2016.
34. American Academy of Family Physicians (AAFP). Choosing Wisely. 2018; <https://www.aafp.org/afp/recommendations/search.htm>. Accessed June, 2018.
35. Mishori R. What Needs to Change to Make Deprescribing Doable. *Family practice management*. 2018;25(3):5-6.
36. American Academy of Family Physicians (AAFP). PCMH/ACO/Primary Care Core Measure Set. 2016; <http://www.aafp.org/practice-management/improvement/measures.html>. Accessed May, 2016.
37. Brown M, Sinsky CA. Medication adherence: we didn't ask and they didn't tell. *Family practice management*. 2013;20(2):25-30.
38. Chambliss ML, Lineberry S, Evans WM, Bibeau DL. Adding health education specialists to your practice. *Family practice management*. 2014;21(2):10-15.
39. Naughton D, Adelman AM, Bricker P, Miller-Day M, Gabbay R. Envisioning new roles for medical assistants: strategies from patient-centered medical homes. *Family practice management*. 2013;20(2):7-12.
40. Mullins A, Mooney J, Fowler R. The benefits of using care coordinators in primary care: a case study. *Family practice management*. 2013;20(6):18-21.
41. Ghorob A. Health Coaching: Teaching Patients to Fish. *Family practice management*. 2013;20(3):40-42.
42. FamilyDoctor.org. Seniors: Managing Your Medications. 2013; <http://familydoctor.org/familydoctor/en/seniors.html>. Accessed July, 2013.
43. Family Caregiver Alliance. Caregivers' Guide to Medications and Aging. 2013; http://www.caregiver.org/caregiver/jsp/content_node.jsp?nodeid=1104. Accessed July, 2013.
44. FamilyDoctor.org. Caregiving: Caring for an Elderly Relative - Managing Medicines. 2012; <http://familydoctor.org/familydoctor/en/seniors/caring-for-an-elderly-relative/managing-medicines.html>. Accessed August, 2013.
45. FamilyDoctor.org. Falls: How to Lower Your Risk. 2000; <http://familydoctor.org/familydoctor/en/prevention-wellness/staying-healthy/first-aid/falls-how-to-lower-your-risk.html>. Accessed August, 2013.