

Dysphagia: Evaluation and Collaborative Management

John M. Wilkinson, MD; Don Chamil Codipilly, MD; and Robert P. Wilfahrt, MD

Mayo Clinic College of Medicine and Science, Rochester, Minnesota

Dysphagia is common but may be underreported. Specific symptoms, rather than their perceived location, should guide the initial evaluation and imaging. Obstructive symptoms that seem to originate in the throat or neck may actually be caused by distal esophageal lesions. Oropharyngeal dysphagia manifests as difficulty initiating swallowing, coughing, choking, or aspiration, and it is most commonly caused by chronic neurologic conditions such as stroke, Parkinson disease, or dementia. Symptoms should be thoroughly evaluated because of the risk of aspiration. Patients with esophageal dysphagia may report a sensation of food getting stuck after swallowing. This condition is most commonly caused by gastroesophageal reflux disease and functional esophageal disorders. Eosinophilic esophagitis is triggered by food allergens and is increasingly prevalent; esophageal biopsies should be performed to make the diagnosis. Esophageal motility disorders such as achalasia are relatively rare and may be overdiagnosed. Opioid-induced esophageal dysfunction is becoming more common. Esophagogastroduodenoscopy is recommended for the initial evaluation of esophageal dysphagia, with barium esophagography as an adjunct. Esophageal cancer and other serious conditions have a low prevalence, and testing in low-risk patients may be deferred while a four-week trial of acid-suppressing therapy is undertaken. Many frail older adults with progressive neurologic disease have significant but unrecognized dysphagia, which significantly increases their risk of aspiration pneumonia and malnourishment. In these patients, the diagnosis of dysphagia should prompt a discussion about goals of care before potentially harmful interventions are considered. Speech-language pathologists and other specialists, in collaboration with family physicians, can provide structured assessments and make appropriate recommendations for safe swallowing, palliative care, or rehabilitation. (*Am Fam Physician*. 2021;103(2):97-106. Copyright © 2021 American Academy of Family Physicians.)

Many people occasionally experience difficult or impaired swallowing, but they often adapt their eating patterns to their symptoms and do not seek medical attention.¹ Among those who do seek care, the most common causes are generally benign and self-limited, and serious or life-threatening conditions are rare. However, many older adults with progressive neurologic disease have significant but unrecognized dysphagia, which increases their risk of aspiration pneumonia and malnourishment. In these patients, the diagnosis of dysphagia should prompt a discussion about goals of care. Understanding the basic pathophysiology of swallowing and the etiologies and clinical presentations of dysphagia allows family physicians to distinguish between oropharyngeal and esophageal pathology, make informed management decisions, and collaborate appropriately with specialists.

CME This clinical content conforms to AAFP criteria for CME. See CME Quiz on page 79.

Author disclosure: No relevant financial affiliations.

Patient information: A handout on this topic is available at <https://familydoctor.org/condition/dysphagia>.


Pathophysiology

Swallowing (deglutition) is a complex process involving voluntary and involuntary neuromuscular contractions coordinated to permit breathing and swallowing through the same anatomic pathway (*Figure 1*).² Deglutition is commonly divided into oropharyngeal and esophageal stages. In the oropharyngeal stage, food is chewed and mixed with saliva to form a bolus of appropriate consistency in the mouth. With the initiation of the swallow, the bolus is propelled into the oropharynx by the tongue. Other structures simultaneously seal the nasopharynx and larynx to prevent regurgitation or aspiration, and the lower esophageal sphincter begins to relax. In the esophageal stage, the food bolus passes the upper esophageal sphincter and enters the esophageal body, where it is propelled by peristalsis through the midthoracic and distal esophagus and into the stomach through the now fully relaxed lower esophageal sphincter.^{3,4}

Oropharyngeal Pathology

Oropharyngeal dysphagia is most commonly related to chronic neurologic conditions, particularly Parkinson disease, stroke, and dementia; it is not part of normal aging.⁵

FIGURE 1



Anatomy and physiology of deglutition. (A) Food (shown in green) is first chewed, mixed with saliva to form a bolus of appropriate consistency, and compressed against the hard palate by the tongue in preparation for swallowing. (B) The food bolus is propelled into the oropharynx, where the involuntary swallowing reflex is triggered. This swallowing reflex lasts approximately one second. (C) The epiglottis moves downward to cover the airway while striated pharyngeal muscles contract to continue moving the food bolus past the cricopharyngeus muscle (the physiologic upper esophageal sphincter) and into the proximal esophagus. This portion of the swallowing reflex also lasts approximately one second. (D) As the food bolus is propelled from the pharynx into the esophagus, involuntary contractions of the skeletal muscles of the upper esophagus force the bolus through the midthoracic and distal esophagus. The medulla controls this involuntary swallowing reflex, although voluntary swallowing may be initiated by the cerebral cortex. The lower esophageal sphincter relaxes from the initiation of the swallow until the food bolus is propelled into the stomach, which may take eight to 20 seconds.

Illustrations by Miriam Kirkman-Oh

Reprinted with permission from Palmer JB, Drennan JC, Baba M. Evaluation and treatment of swallowing impairments. *Am Fam Physician*. 2000; 61(8):2455-2456.

It may be the first symptom of a neuromuscular disorder, such as amyotrophic lateral sclerosis or myasthenia gravis.^{6,7}

Some chronic conditions, such as poor dentition, dentures, dry mouth (xerostomia), or medication adverse effects, may be poorly tolerated in patients who also have

progressive oropharyngeal dysfunction. Tardive dyskinesia with choreiform tongue movements related to long-term antipsychotic use may cause decompensation in older adults; it also may cause dysphagia in younger patients.⁸ Chronic cough related to angiotensin-converting enzyme inhibitor use may interfere with swallowing or be mistaken for aspiration.

Structural abnormalities (e.g., Zenker diverticulum, cricopharyngeal bars or tumors, chronic infections with *Candida* or herpes virus) and extrinsic compression from cervical osteophytes or goiter can also interfere with normal swallowing (Table 1).⁵⁻¹⁴

Esophageal Pathology

Gastroesophageal reflux disease (GERD), functional esophageal disorders, and eosinophilic esophagitis are the most common causes of esophageal dysphagia^{5,15} (Table 2^{1,5,13-15}). Less common causes include medications, obstructive lesions, and esophageal motility disorders.

GASTROESOPHAGEAL REFLUX DISEASE

GERD and recurrent acid exposure result in changes ranging from submucosal inflammation

BEST PRACTICES IN GERIATRICS

Recommendations from the Choosing Wisely Campaign

Recommendation	Sponsoring organization
Do not order "formal" swallow evaluation in stroke patients unless they fail their initial swallow screen.	American Academy of Nursing
Do not recommend percutaneous feeding tubes in patients with advanced dementia; instead, offer careful hand feeding.	American Academy of Hospice and Palliative Medicine American Geriatrics Society AMDA – The Society for Post-Acute and Long-Term Care Medicine

Source: For more information on the Choosing Wisely Campaign, see <https://www.choosingwisely.org>. For supporting citations and to search Choosing Wisely recommendations relevant to primary care, see <https://www.aafp.org/afp/recommendations/search.htm>.

SORT: KEY RECOMMENDATIONS FOR PRACTICE

Clinical recommendation	Evidence rating	Comments
Patients younger than 50 years who have esophageal dysphagia and no other worrisome symptoms should undergo a four-week trial of acid suppression therapy before endoscopy is performed. ^{9,26,27}	B	CAG and ACG guidelines; systematic review of seven retrospective cohort studies showing a positive predictive value of less than 1% for malignancy
Patients with apparent oropharyngeal symptoms but a negative evaluation should be referred for EGD to rule out esophageal pathology. ²⁸	C	Retrospective review of 3,668 consecutive patients; distal esophageal pathology was incorrectly perceived as arising from the neck or throat in 15% to 30% of cases
EGD is recommended for the initial assessment of patients with esophageal dysphagia; barium esophagography is recommended as an adjunct if EGD findings are negative. ^{9,29,30}	C	CAG, STS, and WGO guidelines; expert consensus based on limited evidence and cost-analysis; EGD has greater sensitivity and specificity than barium esophagography, with greater cost-effectiveness
For accurate diagnosis of eosinophilic esophagitis, biopsies from normal-appearing mucosa in the midthoracic and distal esophagus should be requested for all patients with unexplained solid food dysphagia. ^{15,17}	B	Expert consensus recommendation based on CAG, ACG, and AGA guidelines; early-stage eosinophilic esophagitis may not exhibit mucosal changes on endoscopy
Older patients with chronic illness or recent pneumonia should be screened for dysphagia; if it is present, the physician and patient should discuss goals of care. ^{22,43}	C	Expert consensus on dysphagia as a geriatric syndrome

ACG = American College of Gastroenterology; AGA = American Gastroenterological Association; CAG = Canadian Association of Gastroenterology; EGD = esophagogastroduodenoscopy; STS = Society of Thoracic Surgeons; WGO = World Gastroenterology Organisation.

A = consistent, good-quality patient-oriented evidence; **B** = inconsistent or limited-quality patient-oriented evidence; **C** = consensus, disease-oriented evidence, usual practice, expert opinion, or case series. For information about the SORT evidence rating system, go to <https://www.aafp.org/afpsort>.

TABLE 1

Causes of Oropharyngeal Dysphagia

Progressive chronic disease (geriatric syndrome)	Structural causes	Oral causes
Stroke	Head and neck cancers	Poor dentition or dentures
Parkinson disease	Recent surgery or radiation for head and neck cancers (altered anatomy)	Dry mouth (i.e., xerostomia)
Alzheimer and other dementias	Chemoradiation-induced mucositis and edema (short term)	Medications causing dry mouth (e.g., alpha and beta blockers, angiotensin-converting enzyme inhibitors, anticholinergics, anti-histamines, anxiolytics, calcium channel blockers, diuretics, muscle relaxants, tricyclic antidepressants)
Sarcopenia	Zenker diverticulum	Antipsychotic medications*
Neuromuscular disease	Cervical osteophytes	
Amyotrophic lateral sclerosis	Lymphadenopathy	
Myasthenia gravis	Goiter	
Multiple sclerosis	Cricopharyngeal bar	
Dermatomyositis/polymyositis (myopathies)		
Antipsychotic medications*		

Note: Causes are listed in order of prevalence. The relative prevalence of oropharyngeal dysphagia varies depending on the population studied; new-onset symptoms have a relatively low prevalence among primary care outpatients, whereas chronic progressive symptoms are increasingly prevalent among frail older patients. Some patients may have more than one cause.

*—Extrapyramidal effects from the use of antipsychotic medications (e.g., tardive dyskinesia with choreiform tongue movements) may cause dysphagia in younger patients and cause or aggravate other oral conditions in older patients.

Information from references 5–14.

TABLE 2

Causes of Esophageal Dysphagia

Gastroesophageal reflux disease and esophagitis (30% to 40%)

Eosinophilic esophagitis*

Functional dysphagia

Functional esophageal disorders (20% to 30%)†

Functional heartburn

Gastroesophageal reflux disease (nonerosive)

Globus pharyngeus‡

Reflux hypersensitivity

Medications (5%)

Pill esophagitis (direct irritation associated with ascorbic acid, bisphosphonates, ferrous sulfate, nonsteroidal anti-inflammatory drugs, potassium chloride, quinidine, and tetracyclines)

Reflux caused by decreased tone of lower esophageal sphincter (associated with alcohol, anticholinergics, benzodiazepines, caffeine, calcium channel blockers, nitrates, and tricyclic antidepressants)

Structural or mechanical conditions (5%)

Esophageal or peptic stricture (caused by erosive esophagitis)

Foreign body or food impaction (acute-onset dysphagia)

Malignancy (esophageal or gastric cancer, mediastinal mass with extrinsic compression)

Schatzki ring

Esophageal motility disorders (< 5%)§

Absent contractility

Achalasia

Distal esophageal spasm

Esophagogastric junction outflow obstruction

Hypercontractile (jackhammer) esophagus

Hypercontractile motility disorders

Opioid-induced esophageal dysfunction¶

Infections (< 5%)

Candida esophagitis

Cytomegalovirus esophagitis

Herpes simplex virus esophagitis

Rheumatologic conditions (< 5%)

Systemic sclerosis (scleroderma)

Note: Percentages are estimates of relative prevalence among primary care outpatients based on survey data and studies in different populations. Some patients may have more than one condition.

*—The prevalence of eosinophilic esophagitis is increasing.

†—Functional esophageal disorders may be underrecognized; actual prevalence is likely higher.

‡—Globus pharyngeus does not cause dysphagia, but the pathophysiology is likely similar to functional disorders.

§—Esophageal motility disorders may be overdiagnosed; patients may actually have other unrecognized conditions.

¶—Opioid-induced esophageal dysfunction is increasingly recognized and often indistinguishable from esophageal motility disorders; the exact prevalence is unclear.

Information from references 1, 5, and 13-15.

and dysmotility to erosive esophagitis and stricture. Patients with GERD may experience dysphagia even in the absence of apparent mucosal damage.¹⁶

EOSINOPHILIC ESOPHAGITIS

Eosinophilic esophagitis is an increasingly common inflammatory condition triggered by food allergens.¹⁵ Chronic eosinophilic infiltration leads to progressive fibrosis, esophageal rings and furrows, and dysmotility.¹⁷

FUNCTIONAL ESOPHAGEAL DISORDERS

Like irritable bowel syndrome or functional dyspepsia, functional esophageal disorders are thought to be caused by abnormalities of gut-brain interaction and central nervous system processing. People with these conditions may be hypervigilant about minor symptoms or hypersensitive to even physiologic amounts of acid.¹⁸ Patients may report dysphagia, although chest pain and heartburn are more common.¹⁹ These disorders may account for many patients who report intermittent difficulties with swallowing but never seek care,¹ as well as those in whom no explanation is found even after extensive investigations.¹⁵

MEDICATIONS

Medications may cause dysphagia as a result of direct mucosal injury (pill esophagitis), impaired esophageal motility, or lower esophageal sphincter relaxation and reflux.

OBSTRUCTIVE LESIONS

Esophageal cancer, strictures, Schatzki rings, and webs must be considered in patients presenting with dysphagia. However, the prevalence of these conditions is relatively low, particularly in patients younger than 50 years.^{9,10}

ESOPHAGEAL MOTILITY DISORDERS

Esophageal motility disorders such as achalasia, distal esophageal spasm, and systemic sclerosis (scleroderma) are rare. Similar to opioid-induced bowel dysfunction and constipation, opioid-induced esophageal dysfunction, although not as common, is being increasingly recognized.²⁰ Because this condition is often indistinguishable from other esophageal motility disorders, its prevalence is unclear.²¹

Initial Evaluation

The first step in the evaluation of a patient with dysphagia is to distinguish between oropharyngeal and esophageal pathology, based on characteristic symptoms. Clinical features and their temporal profile (*Table 3*^{13,14,22}), in addition to physical examination findings (*Table 4*^{13,14}), may suggest specific diagnoses and guide further testing and management.

TABLE 3

Questions to Assist in the Localization and Diagnosis of Dysphagia

"What about swallowing?"

Effective initial question for screening; as effective as longer screening instruments for prompting further questioning.²²

"What happens when you swallow? Does swallowing make you cough?"

Feeling that food is stuck suggests esophageal origin, regardless of the perceived location.

Trouble initiating a swallow, choking, coughing, or nasopharyngeal regurgitation suggests oropharyngeal dysphagia.

"How long has this been occurring? Is it getting worse? Do you hesitate to go out to eat because of it?"

Acute symptoms may be caused by pill esophagitis, infection, or foreign body impaction.

Intermittent symptoms suggest intermittent use of medications or opioids, esophageal motility disorders, or inconsistent denture use.

Progressive or worsening symptoms suggest malignancy.

Dysphagia may lead to social isolation.

"Do you have trouble chewing your food?"

Bruxism and jaw pain suggest temporomandibular joint arthritis or pain disorders.

Trouble chewing food suggests poorly fitting dentures, dental disease, or xerostomia.

Weakness with chewing suggests myasthenia gravis, giant cell arteritis, or myopathy (dermatomyositis).

"What gets stuck? Solids only? Solids and liquids?"

Trouble with liquids and solids suggests an esophageal motility disorder (e.g., achalasia).

Trouble with liquids only suggests oropharyngeal pathology.

Trouble with solids only suggests a mechanical obstruction of the esophagus, either intrinsic (stricture, web, or tumor) or extrinsic (mediastinal mass or thyromegaly).

"Do you have any other symptoms?"

Dyspepsia suggests a complication of reflux (e.g., stricture).

Halitosis with regurgitation may be due to an esophageal diverticulum or achalasia.

Weight loss from dysphagia suggests neoplasia or advanced disease.

"What medications do you take, including over-the-counter medications? Do you smoke? How much alcohol do you drink? Do you use opioids?"

Nonsteroidal anti-inflammatory drugs, potassium chloride, anticholinergics (including over-the-counter antihistamines or medications for overactive bladder), tricyclic antidepressants, tobacco, and alcohol can cause xerostomia and difficulty swallowing.

Opioids can reduce esophageal motility.

Use of proton pump inhibitors or histamine H₂ blockers suggests complications from gastroesophageal reflux disease.

"Do you have asthma or any food or environmental allergies?"

Positive response should prompt consideration of eosinophilic esophagitis; random biopsies should be obtained during esophago-gastroduodenoscopy to evaluate for this diagnosis.

Information from references 13, 14, and 22.

Most patients with dysphagia have esophageal dysfunction caused by benign and self-limited conditions, including functional esophageal disorders.¹ Oropharyngeal symptoms, particularly in patients without known comorbidities, are more worrisome; they may be the initial presentation of malignancy or neurodegenerative illness.⁹

GLOBUS PHARYNGEUS

Globus pharyngeus, an intermittent nonpainful sensation of a lump, foreign body, or phlegm in the neck or throat, should be distinguished from dysphagia. Globus pharyngeus has a benign natural course, typically improves with swallowing, and generally does not require further evaluation beyond a careful history and examination. It is commonly associated with GERD or anxiety.²³

SYMPTOMS AND HISTORY

Patients with oropharyngeal dysphagia often report choking, coughing, drooling, nasal regurgitation, difficulty initiating a swallow, or needing repeated swallows to clear food from the mouth. They may have hoarseness or other voice changes, including a "wet" voice. Patients will usually accurately localize their symptoms to the throat or neck.

Patients with esophageal dysfunction typically do not have difficulty initiating a swallow, but report a sensation of food getting stuck after swallowing. Patients with obstructive lesions report progressive symptoms occurring mainly with solids, whereas those with motility disorders often have intermittent dysphagia with solids and liquids.

Regurgitation of undigested food, particularly at night, is characteristic of achalasia or Zenker diverticulum. Painful swallowing (odynophagia) suggests an infectious process such as esophageal candidiasis or viral esophagitis. Distal esophageal spasm may also cause pain with swallowing, but this condition is much less prevalent.²⁴

RISK ASSESSMENT

Patients with dysphagia who also report weight loss, fever, gastrointestinal bleeding, or odynophagia, or who have unusually severe or rapidly progressive symptoms, especially older adults and those with a history of cancer or surgery, should have a more comprehensive expedited evaluation. In addition to esophagogastroduodenoscopy (EGD) and barium esophagography, the evaluation may include laryngoscopy to check for pharyngeal lesions and computed tomography to detect extrinsic masses.^{9,25}

TABLE 4

Physical Examination for Dysphagia

Examination component	Physical findings	Potential diagnosis or consideration
General		
Mental status	Obtunded or intoxicated	Transient and self-limited dyscoordination of swallow; address underlying causes
Nutritional state	Cachexia	Neoplasia
Overall fitness	Sarcopenia	Chronic disease
Strength	Weakness/fatigability	Myasthenia gravis
Integumentary		
Skin inspection	Needle tracks or sores; cold, clammy skin	Substance use or opioid-induced esophageal dysfunction
	Sausage digits or Raynaud phenomenon	Connective tissue disease (e.g., scleroderma)
Head, eyes, ears, nose, and throat		
Eyes	Ptosis (fatigable), diplopia	ALS
Mouth	Cervical or supraclavicular lymphadenopathy	Infectious esophagitis or malignancy
	Dry mouth (xerostomia)	Connective tissue disease, medication adverse effect, or tobacco use
	Poor dentition, poorly fitting dentures	Inability to comfortably or effectively form a food bolus precludes safe deglutition
	Thyromegaly or goiter	Extrinsic esophageal compression
Tongue deviates, tongue fasciculations		ALS or cranial nerve defects
Observe a swallow	Coughing, choking, or drooling	Drooling and nasopharyngeal regurgitation suggest oropharyngeal localization; coughing confirms an aspiration risk
Speech	Weak or breathy voice, dysarthria	Vocal cord pathology or ALS
	"Wet" voice	Laryngeal aspiration (likely chronic)
Abdominal		
Inspection	Signs of portal hypertension (e.g., varicosities, jaundice, distension)	Esophageal varices causing dysphagia by mass effect and peristalsis disruption
Palpation and percussion	Organomegaly	Malignancy
Neurologic		
Cranial nerves	Absent gag reflex	Cranial nerves IX and X affected
	Asymmetric facial motor findings	Cranial nerves V, VII, and XII share sensory and motor control over the muscles of expression and the tongue
Gait	Abnormal gait	Weakness generalizes to swallowing

ALS = amyotrophic lateral sclerosis.

Information from references 13 and 14.

Isolated dysphagia does not necessarily require immediate testing. Patients younger than 50 years are considered low risk, particularly if they have had intermittent symptoms of GERD or dyspepsia for more than six months. Testing may be safely deferred in these patients pending a treatment trial, such as a four-week course of a proton pump inhibitor.^{9,26,27}

Patients with oropharyngeal symptoms, hoarseness, or coughing provoked by swallowing sips of water must be evaluated for malignancy and other obstructive lesions, as well as aspiration risk. They should be referred to an otorhinolaryngologist or an appropriately trained speech-language pathologist for further imaging or formal swallowing studies.⁹ The specific screening instruments and tests used, as well as the health professionals administering them, vary based on local practice, expertise, and resources.

Patients with persistent symptoms and a negative oropharyngeal evaluation should be referred for EGD to rule out esophageal pathology.²⁸ Because of overlapping sensory innervation, the actual level of obstruction may be lower than perceived by the patient; in up to one-third of cases, symptoms in the throat or neck are caused by lesions in the distal esophagus.²⁸

Esophageal Dysphagia

TESTING

EGD is the recommended initial test for patients with suspected esophageal dysphagia, followed by barium esophagography if EGD findings are negative.^{9,29,30} EGD directly visualizes the esophageal lumen and mucosa; can detect obstructive lesions, infections, inflammatory conditions, and reflux esophagitis; allows for stricture dilatation and biopsy; and is generally more cost-effective than barium esophagography. However, barium esophagography is preferred over EGD for detecting subtle narrowing or esophageal webs

that may be amenable to dilatation, as well as extrinsic compression of the esophagus. Patients on anticoagulant therapy should undergo barium esophagography first to determine whether a dilatation procedure is needed, which requires stopping anticoagulant therapy. However, routine biopsies can generally be done in these patients without discontinuing therapy.³¹

For accurate diagnosis of eosinophilic esophagitis, biopsies from the midthoracic and distal esophagus, even if the mucosa appears to be normal, should be requested for all patients with unexplained dysphagia.^{15,17} Although the endoscopist or radiologist may suspect achalasia or a hypercontractile motility disorder based on EGD or barium esophagography findings,³² high-resolution esophageal manometry is required to definitively diagnose these conditions.^{33,34}

AVOIDING OVERDIAGNOSIS

Recent research has suggested that although high-resolution esophageal manometry is important for the diagnosis of achalasia, it may result in overdiagnosis of esophageal motility disorders and lead to overtreatment with invasive endoscopic therapies.³⁵ Many patients who are thought to have hypercontractile motility disorders may actually have functional esophageal symptoms unrelated to manometric findings.³⁶ Testing for these uncommon motility disorders can be safely deferred for at least a few months to allow for a trial of optimal medical management of more common conditions. Delayed diagnosis of achalasia does not increase the risk of esophageal cancer.³⁷

INITIAL TREATMENT

Patients with GERD symptoms, esophagitis, or peptic stricture should undergo acid suppression therapy with standard doses of proton pump inhibitors for eight to 12 weeks.^{38,39} Patients with eosinophilic esophagitis may also respond to this regimen, but most require elimination diets, topical steroids, or both.^{40,41}

For patients with functional dysphagia, functional chest pain, heartburn, or globus pharyngeus, reassurance about the benign and self-limited nature of these conditions, mindful eating, avoidance of trigger foods or situations, and a trial of acid suppression may be helpful. Tricyclic antidepressants, which modulate esophageal visceral hypersensitivity and hypervigilance, are somewhat effective in reducing symptoms.¹⁹ The dosages used in dyspepsia trials were 25 mg of amitriptyline or 50 mg of imipramine per day.⁴² Cognitive behavior therapy is helpful in patients with functional dyspepsia²⁷ and may be considered if medical therapy is ineffective.¹⁹

Patients with pill esophagitis should take their medication with copious amounts of water and then remain upright for

30 minutes. If opioid-induced esophageal dysfunction is suspected, the patient should be helped to discontinue use or at least reduce the dosage.^{20,21}

Oropharyngeal Dysphagia

Up to one-half of debilitated and frail older adults have some degree of dysphagia and silent aspiration, although they often are not aware of the problem. Patients who have had a stroke and those with Parkinson disease, dementia, or sarcopenia are at particular risk.^{11,12} Dysphagia may be considered a geriatric syndrome. It is multifactorial and may be triggered by acute insults or gradual decline; it leads to poor outcomes such as malnutrition, social isolation, dehydration, weight loss, and aspiration pneumonia; and treatment requires multidisciplinary interventions.⁴³

SCREENING

Although hospitalized patients are routinely assessed for dysphagia, particularly after a stroke, impairments in community-dwelling older adults may not be recognized. All patients with chronic illness or recent pneumonia should be periodically screened for dysphagia.⁴³ The single question “What about swallowing?” may be as effective as more detailed screening tools.²² Patients who are sufficiently alert and able to follow directions may also be observed swallowing a few sips of water.

AVOIDING OVERTREATMENT

In older patients with progressive chronic illness, a diagnosis of oropharyngeal dysphagia should prompt a discussion about goals of care. The family physician is well-suited to provide anticipatory guidance about the potential consequences of dysphagia, as well as a realistic assessment of the patient’s overall condition and long-term prospects. A formal swallowing evaluation may be needed to guide this discussion (*Table 5*).^{22,44–49}

Interventions for oropharyngeal dysphagia have limited benefit because of the inevitable decline in most patients. Nasogastric tube feeding does not result in any survival benefit or reduce rates of aspiration pneumonia,^{50,51} and it is associated with significant harms.^{52,53} Hand feeding is as effective and is generally recommended; caregivers and care settings must promote choice and respect the preferences of patients and their surrogates.⁵⁴

MULTIDISCIPLINARY EVALUATION AND REHABILITATION

Speech-language pathologists use various tests ranging from bedside assessment to instrumented swallowing studies to determine specific deficits, the patient’s potential for improvement, and the most appropriate dietary

TABLE 5

Swallowing Evaluation for Oropharyngeal Dysphagia

Video fluoroscopic swallowing study (modified barium swallow)

Preferred diagnostic test; more accurate than bedside swallow assessment for detection of aspiration and allows for more precise treatment recommendations^{44,45}

Performed by radiologist and speech-language pathologist in video fluoroscopy suite⁴⁶

Fiber-optic endoscopic evaluation

Supplementary tool to video fluoroscopic swallowing study; good correlation in detection of aspiration and bolus residue in the pharynx and pharyngeal-laryngeal area

Trial of compensatory maneuvers using different viscosities (thin liquid, thick liquid, puree, solid food) to potentially improve swallowing effectiveness

Performed by speech-language pathologist in outpatient/bedside setting with physician review; easy to perform and well tolerated⁴⁶

Bedside swallow assessment

Structured observation of eating and drinking by speech-language pathologist or other trained observer

Trial swallows using different viscosities (thin liquid, thick liquid, puree, solid food): check for coughing, choking, "wet" voice, or piecemeal swallowing (i.e., multiple swallows per bolus)

Water-swallowing test (5 to 30 mL) or repetitive saliva-swallowing test: check for coughing, choking, or "wet" voice

Pulse oximetry combined with trial swallows or water-swallowing test: check for oxygen desaturation > 2% to 3%

Patient self-evaluation: selected questionnaires to assess health status, dysphagia severity, and quality of life

Eating Assessment Tool⁴⁷

Single question ("What about swallowing?") may be as effective as more detailed screening tools²²

Swallowing Quality of Life questionnaire⁴⁸

Sydney Swallow Questionnaire⁴⁹

Information from references 22 and 44-49.

TABLE 6

Management of Oropharyngeal Dysphagia

Diet modifications

Mindful eating

Avoiding foods likely to cause dysphagia

Chewing carefully

Cutting food into smaller pieces

Drinking liquids to dilute food bolus

Eating slowly

Lubricating with sauces

Taking smaller bites

Mechanical soft diet to compensate for impaired chewing due to weakness or poor dentition

Modified consistency diet to compensate for impaired swallowing (thickened food and liquids with increased viscosity and cohesiveness slow transport speed)

Pureed diet to compensate for impaired chewing due to weakness, poor dentition, or dry mouth with impaired bolus formation

Tasting diet⁵⁵

Patient-centered approaches to feeding and environments conducive to eating should be part of usual care for all older adults with chronic debilitating illness²²

Swallowing rehabilitation*

Muscular reconditioning exercises to strengthen jaw, lips, and tongue

In clinically stable patients who have potential for improvement (e.g., after a stroke), the goal is long-term change in control of swallowing via neuroplasticity

In patients with progressive conditions (e.g., Parkinson disease), the goal is maintaining current swallowing status as long as possible

Compensatory safe-swallow techniques (positioning to compensate for weakness and dysfunction)

Eating upright

Chin-tuck maneuver for patients with stroke or degenerative disease changes pharyngeal dimensions to direct the food bolus toward the pharynx and esophagus, compensates for delay of glottic closure, and reduces aspiration risk during swallowing⁵⁶

Head-turn maneuver for patients with unilateral weakness (turning head toward weaker side) uses gravity to push the bolus toward the stronger side⁵⁷

Enteral feeding†

Nasogastric tube feeding: typically placed during the first week after stroke to allow for administration of nutrition and medication

Percutaneous endoscopic gastrostomy: used longer term if dysphagia persists after the first week; best for patients with some degree of rehabilitation potential; does not improve mortality or reduce aspiration risk

*—Patients with potential for recovery and the ability to remember and follow instructions are more likely to benefit from swallowing rehabilitation than those with progressive symptoms.

†—Should be preceded by a discussion about goals of care and shared decision-making, balancing aspiration risk with quality of life.

Information from references 22 and 55-57.

modifications and swallow therapies. Thickened liquids and foods with specific textures are often helpful in reducing aspiration risk. Patients with the ability to remember and follow instructions may be taught compensatory maneuvers of the head, neck, and chin, as well as rehabilitative exercises to promote safer swallowing (Table 6).^{22,55-57}

Palliative care specialists can help facilitate patient-centered feeding. One hospice, in collaboration with professional chefs, maintains a website featuring recipes that have been adapted to emphasize pleasurable textures and tastes.⁵⁵ Social connections and rituals involving food remain important even when swallowing is no longer possible.

This article updates previous articles on this topic by Spieker,¹³ and by Palmer, et al.²

Data Sources: We searched PubMed and Google Scholar using the following search terms alone and in various combinations: dysphagia, esophageal spasm, gastroesophageal reflux disease, eosinophilic esophagitis, functional esophageal disorders, achalasia, esophageal motility disorder, and aspiration. We examined clinical trials, meta-analyses, review articles, and clinical guidelines, as well as the bibliographies of selected articles. The Cochrane database and Essential Evidence Plus were also searched. Search dates: August 2019 to July 2020.

The authors thank Darlene E. Graner, SLPD, CCC-SLP, and Emily A. Hosfield, MS, CCC-SLP, for their review of the manuscript.

The Authors

JOHN M. WILKINSON, MD, is a consultant in the Department of Family Medicine and an associate professor in the Mayo Clinic Alix School of Medicine, Mayo Clinic College of Medicine and Science, Rochester, Minn.

DON CHAMIL CODIPILLY, MD, is a fellow in the Division of Gastroenterology and Hepatology at Mayo Clinic College of Medicine and Science.

ROBERT P. WILFAHRT, MD, is a consultant in the Department of Family Medicine and an assistant professor in the Mayo Clinic Alix School of Medicine, Mayo Clinic College of Medicine and Science.

Address correspondence to John M. Wilkinson, MD, Mayo Clinic Alix School of Medicine, Mayo Clinic College of Medicine and Science, 200 1st St. SW, Rochester, MN 55905 (email: wilkinson.john@mayo.edu). Reprints are not available from the authors.

References

- Wilkins T, Gillies RA, Thomas AM, et al. The prevalence of dysphagia in primary care patients: a HarnesNet Research Network study. *J Am Board Fam Med*. 2007;20(2):144-150.
- Palmer JB, Drennan JC, Baba M. Evaluation and treatment of swallowing impairments. *Am Fam Physician*. 2000;61(8):2453-2462. Accessed May 2, 2020. <https://www.aafp.org/afp/2000/0415/p2453.html>
- Sasegbon A, Hamdy S. The anatomy and physiology of normal and abnormal swallowing in oropharyngeal dysphagia. *Neurogastroenterol Motil*. 2017;29(11):e13100.
- Shaw SM, Martino R. The normal swallow: muscular and neurophysiological control. *Otolaryngol Clin North Am*. 2013;46(6):937-956.
- Cho SY, Choung RS, Saito YA, et al. Prevalence and risk factors for dysphagia: a USA community study. *Neurogastroenterol Motil*. 2015;27(2):212-219.
- Traynor BJ, Codd MB, Corr B, et al. Clinical features of amyotrophic lateral sclerosis according to the El Escorial and Airlie House diagnostic criteria: a population-based study. *Arch Neurol*. 2000;57(8):1171-1176.
- Scherer K, Bedlack RS, Simel DL. Does this patient have myasthenia gravis? *JAMA*. 2005;293(15):1906-1914.
- Miarons Font M, Rofes Salsench L. Antipsychotic medication and oropharyngeal dysphagia: systematic review. *Eur J Gastroenterol Hepatol*. 2017;29(12):1332-1339.
- Liu LWC, Andrews CN, Armstrong D, et al. Clinical practice guidelines for the assessment of uninvestigated esophageal dysphagia. *J Can Assoc Gastroenterol*. 2018;1(1):5-19.
- Roden DF, Altman KW. Causes of dysphagia among different age groups: a systematic review of the literature. *Otolaryngol Clin North Am*. 2013;46(6):965-987.
- Serra-Prat M, Hinojosa G, López D, et al. Prevalence of oropharyngeal dysphagia and impaired safety and efficacy of swallow in independently living older persons. *J Am Geriatr Soc*. 2011;59(1):186-187.
- Lin LC, Wu SC, Chen HS, et al. Prevalence of impaired swallowing in institutionalized older people in Taiwan. *J Am Geriatr Soc*. 2002;50(6):1118-1123.
- Spieker MR. Evaluating dysphagia. *Am Fam Physician*. 2000;61(12):3639-3648. Accessed May 4, 2020. <https://www.aafp.org/afp/2000/0615/p3639.html>
- Jansson-Knodell CL, Codipilly DC, Leggett CL. Making dysphagia easier to swallow: a review for the practicing clinician. *Mayo Clin Proc*. 2017;92(6):965-972.
- Kidambi T, Toto E, Ho N, et al. Temporal trends in the relative prevalence of dysphagia etiologies from 1999-2009. *World J Gastroenterol*. 2012;18(32):4335-4341.
- Richter JE, Rubenstein JH. Presentation and epidemiology of gastroesophageal reflux disease. *Gastroenterology*. 2018;154(2):267-276.
- Straumann A, Katzka DA. Diagnosis and treatment of eosinophilic esophagitis. *Gastroenterology*. 2018;154(2):346-359.
- Drossman DA. Functional gastrointestinal disorders: history, pathophysiology, clinical features and Rome IV. *Gastroenterology*. 2016;150(6):1262-1279.e2.
- Aziz Q, Fass R, Gyawali CP, et al. Functional esophageal disorders. *Gastroenterology*. 2016;150(6):1368-1379.
- Kraichely RE, Arora AS, Murray JA. Opiate-induced esophageal dysmotility. *Aliment Pharmacol Ther*. 2010;31(5):601-606.
- Ratuapli SK, Crowell MD, DiBaise JK, et al. Opioid-induced esophageal dysfunction (OIED) in patients on chronic opioids. *Am J Gastroenterol*. 2015;110(7):979-984.
- Heijnen BJ, Speyer R, Bülow M, et al. 'What about swallowing?' Diagnostic performance of daily clinical practice compared with the Eating Assessment Tool-10. *Dysphagia*. 2016;31(2):214-222.
- Lee BE, Kim GH. Globus pharyngeus: a review of its etiology, diagnosis and treatment. *World J Gastroenterol*. 2012;18(20):2462-2471.
- Khalaf M, Chowdhary S, Elias PS, et al. Distal esophageal spasm: a review. *Am J Med*. 2018;131(9):1034-1040.
- Rhatigan E, Tyrmpas I, Murray G, et al. Scoring system to identify patients at high risk of esophageal cancer. *Br J Surg*. 2010;97(12):1831-1837.
- Vakil N, Moayyedi P, Fennerty MB, et al. Limited value of alarm features in the diagnosis of upper gastrointestinal malignancy: systematic review and meta-analysis. *Gastroenterology*. 2006;131(2):390-401.

27. Moayyedi P, Lacy BE, Andrews CN, et al. ACG and CAG clinical guideline: management of dyspepsia [published correction appears in *Am J Gastroenterol*. 2017;112(9):1484]. *Am J Gastroenterol*. 2017;112(7):988-1013.
28. Ashraf HH, Palmer J, Dalton HR, et al. Can patients determine the level of their dysphagia? *World J Gastroenterol*. 2017;23(6):1038-1043.
29. Varghese TK Jr, Hofstetter WL, Rizk NP, et al. The Society of Thoracic Surgeons guidelines on the diagnosis and staging of patients with esophageal cancer. *Ann Thorac Surg*. 2013;96(1):346-356.
30. Malagelada J, Bazzoli F, Boeckxstaens G, et al. World Gastroenterology Organisation global guidelines: dysphagia. Updated September 2014. Accessed March 17, 2020. <https://www.worldgastroenterology.org/guidelines/global-guidelines/dysphagia/dysphagia-english>
31. Acosta RD, Abraham NS, Chandrasekhara V, et al.; ASGE Standards of Practice Committee. The management of antithrombotic agents for patients undergoing GI endoscopy [published correction appears in *Gastrointest Endosc*. 2016;83(3):678]. *Gastrointest Endosc*. 2016;83(1):3-16.
32. McCormick SE, Kozarek RA. Endoscopic evaluation of esophageal motility disorders. *GI Motility Online*. 2006. Accessed March 17, 2020. <https://www.nature.com/gimo/contents/pt1/full/gimo29.html>
33. Kahrilas PJ, Bredenoord AJ, Fox M, et al.; International High Resolution Manometry Working Group. The Chicago Classification of esophageal motility disorders, v3.0. *Neurogastroenterol Motil*. 2015;27(2):160-174.
34. Almansa C, Heckman MG, DeVault KR, et al. Esophageal spasm: demographic, clinical, radiographic, and manometric features in 108 patients. *Dis Esophagus*. 2012;25(3):214-221.
35. Mion F, Marjoux S, Subtil F, et al. Botulinum toxin for the treatment of hypercontractile esophagus: results of a double-blind randomized sham-controlled study. *Neurogastroenterol Motil*. 2019;31(5):e13587.
36. Bredenoord AJ. Hypercontractile esophageal motility disorder or functional esophageal symptoms and unrelated hypercontractility? *Neurogastroenterol Motil*. 2019;31(5):e13594.
37. Zendehelel K, Nyrén O, Edberg A, et al. Risk of esophageal adenocarcinoma in achalasia patients, a retrospective cohort study in Sweden. *Am J Gastroenterol*. 2011;106(1):57-61.
38. Katz PO, Gerson LB, Vela MF. Guidelines for the diagnosis and management of gastroesophageal reflux disease [published correction appears in *Am J Gastroenterol*. 2013;108(10):1672]. *Am J Gastroenterol*. 2013;108(3):308-328.
39. Weijenborg PW, Cremonini F, Smout AJ, et al. PPI therapy is equally effective in well-defined non-erosive reflux disease and in reflux esophagitis: a meta-analysis. *Neurogastroenterol Motil*. 2012;24(8):747-757.e350.
40. Dellon ES, Katzka DA, Collins MH, et al.; MP-101-06 Investigators. Budesonide oral suspension improves symptomatic, endoscopic, and histologic parameters compared with placebo in patients with eosinophilic esophagitis. *Gastroenterology*. 2017;152(4):776-786.e5.
41. Lucendo AJ, Arias Á, González-Cervera J, et al. Empiric 6-food elimination diet induced and maintained prolonged remission in patients with adult eosinophilic esophagitis: a prospective study on the food cause of the disease. *J Allergy Clin Immunol*. 2013;131(3):797-804.
42. Ford AC, Luthra P, Tack J, et al. Efficacy of psychotropic drugs in functional dyspepsia: systematic review and meta-analysis. *Gut*. 2017;66(3):411-420.
43. Baijens LW, Clavé P, Cras P, et al. European Society for Swallowing Disorders - European Union Geriatric Medicine Society white paper: oropharyngeal dysphagia as a geriatric syndrome. *Clin Interv Aging*. 2016;11:1403-1428.
44. Braun T, Juenemann M, Viard M, et al. What is the value of fibre-endoscopic evaluation of swallowing (FEES) in neurological patients? A cross-sectional hospital-based registry study. *BMJ Open*. 2018;8(3):e019016.
45. Alagiakrishnan K, Bhanji RA, Kurian M. Evaluation and management of oropharyngeal dysphagia in different types of dementia: a systematic review. *Arch Gerontol Geriatr*. 2013;56(1):1-9.
46. Brady S, Donzelli J. The modified barium swallow and the functional endoscopic evaluation of swallowing. *Otolaryngol Clin North Am*. 2013;46(6):1009-1022.
47. Belafsky PC, Mouadeb DA, Rees CJ, et al. Validity and reliability of the Eating Assessment Tool (EAT-10). *Ann Otol Rhinol Laryngol*. 2008;117(12):919-924.
48. McHorney CA, Robbins J, Lomax K, et al. The SWAL-QOL and SWAL-CARE outcomes tool for oropharyngeal dysphagia in adults: III. Documentation of reliability and validity. *Dysphagia*. 2002;17(2):97-114.
49. Wallace KL, Middleton S, Cook IJ. Development and validation of a self-report symptom inventory to assess the severity of oropharyngeal dysphagia. *Gastroenterology*. 2000;118(4):678-687.
50. Sampson EL, Candy B, Jones L. Enteral tube feeding for older people with advanced dementia. *Cochrane Database Syst Rev*. 2009;(2):CD007209.
51. Teno JM, Gozalo PL, Mitchell SL, et al. Does feeding tube insertion and its timing improve survival? *J Am Geriatr Soc*. 2012;60(10):1918-1921.
52. Lubart E, Leibovitz A, Dror Y, et al. Mortality after nasogastric tube feeding initiation in long-term care elderly with oropharyngeal dysphagia—the contribution of refeeding syndrome. *Gerontology*. 2009;55(4):393-397.
53. Givens JL, Selby K, Goldfeld KS, et al. Hospital transfers of nursing home residents with advanced dementia. *J Am Geriatr Soc*. 2012;60(5):905-909.
54. American Geriatrics Society Ethics Committee and Clinical Practice and Models of Care Committee. American Geriatrics Society feeding tubes in advanced dementia position statement. *J Am Geriatr Soc*. 2014;62(8):1590-1593.
55. L'Institut de Tourisme et d'Hotellerie du Quebec; Palliative Care McGill, Department of Family Medicine, McGill University; The International Congress on Palliative Care; et al. Eating well, always. 2019. Accessed March 17, 2020. https://www.mcgill.ca/palliativecare/files/palliative_care/pal2019_recipe_book_eng_final_0.pdf
56. Saconato M, Chiari BM, Lederman HM, et al. Effectiveness of chin-tuck maneuver to facilitate swallowing in neurologic dysphagia. *Int Arch Otorhinolaryngol*. 2016;20(1):13-17.
57. Vose A, Nonnenmacher J, Singer ML, et al. Dysphagia management in acute and sub-acute stroke. *Curr Phys Med Rehabil Rep*. 2014;2(4):197-206.