

U.S. Preventive Services Task Force

Screening for High Blood Pressure in Children and Adolescents: U.S. Preventive Services Task Force Recommendation Statement

Summary of Recommendation

The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of screening for high blood pressure in children and adolescents (*Table 1*). **I statement.**

Importance

The prevalence of hypertension (both primary and secondary) in children and adolescents in the United States ranges from 3% to 4% in most studies.¹⁻⁵ Primary hypertension in children and adolescents occurs primarily in children older than 13 years and has no known cause but is associated with several risk factors, including family history and higher body mass index.¹ Secondary hypertension occurs primarily in younger children and is most commonly caused by genetic disorders, renal disease, endocrine disorders, or cardiovascular abnormalities.^{1,6}

USPSTF Assessment of Magnitude of Net Benefit

The USPSTF concludes that the evidence to support screening for high blood pressure in children and adolescents aged 3 to 18 years is insufficient and that the balance of benefits and harms cannot be determined.

See *Table 2* for more information on the USPSTF recommendation rationale and assessment. For more details on the methods the USPSTF uses to determine the net benefit, see the USPSTF Procedure Manual.⁷

As published by the USPSTF.

This summary is one in a series excerpted from the Recommendation Statements released by the USPSTF. These statements address preventive health services for use in primary care clinical settings, including screening tests, counseling, and preventive medications.

The complete version of this statement, including supporting scientific evidence, evidence tables, grading system, members of the USPSTF at the time this recommendation was finalized, and references, is available on the USPSTF website at <https://www.uspreventiveservicestaskforce.org/>.

This series is coordinated by Kenny Lin, MD, MPH, deputy editor.

A collection of USPSTF recommendation statements published in *AFP* is available at <https://www.aafp.org/afp/uspstf>.

Practice Considerations

PATIENT POPULATION UNDER CONSIDERATION

This recommendation applies to children and adolescents not known to have hypertension.

DEFINITIONS OF HYPERTENSION

Primary hypertension has no known cause. Secondary hypertension in children is most commonly caused by renal or renovascular disease. Other causes of secondary hypertension include congenital cardiovascular abnormalities, genetic disorders, and endocrine disorders.^{1,6}

The American Academy of Pediatrics revised its definitions of abnormal blood pressure in children and adolescents in 2017.⁸ Hypertension is defined as sustained high blood pressure. Elevated blood pressure (previously known as prehypertension) in children aged 1 to 13 years is defined as 3 auscultatory blood pressure measurements between the 90th and 94th percentile based on age, height, and sex, or 120 to 129/< 80 mm Hg.⁸ For children aged 1 to 13 years, hypertension is defined as 3 auscultatory blood pressure measurements at 3 different visits that are either equal to or above the 95th percentile based on age, height, and sex, or above 130/80 mm Hg, whichever measurement is lower.^{1,8} The thresholds for adolescents 13 years or older follow the adult guidelines from the American Heart Association and American College of Cardiology, regardless of sex and height: Stage 1 hypertension is defined as a blood pressure measurement of 130/80 to 139/89 mm Hg, and stage 2 hypertension is defined as a blood pressure measurement above 140/90 mm Hg.^{1,9}

SCREENING TESTS

Best practice is to measure blood pressure by auscultation of the upper right arm with an appropriately sized cuff^{1,8} at 3 different visits. (The right arm is preferred in children because standardized blood pressure tables reflect right arm readings and because low readings in the left arm could be a result of coarctation of the aorta.) Some organizations suggest that, as in adults, the diagnosis should be confirmed by ambulatory blood pressure monitoring. Ambulatory blood pressure monitoring uses portable devices that record blood pressure measurements at regular intervals (usually every 20 to 30 minutes) over 12 to 24 hours.⁸

TABLE 1

Screening for High Blood Pressure in Children and Adolescents: Clinical Summary of the USPSTF Recommendation

What does the USPSTF recommend?	For children and adolescents: The USPSTF found that the evidence is insufficient to assess the balance of benefits and harms of screening for high blood pressure in children and adolescents. More research is needed. I statement
To whom does this recommendation apply?	Children and adolescents not known to have hypertension.
What's new?	This recommendation is consistent with the 2013 USPSTF statement.
How to implement this recommendation?	There is insufficient evidence to recommend for or against screening for high blood pressure in children and adolescents. Clinicians should remain alert to the signs and/or symptoms of high blood pressure and evaluate as appropriate.
What are the other relevant USPSTF recommendations?	The USPSTF has issued other relevant recommendation statements at www.uspreventiveservicestaskforce.org , including: Screening for obesity in children and adolescents Screening for lipid disorders in children and adolescents
Where to read the full recommendation statement?	Visit the USPSTF website (https://www.uspreventiveservicestaskforce.org) to read the full recommendation statement. This includes more details on the rationale of the recommendation, including benefits and harms; supporting evidence; and recommendations of others.

USPSTF = U.S. Preventive Services Task Force.

SCREENING INTERVALS

There is no good evidence about how often blood pressure should be measured in children and adolescents.

TREATMENT

Hypertension in children is often treated with lifestyle and pharmacologic interventions. Lifestyle interventions for hypertension include weight loss in children who are overweight or obese, increased physical activity, and the Dietary Approaches to Stop Hypertension (DASH) diet, as well as education and counseling. Medication is generally used to treat hypertension that is unresponsive to lifestyle modification or that has a secondary cause.¹

SUGGESTIONS FOR PRACTICE REGARDING THE I STATEMENT

In deciding whether to screen for high blood pressure in children and adolescents, clinicians should consider the following.

Potential Preventable Burden. Important risk factors for primary hypertension in children and

adolescents are higher body mass index and a family history of hypertension. Other risk factors include low birth weight, male sex, and African American or Hispanic/Latino race/ethnicity. The increasing prevalence of hypertension in children and adolescents, possibly driven by childhood obesity, suggests that identification and treatment of hypertension may become a significant health care issue. The goal of identifying and treating children and adolescents with primary hypertension can be viewed within a larger framework of adult cardiovascular risk reduction, which includes addressing other biometric risk factors such as higher body mass index and lipid profiles and hyperglycemia. However, the variables for cardiovascular risk reduction in children are less understood than in adults.

Extending the adult framework for cardiovascular risk reduction to children and adolescents is limited by several methodological challenges that complicate the determination of the potential preventable burden. There is limited evidence about the clinical and epidemiologic significance of percentile thresholds used in children in terms

TABLE 2

Summary of USPSTF Rationale*

Rationale	Assessment
Detection	There is inadequate evidence about the accuracy of screening for high blood pressure in children and adolescents.
Benefits of early detection and intervention and treatment (based on direct or indirect evidence)	There is no direct evidence that screening for high blood pressure in children and adolescents delays or reduces adverse health outcomes. There is adequate evidence about the longitudinal association between high blood pressure in children and adolescents and high blood pressure and other intermediate outcomes in adults. There is inadequate evidence on the long-term effectiveness of interventions (pharmacologic, lifestyle, or lifestyle interventions combined with pharmacotherapy) in the treatment of high blood pressure in children and adolescents resulting in reduced blood pressure or improved intermediate outcomes in children or adults. There is inadequate evidence on the effectiveness of interventions in the treatment of high blood pressure in children and adolescents resulting in reduced adverse health outcomes associated with high blood pressure in adults.
Harms of early detection and intervention and treatment	There is inadequate evidence to assess the potential harms of screening for high blood pressure in children and adolescents. There is inadequate evidence to assess the potential harms of pharmacologic, lifestyle, or lifestyle interventions combined with pharmacotherapy for treatment of high blood pressure in children and adolescents.
USPSTF assessment	The benefits and harms of screening for blood pressure in children and adolescents are uncertain, and the balance of benefits and harms cannot be determined

USPSTF = U.S. Preventive Services Task Force.

*—See the eFigure in the Supplement for explanation of USPSTF grades and levels of evidence.

of their association with adult cardiovascular disease. In addition, the performance characteristics of current methods for diagnosing hypertension during childhood are limited and of concern because of possible high false-positive rates (i.e., elevated blood pressure measurements that later normalize).

One justification that has been suggested for screening is to identify secondary hypertension, a relatively rare condition. Younger children (younger than 6 years) are more likely than older children and adolescents to have a secondary cause of hypertension, such as renal disease, coarctation of the aorta, or endocrine disease. Secondary hypertension is unlikely to be the only clinical manifestation of the underlying disorder in these cases, and management is primarily targeted at treating the underlying condition, as well as controlling hypertension.¹

Potential Harms. Clinic-based screening for high blood pressure could result in false-positive results. Thus, unnecessary secondary evaluations

or treatments may be common, particularly with frequent blood pressure screening. Pharmacologic interventions have been shown to be well tolerated, but this has only been evaluated in studies over relatively short periods.¹ Treatment of hypertension in childhood and adolescence with pharmacologic agents is done for a much longer period, and therefore, adverse effects may be more likely to occur.

Current Practice. Current practice for blood pressure screening typically involves measurement in office-based health care settings as part of well-child or sports participation examinations, often in conjunction with other vital signs and growth parameters. Several organizations recommend routine measurement of blood pressure at well-child visits starting at age 3 years, based on consensus.¹

OTHER RELATED USPSTF RECOMMENDATIONS

The USPSTF recommends that clinicians screen for obesity in children 6 years or older and offer

or refer them to a comprehensive, intensive behavioral intervention to improve weight status (B recommendation).¹⁰ The USPSTF found insufficient evidence on screening for lipid disorders in children and adolescents 20 years or younger (I statement).¹¹ These recommendations are available on the USPSTF website (<https://www.uspreventiveservicestaskforce.org>).

This recommendation statement was first published in *JAMA*. 2020;324(18):1884-1895.

The "Update of Previous USPSTF Recommendation," "Supporting Evidence," "Research Needs and Gaps," and "Recommendations of Others" sections of this recommendation statement are available at <https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/blood-pressure-in-children-and-adolescents-hypertension-screening#full-recommendationstart>.

The USPSTF recommendations are independent of the U.S. government. They do not represent the views of the Agency for Healthcare Research and Quality, the U.S. Department of Health and Human Services, or the U.S. Public Health Service.

References

1. Gartlehner G, Vander Schaaf EB, Orr C, et al. Screening for high blood pressure in children and adolescents: systematic review for the U.S. Preventive Services Task Force. Evidence synthesis no. 193. AHRQ publication no. 20-05261-EF-1. Agency for Healthcare Research and Quality; 2020.
2. Dobson CP, Eide M, Nylund CM. Hypertension prevalence, cardiac complications, and antihypertensive medication use in children. *J Pediatr*. 2015;167(1):92-97.e1.
3. McNiece KL, Poffenbarger TS, Turner JL, et al. Prevalence of hypertension and pre-hypertension among adolescents. *J Pediatr*. 2007;150(6):640-644, 644.e1.
4. Hansen ML, Gunn PW, Kaelber DC. Underdiagnosis of hypertension in children and adolescents. *JAMA*. 2007; 298(8):874-879.
5. Chen X, Wang Y. Tracking of blood pressure from childhood to adulthood: a systematic review and meta-regression analysis. *Circulation*. 2008;117(25):3171-3180.
6. Flynn JT, Pierce CB, Miller ER III, et al.; Chronic Kidney Disease in Children Study Group. Reliability of resting blood pressure measurement and classification using an oscillometric device in children with chronic kidney disease. *J Pediatr*. 2012;160(3):434-440.e1.
7. U.S. Preventive Services Task Force. Procedure manual. Accessed April 3, 2020. <https://www.uspreventiveservices taskforce.org/uspstf/procedure-manual>
8. Flynn JT, Kaelber DC, Baker-Smith CM, et al.; Subcommittee on Screening and Management of High Blood Pressure in Children. Clinical practice guideline for screening and management of high blood pressure in children and adolescents [published corrections appear in *Pediatrics*. 2017; 140(6):e20173035 and *Pediatrics*. 2018;142(3):e20181739]. *Pediatrics*. 2017;140(3):e20171904.
9. Whelton PK, Carey RM, Aronow WS, et al. 2017 ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2018;138(17):e426-e483.
10. US Preventive Services Task Force. Screening for obesity in children and adolescents: US Preventive Services Task Force recommendation statement. *JAMA*. 2017;317(23): 2417-2426.
11. US Preventive Services Task Force. Screening for lipid disorders in children and adolescents: US Preventive Services Task Force recommendation statement [published correction appears in *JAMA*. 2016;316(10):1116]. *JAMA*. 2016; 316(6):625-633. ■