

## ADHD Screeners and Quality of Life Assessments

### ADHD Screeners:

Screener	Details
<b>ADHD Rating Scale IV (ADHD-RS-IV) With Adult Prompts</b>	<ul style="list-style-type: none"> <li>• Rates the frequency and severity of ADHD symptoms</li> <li>• Consists of 18 items: 9 assess inattentive symptoms and 9 assess hyperactive-impulsive symptoms</li> <li>• Scoring is based on a 4-point frequency scale ranging from 0 (never) to 3 (very often)</li> <li>• To access, email: <a href="mailto:adultadhd2@nyulangone.org">adultadhd2@nyulangone.org</a></li> </ul>
<b>Adult ADHD Clinical Diagnostic Scale (ACDS) v1.2</b>	<ul style="list-style-type: none"> <li>• Provides information on current adult symptoms of ADHD</li> <li>• Based on a semi-structured clinical interview</li> <li>• Consists of 18 items</li> <li>• To access, email: <a href="mailto:adultADHD@med.nyu.edu">adultADHD@med.nyu.edu</a></li> </ul>
<b>Adult ADHD Investigator Rating Scale (AISRS)</b>	<ul style="list-style-type: none"> <li>• Based on a semi-structured clinical interview</li> <li>• Developed to evaluate symptoms at baseline and during treatment</li> <li>• Consists of 18 items: 9 for inattentive symptoms and 9 for hyperactive-impulsive symptoms</li> <li>• Scoring is based on 4-point scale from 0 (none) to 3 (severe)</li> <li>• Need an account to access; potential fee associated with this instrument</li> <li>• To access, email: <a href="mailto:adultadhd2@nyulangone.org">adultadhd2@nyulangone.org</a></li> </ul>
<b>Adult ADHD Self-Report Scale (ASRS) v1.1</b>	<ul style="list-style-type: none"> <li>• Developed as a 6-question subset of the ADHD ASRS Symptom Checklist</li> <li>• Used as an initial self-assessment tool</li> <li>• Consists of 6 items: 4 for inattentive symptoms and 2 for hyperactive-impulsive symptoms</li> <li>• Scoring is based on symptom frequency</li> <li>• <a href="https://www.hcp.med.harvard.edu/ncs/ftpd/ahd/6Q_ASRS_English.pdf">https://www.hcp.med.harvard.edu/ncs/ftpd/ahd/6Q_ASRS_English.pdf</a></li> </ul>
<b>Adult ADHD Self-Report Screening Scale for DSM-5 (ASRS DSM-5) Screener</b>	<ul style="list-style-type: none"> <li>• Developed as an update to ASRS v1.1</li> <li>• Used as a self-report</li> <li>• Consists of 6 items</li> <li>• Scoring is based on a 5-point scale from 0 (never) to 4 (very often)</li> <li>• <a href="https://www.hcp.med.harvard.edu/ncs/ftpd/ahd/ASRS-5_English.pdf">https://www.hcp.med.harvard.edu/ncs/ftpd/ahd/ASRS-5_English.pdf</a></li> </ul>
<b>Adult ASRS Symptom Checklist v1.1</b>	<ul style="list-style-type: none"> <li>• Used as an initial symptom assessment to identify adults who may have ADHD</li> <li>• Consists of 18 items</li> <li>• <a href="https://med.nyu.edu/psych/sites/default/files/psych/psych_adhd_checklist_0.pdf">https://med.nyu.edu/psych/sites/default/files/psych/psych_adhd_checklist_0.pdf</a></li> </ul>
<b>Barratt Impulsiveness Scale (BIS-11)</b>	<ul style="list-style-type: none"> <li>• Developed as a measure of impulsiveness</li> <li>• Used as a self-report</li> <li>• Consists of 30 items</li> <li>• Scoring is based on a 4-point scale from 1 (rarely/never) to 4 (almost always/always)</li> <li>• <a href="http://www.impulsivity.org/measurement/bis11">http://www.impulsivity.org/measurement/bis11</a></li> </ul>
<b>Clinical Global Impression (CGI)</b>	<ul style="list-style-type: none"> <li>• Consists of 3 items</li> <li>• Observer rated</li> <li>• One item measures illness severity, another measures global improvement or change, and last one measures therapeutic response</li> <li>• <a href="https://www.psywellness.com.sg/docs/CGI.pdf">https://www.psywellness.com.sg/docs/CGI.pdf</a></li> </ul>

**ADHD Screeners (cont.):**

<p><b>Brown Attention-Deficit Disorder Symptom Assessment Scale (BADDS) for Adults</b></p>	<ul style="list-style-type: none"> <li>• Used as a self-report or as part of a clinical interview</li> <li>• Consists of 40 items representing 5 symptom areas: activation, attention, effort, affect and memory</li> <li>• Asks about the adult’s clinical history, work and/or school functioning, leisure, mood, and sensitivity to criticism</li> <li>• Scoring is based on a 4-point frequency scale from 0 (never) to 3 (almost daily)</li> <li>• There is a fee associated with the use of this instrument</li> <li>• <a href="https://www.pearsonassessments.com/store/usassessments/en/Store/Professional-Assessments/Behavior/Attention-ADHD/Brown-Attention-Deficit-Disorder-Scales/p/100000456.html">https://www.pearsonassessments.com/store/usassessments/en/Store/Professional-Assessments/Behavior/Attention-ADHD/Brown-Attention-Deficit-Disorder-Scales/p/100000456.html</a></li> </ul>
<p><b>Conners’ Adult ADHD Rating Scales (CAARS)</b></p>	<ul style="list-style-type: none"> <li>• Measures the presence and severity of ADHD symptoms</li> <li>• Self-report and observer rating forms available</li> <li>• Consists of 26 items for the short form; 66 items for the long form; 30 items for the screening</li> <li>• There is a fee associated with the use of this instrument</li> <li>• <a href="https://storefront.mhs.com/collections/caars#:~:text=The%20Conners'%20Adult%20ADHD%20Rating,ad%20severity%20of%20ADHD%20symptoms">https://storefront.mhs.com/collections/caars#:~:text=The%20Conners'%20Adult%20ADHD%20Rating,ad%20severity%20of%20ADHD%20symptoms</a></li> </ul>
<p><b>Diagnostic Interview for ADHD in Adults (DIVA) 2.0</b></p>	<ul style="list-style-type: none"> <li>• Consists of three parts – attention deficit, hyperactivity-impulsivity, and age of onset and impairment</li> <li>• Based on a semi-structured clinical interview</li> <li>• Usually takes around 1 to 1.5 hours to complete</li> <li>• There is a potential fee associated with the use of this instrument (dependent on mechanism of delivery)</li> <li>• <a href="http://www.divacenter.eu/Content/VertalingPDFs/DIVA_2_EN_FORM%20-%20involbaar.pdf">http://www.divacenter.eu/Content/VertalingPDFs/DIVA_2_EN_FORM%20-%20involbaar.pdf</a></li> </ul>
<p><b>Wender Utah Rating Scale (WURS)</b></p>	<ul style="list-style-type: none"> <li>• Assess adults for ADHD with a subset of 25 questions</li> <li>• Full assessment consists of 61 questions about the adult’s childhood behavior</li> <li>• Scoring is based on a 5-point scale from 0 (not at all or very slightly) to 4 (very much)</li> <li>• <a href="http://neurosciencecmc.com/library/rating_scales/adhd_wender.pdf">http://neurosciencecmc.com/library/rating_scales/adhd_wender.pdf</a></li> </ul>

**Quality of Life Assessments:**

Assessment	Details
<p><b>Adult ADHD Quality of Life Measure (AAQoL) [2]</b></p>	<ul style="list-style-type: none"> <li>• Consists of 29 questions</li> <li>• Four subscale scores are derived: Productivity, Life Outlook, Relationships, and Health (psychological)</li> <li>• Scoring is based on a 5-point Likert-type scale for frequency of occurrence</li> <li>• Lower scores indicate greater impairment in functioning</li> <li>• <a href="http://www.dr.lancelevy.com/PDF-astools/ADHD-new-quality-of-life-questionnaire.pdf">http://www.dr.lancelevy.com/PDF-astools/ADHD-new-quality-of-life-questionnaire.pdf</a></li> </ul>
<p><b>Driving Behavior Survey (DBS) [4]</b></p>	<ul style="list-style-type: none"> <li>• 21-item scale indexing the frequency of anxious driving behavior</li> <li>• Across three domains: exaggerated safety/caution behavior, anxiety-based performance deficits, and hostile/aggressive behavior</li> <li>• Scoring is based on a 7-point scale from 1 (never) to 7 (always)</li> <li>• Higher scores indicate greater frequency of anxious behavior</li> <li>• <a href="https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3006470/table/T6/?report=objectonly">https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3006470/table/T6/?report=objectonly</a></li> </ul>
<p><b>Work Productivity and Activity Impairment Questionnaire General Health V2.0 (WPAI:GH)</b></p>	<ul style="list-style-type: none"> <li>• Consists of 6 items</li> <li>• Evaluates absenteeism (work time missed); presenteeism (impairment at work / reduced on-the-job effectiveness); work productivity loss (overall work impairment / absenteeism plus presenteeism); and activity impairment</li> <li>• <a href="http://www.reillyassociates.net/WPAI_General.html">http://www.reillyassociates.net/WPAI_General.html</a></li> <li>• <a href="http://oml.eular.org/sysModules/obxOml/docs/ID_98/WPAI-GH_English_US_V2.pdf">http://oml.eular.org/sysModules/obxOml/docs/ID_98/WPAI-GH_English_US_V2.pdf</a></li> </ul>